

Online behavioural measures are better than eye movements for predicting ADHD

Integrating objective measures of ADHD

Deborah Apthorp^{1,2, }
@deborahapthorp
dapthorp@une.edu.au

Kathleen Crocker¹
kcrocke2@myune.edu.au

¹ School of Psychology, University of New England, NSW, Australia
² School of Medicine and Psychology, Australian National University, ACT, Australia

Introduction

Diagnosis of Attention Deficit Hyperactivity Disorder (ADHD) is currently quite fragmented and ad hoc (Planton et al. 2021). Diagnostic criteria rely on either self-report, parent-report or both; the main validated survey for adults is the Adult ADHD Self-Report Scale (ASRS; Kessler et al. 2005). There is increasing interest in more objective measures to increase precision and reliability in diagnosis; since ADHD is construed as a disorder which affects attention, behavioural tasks which are designed to measure various aspects of attention should be applicable. The Continuous Performance Test (CPT) is commonly used (Epstein et al. 1998). In addition, eye tracking measures are a promising line of enquiry and have shown differences in lab-based studies between individuals with ADHD and controls (Conrad et al. 2023). These have the potential to show early warning signs even in babies too young to perform behavioural tasks. We investigated the potential to use online webcam-based eye tracking and behavioural tasks for remote diagnosis. This study was preregistered at the OSF: osf.io/ar6k2/

Objectives

1. Recruit online participants with and without diagnosed ADHD
2. Collect survey data (ASRS + demographics)
3. Online eye tracking (RealEye)
4. Online behavioural data collection (Continuous Performance Task - Non-X)

Methods

Participants

Table 1: Descriptive statistics for age and gender by ADHD diagnosis

	No	Yes	p
n	84	35	
age (mean (SD))	35.04 (11.22)	36.06 (12.45)	0.662
gender (%)			0.168
Male	14 (16.7)	10 (28.6)	
Female	66 (78.6)	25 (71.4)	
Other	4 (4.8)	0 (0.0)	

- 118 completed ASRS (Age: 18-60, M = 36; 24 male)
- 58 also completed eye tracking and CPT
- Dropout may have been caused by webcam and browser restrictions

Measures

- Adult ADHD self-report scale (ASRS- ref)
- CPT non-X (Pavlov/Py) - errors of omission, errors of commission, d', mean RT
- Eye tracking (RealEye) - fixation duration, fixation count, scanpath length

Procedure

- Qualtrics survey (information, consent, demographics, ASRS)
- RealEye online eye tracking - while doing practice task for CPT non-X (ask me why)
- CPT non-X - Pavlov/Py - see Figure 1




Figure 1: An illustration of the CPT non-X paradigm

Results

Participants who had been diagnosed with ADHD were higher on the ASRS (total and subscales), although there was some overlap - see Figure 2.




Figure 2: ASRS part A (screening section) and subscales for diagnosed and non-diagnosed participants

Because so few participants diagnosed with ADHD completed the eye tracking and CPT non-X, and since the study attracted the interest of many who suspected they might have ADHD (reflected in the qualitative data and the proportion who scored above the clinical cutoff for Section A), we thought it would be interesting to look at correlations between the ASRS subscales and the behavioural and eye tracking data (also preregistered). There were strong correlations for the behavioural data, but not for any of the eye tracking measures - see Figure 3




Figure 3: Correlations between ASRS measures and behavioural (top) and eye tracking measures (bottom)

Due to the low number of participants diagnosed with ADHD who completed the CPT non-X (11) and eye tracking (12), we diverged from the preregistration for the final analysis (hierarchical logistic regression) by using the ASRS clinical cutoff to assign non-diagnosed individuals to the ADHD group based on their scores. This resulted in groups of 31 probable/diagnosed ADHD participants and 27 controls.

A hierarchical logistic regression showed that, after controlling for age and gender, CPT variables errors of commission, errors of omission and reaction times predicted group membership, but no eye tracking variables were significant in the model. Classification accuracy was 85%, with 90% sensitivity and 78% specificity, AUC = 0.93. See Table 2.

Table 2: Odds ratio table for the logistic regression fit

term	estimate	std.error	statistic	pvalue
age	9.420000e-01	0.047	-1.280	0.200
gender	1.549000e+00	0.755	0.579	0.562
errors_omission	0.000000e+00	16.135	-2.362	0.018
errors_commission	6.171004e+06	4.903	3.189	0.001
r_ts	6.424268e+15	15.392	2.365	0.018
log_scanpath_length	2.396200e+01	4.705	0.675	0.500
fixation_duration_ms	9.900000e-01	0.020	-0.509	0.611
fixation_count	9.860000e-01	0.027	-0.517	0.605

Discussion

- CPT non-X is feasible to use online as an ADHD diagnostic tool
- Online eye tracking is not there yet (at least not RealEye)
- Tracking for a longer time across entire task would have been better (but calibration may have been an issue?)
- Integrating more objective measures in ADHD diagnosis is feasible, even in an online environment
- Could be useful for rural/remote diagnosis where clinical services are limited

References

Conrad, Alexis, Jason Brunt, and John Williams. 2023. "B - 03 Inhibitory Control and Saccadic Eye Movement in Adults with Attention-Deficit/Hyperactivity Disorder: An Eye-Tracking Analysis." *Archives of Clinical Neuropsychology* 38 (7): 1367-67. <https://doi.org/10.1093/arclin/acad072.0>.
Epstein, Jeffery N., C. Keith Conners, Gill Sitarenios, and Drew Erhardt. 1998. "Continuous Performance Test Results of Adults with Attention Deficit Hyperactivity Disorder." *The Clinical Neuropsychologist* 12 (2): 155-68. <https://doi.org/10.1076/clin.12.2.155.2000>.
Kessler, Ronald C., Lenard Adler, Minnie Ames, et al. 2005. "The World Health Organization Adult ADHD Self-Report Scale (ASRS): A Short Screening Scale for Use in the General Population." *Psychological Medicine* 35 (2): 245-56. <https://doi.org/10.1017/S0033291704002892>.
Planton, M., B. Lemesle, M. Cousineau, et al. 2021. "The Role of Neuropsychological Assessment in Adults with Attention Deficit/Hyperactivity Disorders." *Revue Neurologique* 177 (4): 341-48. <https://doi.org/10.1016/j.neurol.2021.01.006>.

