

Bifurcations in continuous time piecewise smooth dynamical systems

Debsankha Manik

December 10, 2012

A BRIEF SUMMARY ON PWS FLOWS

A simple piecewise smooth function:

$$\dot{x} = \begin{cases} F_1(x) & : H(x) < 0 \\ F_2(x) & : H(x) > 0 \end{cases}$$


Switching manifold: $H(x) = 0$

The flows of F_1 and F_2 are $\varphi_1(x)$ and $\varphi_2(x)$ respectively, defined in respective regions and also in the neighbourhood of the switching manifold:

$$\dot{\varphi}_1 = F_1 \tag{1}$$

$$\dot{\varphi}_2 = F_2 \tag{2}$$

Figure : Example: Impact Oscillator



HOW TO ANALYZE NONLINEAR SYSTEMS?

Figure : Look at the vector field






Figure : Possible scenarios



Bifurcation of equilibria

Grazing bifurcation



Sliding bifurcation

Choose a coordinate system such that:

$$\dot{\mathbf{x}} = \begin{cases} F_1(\mathbf{x}, \mu) & : x_n < 0 \\ F_2(\mathbf{x}, \mu) & : x_n > 0 \end{cases}$$

and $\mathbf{x} = \mathbf{0}$ is a grazing point.

x_n == n th component of \mathbf{x} .

Locally linearize:

$$\dot{\mathbf{x}} = \begin{cases} \mathbf{A}_1 \mathbf{x} + \mathbf{B} \mu & : x_n < 0 \\ \mathbf{A}_2 \mathbf{x} + \mathbf{B} \mu & : x_n > 0 \end{cases}$$

Where:

$$\mathbf{A}_i = \left. \frac{\partial \mathbf{F}_i}{\partial \mathbf{x}} \right|_{\mathbf{x}=\mathbf{0}}$$

and $\mathbf{B} = \left. \frac{\partial \mathbf{F}_1}{\partial \mu} \right|_{\mu=0} = \left. \frac{\partial \mathbf{F}_2}{\partial \mu} \right|_{\mu=0}$ (Due to continuity).

Also, \mathbf{A}_1 and \mathbf{A}_2 can differ only in the n -th column (Again due to continuity).

Let: $A_1 \mathbf{x}_1^* + \mathbf{B}\mu = \mathbf{0}$, $A_2 \mathbf{x}_2^* + \mathbf{B}\mu = \mathbf{0}$.

Assuming A_i 's are invertible:

$$\mathbf{x}_i^* = -\mathbf{A}_i^{-1} \mathbf{B}\mu = -\frac{\text{adj}(\mathbf{A}_i)}{\det(\mathbf{A}_i)} \mathbf{B}\mu$$

The solutions exist iff:

$$x_{1_n < 0}^* < 0, x_{2_n}^* > 0.$$

Now,

$$x_{1_k}^* = \frac{c_{1_k}^*}{\det(\mathbf{A}_1)} \mu, x_{2_k}^* = \frac{c_{2_k}^*}{\det(\mathbf{A}_2)}$$

Where,

$$c_{i_k}^* = [-\text{adj}(\mathbf{A}_i) \mathbf{B}]_k = [-\text{adj}(\mathbf{A}_i)_{kj} \mathbf{B}_j]$$


Because A_1 differs from A_2 only in n -th column, $\text{adj}(A_1)$ and $\text{adj}(A_2)$ shares a common n -th row, $c_{1_n}^* = c_{2_n}^* := C$

CONDITION FOR BORDER CROSSING OF EQUILLIBRIA

$$x_{1_n}^* = \frac{C}{\det(\mathbf{A}_1)}\mu, x_{2_k}^* = \frac{C}{\det(\mathbf{A}_2)}\mu$$

Cases:

1. $\det(\mathbf{A}_1)\det(\mathbf{A}_2) < 0$. $x_{1_n}^*$ and $x_{2_n}^*$ always have opposite signs.
2. $\det(\mathbf{A}_1)\det(\mathbf{A}_2) > 0$. $x_{1_n}^*$ and $x_{2_n}^*$ always have same signs.




Case 1

Case 2

GRAZING ORBITS

Figure :



GRAZING ORBITS


Case 1: $F_1(x) \neq F_2(x)$ at switching manifold: The Poincare map (given by Nordmak et al[4], using the ZDM formalism; also by Molenaar, without ZDM) is of the form:

$$\left. \begin{aligned} x_{n+1} &= ax_n + y_n + \rho \\ y_{n+1} &= -bx_n \end{aligned} \right\} \quad x \leq 0$$

$$\left. \begin{aligned} x_{n+1} &= -c\sqrt{x_n} + y_n + \rho \\ y_{n+1} &= -dx_n \end{aligned} \right\} \quad x > 0$$

- ▶ Jacobian of the system is singular.
- ▶ Infinite stretching of the phase space.
- ▶ In some cases, period-adding bifurcation occurs.
- ▶ This singularity affects only the trace of J , not the determinant. [3]

Figure : Bifurcation in a map with square root singularity




Case 2: $F_1(x) = F_2(x)$ at switching manifold: The Poincare map (given by Dankowicz and Nordmak) is of the form:

$$x_{n+1} = \begin{cases} ax_n + \rho & x \leq 0 \\ ax_n - bx_n^{3/2} + \rho & x > 0 \end{cases}$$

- ▶ Jacobian of the system is continuous.
- ▶ No chance of border collision bifurcation (Elaborate later).
- ▶ Both trace and determinant of J vary continuously.

DISCONTINUOUS JACOBIAN IS NECESSARY FOR PERIOD DOUBLING ON BC

Figure :



Suppose for the parameter value $\mu = 0$, the fixed point of the left hand side map crosses the boundary $x = 0$. Suppose a period doubling occurs.

Now the fixed point of both the maps are at δ^2 , say. Then:

$$f_2(f_1(-\epsilon^2)) = -\epsilon^2$$

$$f_2(\delta^2 + \dot{f}_1(\delta^2)(\delta^2 + \epsilon^2)) = -\epsilon^2$$

$$\delta^2 + \dot{f}_2(\delta^2)\dot{f}_1(\delta^2)(\delta^2 + \epsilon^2) = -\epsilon^2$$

$$\dot{f}_2(\delta^2)\dot{f}_1(\delta^2) = -1$$

$$\dot{f}_2(0)\dot{f}_1(0) \approx -1$$

Therefore the discontinuity is evident.

HOW TO UNDERSTAND A BORDER COLLISION BIFURCATION

What do we do in smooth bifurcations?

1. Take Poincare section.
2. Calculate fixed points.
3. Calculate jacobian and its eigenvalues. [If the absolute values of the eigenvalues are < 1 , we will see attractive behaviour, otherwise repulsive.]

What's the problem now?


-There's no easy way to get the Poincare map in a closed form.

However, we can still compute eigenvalues by searching for a direction which remains unchanged in each stroboscopic slice.

-But we need to find out the fixed points first.

A BORDER CROSSING ORBIT

Figure :



Supposing the orbit crosses the border only once.
Then finding out the orbit boils down to:

$$x_1 = \varphi_1(x_0, 0, \tau_0) \quad (3)$$

$$H(x_1) = 0 \quad (4)$$

$$x_2 = \varphi_2(x_1, \tau_0, \tau_1) \quad (5)$$

$$H(x_2) = 0 \quad (6)$$

$$x_0 = \varphi_1(x_2, \tau_1, T - \tau_0 - \tau_1) \quad (7)$$

This is a set of $3n + 2$ equations in $3n + 2$ unknowns, so this can be tackled with standard Newton's method of root finding:[3]

$$y_{n+1} = \frac{G(y_n)}{J(\bar{y}_n)}$$

Here $y := \{x_0, x_1, x_2, \tau_0, \tau_1\}$.

THEORY OF BORDER COLLISION BIFURCATIONS

For a PWS system, we can now:

1. Find out all the periodic orbits. But finding out a T_j orbit involves inverting a $(2j + 1)n + 2 * j$ dimensional matrix. Since best known matrix inversion algorithms are $O(n^2)$, it may not be feasible to do this for orbits that cross the border many times.

THEORY OF BORDER COLLISION BIFURCATIONS

For a PWS system, we can now:

1. Find out all the periodic orbits. But finding out a T_j orbit involves inverting a $(2j + 1)n + 2 * j$ dimensional matrix. Since best known matrix inversion algorithms are $O(n^2)$, it may not be feasible to do this for orbits that cross the border many times.
2. Then we can find out the eigenvalues of the Poincare map at the fixed points by taking small deviations from the periodic orbit and doing a least-square style fitting for the parameters for the locally linear map. (Only δ and τ in case of 2-d) Interesting phenomena will occur when fixed points change nature after border collision.

THEORY OF BORDER COLLISION BIFURCATIONS

For a PWS system, we can now:

1. Find out all the periodic orbits. But finding out a T_j orbit involves inverting a $(2j + 1)n + 2 * j$ dimensional matrix. Since best known matrix inversion algorithms are $O(n^2)$, it may not be feasible to do this for orbits that cross the border many times.
2. Then we can find out the eigenvalues of the Poincare map at the fixed points by taking small deviations from the periodic orbit and doing a least-square style fitting for the parameters for the locally linear map. (Only δ and τ in case of 2-d) Interesting phenomena will occur when fixed points change nature after border collision.
3. Then we apply standard results regarding the stability of fixed learnt in case of smooth bifurcations to explain the phenomenon.

The locally linear map will be, in 2-D:

$$\begin{bmatrix} x \\ y \end{bmatrix} \rightarrow \begin{cases} \begin{bmatrix} \tau_L & 1 \\ -\delta_L & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \mu \begin{bmatrix} 1 \\ 0 \end{bmatrix} & x \leq 0 \\ \begin{bmatrix} \tau_R & 1 \\ -\delta_R & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \mu \begin{bmatrix} 1 \\ 0 \end{bmatrix} & x > 0 \end{cases}$$

The parameter space of $\delta_L, \delta_R, \tau_L, \tau_R$ can be divided into regions exhibiting different behaviours on border collision. [1]

- [1] Soumitro Banerjee, Priya Ranjan, and Celso Grebogi. Bifurcations in two-dimensional piecewise smooth maps - theory and applications in switching circuits, 2000.
- [2] M. Di Bernardo, F. Garofalo, L. Iannelli, and F. Vasca. Bifurcations in piecewise-smooth feedback systems. *International Journal of Control*, 75(16-17):1243–1259, 2002.
- [3] Yue Ma, James Ing, Soumitro Banerjee, Marian Wiercigroch, and Ekaterina Pavlovskaja. The nature of the normal form map for soft impacting systems. *International Journal of Non-Linear Mechanics*, 43(6):504 – 513, 2008.
 ;ce:title;Non-linear Dynamics of Engineering Systems;/ce:title;.
- [4] A.B. Nordmark. Non-periodic motion caused by grazing incidence in an impact oscillator. *Journal of Sound and Vibration*, 145(2):279 – 297, 1991.