

Assignment 8: Complex Numbers 1

Declan Murphy Zink

10/5/2020

3

3.a Find the real and imaginary parts of $(\sqrt{3} - i)^{10}$ and $(\sqrt{3} - i)^{-7}$. For which values of n is $(\sqrt{3} - i)^n$ real?

$$(\sqrt{3} - i)^{10} = (2e^{-\frac{\pi}{6}i})^{10} = 2^{10}e^{-\frac{10\pi}{6}i} = 2^9 \cdot 2e^{\frac{\pi}{3}i} = 2^9(1 + \sqrt{3}i).$$

$$(\sqrt{3} - i)^{-7} = 2^{-7}e^{\frac{7\pi}{6}i} = 2^{-8}(-\sqrt{3} - i).$$

for $(\sqrt{3} - i)^n$ to be real, $\frac{n\pi}{6} = k\pi, k \in \mathbb{Z}$, so n must be a multiple of 6.

3.b What is \sqrt{i}

$$\sqrt{i} = (e^{\frac{\pi}{2}i})^{\frac{1}{2}} = e^{\frac{\pi}{4}i} \text{ or } e^{\frac{5\pi}{4}i}$$

3.c Find all tenth roots of i . Which one is nearest to i in the Argand diagram?

$$\sqrt[10]{i} = (e^{\frac{\pi}{2}i})^{\frac{1}{10}} = e^{\frac{\pi}{20}i}, e^{\frac{5\pi}{20}i}, e^{\frac{9\pi}{20}i}, e^{\frac{13\pi}{20}i}, e^{\frac{17\pi}{20}i}, e^{\frac{21\pi}{20}i}, e^{\frac{25\pi}{20}i}, e^{\frac{29\pi}{20}i}, e^{\frac{33\pi}{20}i}, e^{\frac{37\pi}{20}i}.$$

$e^{\frac{9\pi}{20}i}$ is closest to i .

3.d Find the seven roots of the equation $z^7 - \sqrt{3} + i = 0$. Which one of these roots is closest to the imaginary axis?

$$z^7 = \sqrt{3} - i, \text{ so for } k \in \mathbb{Z}, 0 \leq k < 7, z = (2e^{(-\frac{\pi}{6} + 2k\pi)i})^{\frac{1}{7}} = 2^{\frac{1}{7}}e^{(-\frac{\pi}{42} + \frac{12k\pi}{42})i}$$

Closest to the imaginary axis $= 2^{\frac{1}{7}}e^{\frac{23\pi}{42}i}$.

4 Prove the "Triangle Inequality" for complex numbers: $|u + v| \leq |u| + |v|$ for all $u, v \in \mathbb{C}$.

For $a, b, c, d \in \mathbb{R}$:

$$|a + bi + c + di| \leq |a + bi| + |c + di|$$

$$\sqrt{(a + c)^2 + (b + d)^2} \leq \sqrt{a^2 + b^2} + \sqrt{c^2 + d^2}$$

$$(a + c)^2 + (b + d)^2 \leq a^2 + b^2 + c^2 + d^2 + 2\sqrt{(a^2 + b^2)(c^2 + d^2)}$$

$$\begin{aligned}
2ac + 2bd &\leq 2\sqrt{(a^2 + b^2)(c^2 + d^2)} \\
ac + bd &\leq \sqrt{a^2c^2 + a^2d^2 + b^2c^2 + b^2d^2} \\
a^2c^2 + 2abcd + b^2d^2 &\leq a^2c^2 + a^2d^2 + b^2c^2 + b^2d^2 \\
0 &\leq a^2d^2 - 2abcd + b^2c^2 \\
0 &\leq (ad - bc)^2
\end{aligned}$$

This must be true since $a, b, c, d \in \mathbb{R}$.

5 Let z be a non-zero complex number. Prove that the three cube roots of z are the corners of an equilateral triangle in the Argand diagram.

Let $z = re^{(\theta+2k\pi)i}$ for $k \in \mathbb{Z}, 0 \leq k < 3$. Thus $\sqrt[3]{z} = r^{\frac{1}{3}}e^{(\frac{\theta}{3}+\frac{2k\pi}{3})i}$, which means that each cubic root is $\frac{2\pi}{3}$ radians from the other two roots. Since the modulus of each root is the same, $r^{\frac{1}{3}}$, this forms an equilateral triangle.