Course code and name:	B38EM Introduction to Electricity and Magnetism	
Type of assessment:	Individual	
Coursework Title:	Take home Assignment 2	
Student Name:	MA XUNCHI	
Student ID Number:	H00392669 21012100015	

Declaration of authorship. By signing this form:

- I declare that the work I have submitted for individual assessment OR the work I have contributed to a group assessment, is entirely my own. I have NOT taken the ideas, writings or inventions of another person and used these as if they were my own. My submission or my contribution to a group submission is expressed in my own words. Any uses made within this work of the ideas, writings or inventions of others, or of any existing sources of information (books, journals, websites, etc.) are properly acknowledged and listed in the references and/or acknowledgements section.
- I confirm that I have read, understood and followed the University's Regulations on plagiarism as published on the <u>University's website</u>, and that I am aware of the penalties that I will face should I not adhere to the University Regulations.
- I confirm that I have read, understood and avoided the different types of plagiarism explained in the University guidance on <u>Academic Integrity and Plagiarism</u>

Student Signature: MA XUNCH1

Date: 04 -20-2023


Copy this page and insert it into your coursework file in front of your title page. For group assessment each group member must sign a separate form and all forms must be included with the group submission.

Your work will not be marked if a signed copy of this form is not included with your submission.

Introduction to Electricity and Magnetism B38EM Assignment #2

$$\varepsilon_0 = 8.85 \times 10^{-12} \,\text{Fm}^{-1}$$
, $e = 1.6 \times 10^{-19} \,\text{C}$, $\mu_0 = 4 \pi \, 10^{-7} \,\text{N/A}^2$

1. Consider the system shown in **Figure Q1** where a metal bar of mass *m* slides frictionless on two parallel conducting rails separated by a distance *l*. A resistor *R* is connected across the rails. If a uniform magnetic field B, pointing into the page, fills the entire region, find the following:



- (a) If the bar moves to the right at speed v, what is the current in the resistor? In which direction does it flow? (3 marks)
- (b) Determine the magnetic force. What is the direction of the force? (3 marks)
- (c) If the bar starts out with speed v_0 at t=0, and it is left to slide, what is its speed at a later t? (4 marks)

(3 marks)

(3 marks)

(d)
$$V = \int_{0}^{\infty} \left(\frac{B_{1}V}{R}\right)^{2} R dt$$

Since $V = V_{0}e^{-\frac{B^{2}RV}{Rm}t}$

(b) $F = BI = \frac{B^{2} I^{2}V}{R}$

The current in the conductor goes upwards, So the direction of magnetic force is left

(C) $F = \frac{B^{2}I^{2}V}{R}$

Proved

$$M \frac{dV}{dt} = -\frac{B^{2}I^{2}V}{Rm} dt \implies V = V_{0}e^{\frac{B^{2}R^{2}V}{Rm}t}$$

- 2. The transformer shown in Figure Q2 consists of a long wire coincident with the z-axis carrying a current $I = I_0 \cos \omega t$, coupling magnetic energy to a toroidal coil situated in the x-y plane and centred at the origin. The toroidal core uses iron material with relative permeability, μ_r , around which 100 turns of a tightly wound coil serves to induce a voltage V_{emf} , as shown in the figure.
 - (a) Develop an expression for V_{emf} (3 marks)
 - (b) Calculate V_{emf} , for the parameters in **Table Q2** (2 marks)

$$\begin{array}{l}
(a) \\
E = \oint \overrightarrow{B} \cdot dS \\
= \int_{a}^{b} \frac{\mu I}{2\pi \gamma} \cdot C d\gamma \\
= \frac{\mu c I}{2\pi} \ln(\frac{b}{a})
\end{array}$$

2.




Figure Q2

$=\frac{\mu_{CNWI_0}}{2\pi}\left(h(\frac{b}{a})\sin\frac{b}{a}\right)$	۷.
--	----

Table Q2		
Parameter	Value	
Frequency, f	60 Hz	
Relative permeability, μ_r	4000	
a	5 cm	
ь	6 cm	
С	2 cm	
Current amplitude, I_0	50 A	

(b) according to a

Vemf ~5.5 Sin 377t

- 3. In a Cartesian coordinate, a plane wave is polarized with its electric vector along z. The wave propagates along the y-axis. The electric field is given by $E_z(y,t) = E_0 e^{i(ky-\omega t)}$ Volts/metre. This wave is propagating in vacuum; its amplitude is $E_0 = 7$ V/m and its wavelength is $\lambda = 0.3$ meters.
- a) What is the frequency of the wave? (2 marks)
- b) How large is the magnetic field associated with this wave (2 marks) and in what direction is it oriented (2 marks)?
- c) What is the average rate at which energy is transported by this wave (per square metres)?

$$f = \frac{C}{\lambda} = \frac{3 \times 10^8 \, \text{m/s}}{0.3 \, \text{m}}$$

$$= 10^9 \, \text{Hz}$$

$$(C) \, I = \frac{1}{2} \, C \, \epsilon_0 \, E_0^2 \approx 0.065 \, \text{W/m}^3$$

- 4. A coaxial cable that connects a radar receiver to its antenna is 30-m-long and it is lossless. The cable has a characteristic impedance $Z_0 = 50 \Omega$ and operates at 2 MHz. The cable is terminated at an unmatched antenna that has an impedance of $Z_L = 60 + j40 \Omega$. If the phase velocity on this transmission line is u = 0.6c (where c is the speed of light in vacuum), find:
- d) The complex reflection coefficient Γ at the cable and antenna interface; (2 marks)
- e) The voltage standing wave ratio; (2 marks)
- f) The input impedance of this transmission line. (1 marks)

$$U_{11}d_{1}T = \frac{2c-20}{2c+20} = \frac{10tj40}{110tj40} \approx 0.352 255.98^{\circ}$$

(e)
$$S = \frac{[+|T|]}{[-|T|]} = \frac{[+0.352]}{[-0.352]} \approx 2.086$$

$$\beta = \frac{2\pi f}{u}$$
(f) $2 = \frac{2\pi f}{20 + j 2 \tan \beta l} \sim (23.973 + j)(352) 52$