



Signature and Log-signature for the Study of Empirical Distributions Generated with GANs.


J. de Curtò y DÍAz.

I. de Zarzà y Cubero.


29-03-2022.

c@decurto.be

z@dezarza.be



Signature and Log-signature for the Study of Empirical Distributions Generated with GANs.



De Curtò y DíAz.
De Zarzà y Cubero.

Signature and Log-signature for the Study of Empirical Distributions Generated with GANs

<https://arxiv.org/abs/2203.03226>

<https://github.com/decurtoydiaz/signatures>

Acknowledgements

This work is part of CIMDA (Centre for Intelligent Multidimensional Data Analysis), HK Science Park, HK.

A joint Center between City University of Hong Kong and the University of Oxford.

Our work has been supported by HK Innovation and Technology Commission (InnoHK Project CIMDA) and HK Research Grants Council (Project CityU 11204821).

c@decurto.be

z@dezarza.be





De Curtò y DíAz.
De Zarzà y Cubero.

Signature and Log-signature for the Study of Empirical Distributions Generated with GANs.

Key contributions:

- We propose two frameworks to assess GAN convergence based on analytical measures, one based on statistical tests and the other leveraging tools from harmonic analysis (a recently proposed generalization of Fourier).

Since its first appearance in 2014, measures to assess GAN convergence have been mainly based on empirical evaluations (MS-SSIM and FID).

- We also propose a PCA-Adaptive t-SNE for image visualization on several domains.

Goodfellow, I., Pouget-Abadie, J., Mirza, M., Xu, B., Warde-Farley, D., Ozair, S., Courville, A., Bengio, Y.: Generative adversarial networks. NIPS **27** (2014)

van der Maaten, L., Hinton, G.: Visualizing data using t-SNE. JMLR **9**(86), 2579–2605 (2008)

c@decurto.be

z@dezarza.be



Signature and Log-signature for the Study of Empirical Distributions Generated with GANs.



De Curtò y DíAz.
De Zarzà y Cubero.

Visualization pipeline.






Fig. 2: K-means Clustering (left) and t-SNE (right) on images from NASA Curiosity (upper figures) and Perseverance (lower figures).

c@decurto.be
z@dezarza.be



Signature and Log-signature for the Study of Empirical Distributions Generated with GANs.



De Curtò y Díaz.
De Zarzà y Cubero.

Table 4: Evaluation of the statistical test measures of homoscedasticity (T1), normality (T2) and goodness of fit (T3) on AFHQ and MetFaces using state-of-the-art pretrained models of Stylegan2-ada [21] and Stylegan3-ada [22] and NASA Perseverance.

Model	Dataset	T1	T2	T3
Stylegan2-ada	NASA Perseverance	✗	✓	✗
	AFHQ	Cat ✗	✗	✓
		Dog ✗	✓	✗
		Wild ✗	✗	✓
r-Stylegan3-ada	MetFaces	✗	✗	✗
t-Stylegan3-ada		✗	✗	✗

c@decurto.be
z@dezarza.be



Signature and Log-signature for the Study of Empirical Distributions Generated with GANs.




De Curtò y DíAz.
De Zarzà y Cubero.

Following [2], the truncated signature of order N of the path \mathbf{x} is defined as a collection of coordinate iterated integrals

$$S^N(\mathbf{x}) = \left(\left(\int \cdots \int \prod_{c=1}^a \frac{df_{z_c}}{dt}(t_c) dt_1 \cdots dt_a \right)_{1 \leq z_1, \dots, z_a \leq d} \right)_{1 \leq a \leq N}. \quad (1)$$

Bonnier, P., Kidger, P., Arribas, I.P., Salvi, C., Lyons, T.: Deep signature transforms. NIPS (2019)

Lyons, T.: Rough paths, signatures and the modelling of functions on streams. Proceedings of the International Congress of Mathematicians (2014)



c@decurto.be
z@dezarza.be

Model	Dataset	RMSE \tilde{S}^3	MAE \tilde{S}^3	RMSE $\log \tilde{S}^3$	MAE $\log \tilde{S}^3$
Stylegan2-ada	Cat	61450	45968	29201	22297
	Dog	38861	30441	31686	24612
	Wild	33306	25578	26622	20359
r-Stylegan3-ada	MetFaces	33247	23428	25685	18071
t-Stylegan3-ada		34977	22799	24707	16539
		30894	19872	21560	13761



Signature and Log-signature for the Study of Empirical Distributions Generated with GANs.



De Curtò y DíAz.
De Zarzà y Cubero.

Definition 4. Given n components of the element-wise mean of the signatures $\{y^{(c)}\}_{c=1}^n \subseteq T(\mathbb{R}^d)$ from the model chosen as a source of synthetic samples and the same number of components of the element-wise mean of the signatures $\{x^{(c)}\}_{c=1}^n \subseteq T(\mathbb{R}^d)$ from the original distribution, then we define the Root Mean Squared Error (RMSE) and Mean Absolute Error (MAE) by

$$\text{RMSE} \left(\left\{ x^{(c)} \right\}_{c=1}^n, \left\{ y^{(c)} \right\}_{c=1}^n \right) = \sqrt{\frac{1}{n} \sum_{c=1}^n (y^{(c)} - x^{(c)})^2},$$

and

$$\text{MAE} \left(\left\{ x^{(c)} \right\}_{c=1}^n, \left\{ y^{(c)} \right\}_{c=1}^n \right) = \frac{1}{n} \sum_{c=1}^n |y^{(c)} - x^{(c)}|.$$

The case for log-signature is analogous.

J. de Curtò, I. de Zarzà, and H. Yan, “Signature and Log-signature for the Study of Empirical Distributions Generated with GANs,” *arXiv:2203.03226*, 2022.

c@decurto.be

z@dezarza.be





Fig. 8: Spectrum of the element-wise mean of the signatures (left) and log-signatures (right) of order 3 and size 64×64 of original ('o') against synthetic ('x') samples.



Signature and Log-signature for the Study of Empirical Distributions Generated with GANs.





Fig. 13: Visualization of PCA Adaptive t-SNE on original (left) versus synthetic (right) samples of AFHQ Cat (a,b), Dog (c,d) and Wild (e,f) using Stylegan2-ada.



De Curtò y Díaz.
De Zarzà y Cubero.




Fig. 15: Visualization of PCA Adaptive t-SNE on original (top) versus synthetic (bottom) samples of NASA Perseverance using Stylegan2-ada across several epoch iterations: 193 (b), 371 (c), 596 (d), 798 (e) and 983 (d).

c@decurto.be
z@dezarza.be

