

IMO 2026 — Problem 3

Solution by Kimi K3

2026-07-18

Source: `results/kimi-k3/problem-03/current.md` — autonomous run, no human intervention; solution as written by the model. Multi-round runs: earlier-round audit logs in `scratch/`.

Problem

Let n be a positive integer. Liu Bang and Xiang Yu have a stick of length 1 and want to divide it between themselves. Liu Bang marks at most n points on the stick, and then Xiang Yu marks at most n points on the stick. The marked points are distinct. Then, the stick is cut at all marked points, creating a number of pieces. Afterwards, they take turns claiming any unclaimed piece of the stick, with Liu Bang going first. Each player's goal is to maximise the total length of their own pieces. For each n , determine the largest value c such that Liu Bang may guarantee a total length of at least c , regardless of Xiang Yu's play.

Solution document

Status

solved

Approaches tried

- Modeling: claiming game with “pick any piece” — optimal play = greedy (proved below, brute-force verified). Game reduces to: LB picks a composition of 1 into $\leq n + 1$ pieces, XY subdivides with $\leq n$ cuts; payoff D = alternating sum; LB share = $(1 + D)/2$.
- Grid computation: $c_1 = 2/3, c_2 = 4/7, c_3 = 8/15$ conjecture $c_n = \frac{2^n}{2^{n+1}-1}$, LB plays geometric pieces $\frac{2^{i-1}}{2^{n+1}-1}$. (Confirmed as the answer.)
- Odd-set identity: $D(M) = |\{t : N(t) \text{ odd}\}|$ (layer cake). Basis of the matching formula.
- **Matching formula** (proved below; verified `code/tests2.py`, 300 tests): $D(M) = \min_{\mu} (\sum_{\{x,y\} \in \mu} |x - y| + \sum_{z \text{ unmatched}} z)$.
- **Part A** (LB guarantee): super-increasing family (margin δ , m pieces), subdivided with $\leq m - 1$ cuts $D \geq \delta$. Proof: matching formula + loop-augmented component graph + counting ($e \leq m - 1$ tree component exists) + tree inequality ($|\sum_{i \in C} \epsilon_i a_i| \leq \text{cost}(C)$) + margin. Verified 1500 random trials. (Fix vs. earlier sketch: self-pairs count as *loops*, making their component cyclic; with this, the tree inequality is per-component correct.)

- NM (“some same-parent merge never increases D ”) — FALSE in general ($\{1,3\}$, cut $3 \rightarrow 1.5+1.5$: D drops $2 \rightarrow 1$, only merge raises it). Induction-via-merging abandoned; matching route used instead.
- **Part B** (XY counterstrategy): parity view of cuts: bisect x (=remove x), cancel $x > y$ (=remove x, y , add $x-y$), 1 cut each. Chains = nested abs values on a subset. Chain theorem: any m positive reals with sum S have a nonempty subset with chain value $\leq S/(2^m - 1)$; proved by pigeonhole on 2^m subset sums + **greedy-scheduling realizability lemma** (adjacent gaps are chain-realizable). Verified $n=1..4$ (2000 random each) + adversarial local search (extremum = geometric family, value exactly $T_n S$); AGR checked 180 families, 0 failures.
- Dominance “min-subdiv = min-halve/cancel”: false on coarse grids (grid artifacts only); irrelevant — for Part B we only need halve/cancel sequences, which are valid subdivisions.

Current best

Answer: $c = \frac{2^n}{2^{n+1}-1}$. Proved in full below. Lower bound (LB guarantee): LB cuts into geometric pieces $\frac{2^{i-1}}{2^{n+1}-1}$ ($i = 1, \dots, n+1$); any refinement by $\leq n$ further cuts has alternating sum $D \geq \frac{1}{2^{n+1}-1}$ (Theorem 3.1). Upper bound (XY counterstrategy): for any LB composition, XY either bisects all pieces (if fewer than $n+1$) forcing $D = 0$, or uses a sub- T_n chain (Theorem 4.2, via pigeonhole + realizability) to force $D \leq \frac{1}{2^{n+1}-1}$. Hence LB gets exactly $\frac{1}{2}(1 + \frac{1}{2^{n+1}-1}) = \frac{2^n}{2^{n+1}-1}$ under optimal play.

Full proof

Setup and notation. After all marks are made, the stick is cut into a finite multiset M of positive piece lengths summing to 1. For a multiset $M = \{p_1 \geq p_2 \geq \dots \geq p_m\}$ sorted in decreasing order, define the *alternating sum*

$$D(M) = p_1 - p_2 + p_3 - p_4 + \dots = \sum_{i=1}^m (-1)^{i+1} p_i,$$

and write $\text{odd}(M) = \sum_{i \text{ odd}} p_i$, $\text{even}(M) = \sum_{i \text{ even}} p_i$ for the sums of the odd- and even-indexed pieces. Note $\text{odd} + \text{even} = \sum M$ and $\text{odd} - \text{even} = D(M) \geq 0$ (since $D = (p_1 - p_2) + (p_3 - p_4) + \dots \geq 0$).

1. The claiming game

Lemma 1.1 (value of the claiming game). *In the claiming game played on a multiset M with Liu Bang moving first, Liu Bang can guarantee a total of at least $\text{odd}(M)$, and Xiang Yu can guarantee a total of at least $\text{even}(M)$. Hence under optimal play Liu Bang obtains exactly $\text{odd}(M) = \frac{\sum M + D(M)}{2}$.*

Proof. Induction on $|M|$. If $|M| = 1$ the claims are trivial. Let $|M| \geq 2$ and sort $M = \{p_1 \geq \dots \geq p_m\}$.

Liu Bang’s guarantee. LB first takes p_1 . Suppose XY then takes p_j ($j \geq 2$). The remainder is $M' = M \setminus \{p_1, p_j\}$, on which LB moves first; by induction LB guarantees $\text{odd}(M')$ from M' . The sorted order of M' is $p_2, \dots, p_{j-1}, p_{j+1}, \dots, p_m$; its odd-indexed elements are the p_i with i even,

$2 \leq i < j$, together with the p_i with i odd, $i > j$. Hence LB's total is

$$p_1 + \text{odd}(M') = p_1 + \sum_{\substack{i \text{ even} \\ i < j}} p_i + \sum_{\substack{i \text{ odd} \\ i > j}} p_i = \text{odd}(M) + \left(\sum_{\substack{i \text{ even} \\ i < j}} p_i - \sum_{\substack{i \text{ odd} \\ 3 \leq i \leq j}} p_i \right),$$

where the last sum ends at $i = j - 1$ if j is even and at $i = j$ if j is odd. In either case the bracket is a sum of consecutive differences $(p_2 - p_3) + (p_4 - p_5) + \dots \geq 0$. So LB gets at least $\text{odd}(M)$ no matter what XY does.

Xiang Yu's guarantee. Suppose LB first takes p_j . XY takes a largest remaining piece ℓ ($\ell = p_1$ if $j \neq 1$, else $\ell = p_2$), and then, moving first on $M'' = M \setminus \{p_j, \ell\}$, guarantees $\text{odd}(M'')$ by induction. If $j = 1$: XY's total is $p_2 + \text{odd}(M \setminus \{p_1, p_2\}) = p_2 + \sum_{i \text{ odd} \geq 3} p_i = \text{even}(M) + \sum_{i \text{ odd} \geq 3} (p_i - p_{i+1}) \geq \text{even}(M)$. If $j \geq 2$: XY's total is $p_1 + \text{odd}(M \setminus \{p_1, p_j\}) \geq \text{odd}(M) \geq \text{even}(M)$ by the computation above (with the roles exchanged) and $D(M) \geq 0$. ■

Since $\text{odd}(M) + \text{even}(M) = \sum M = 1$, LB's share under optimal play equals $\text{odd}(M) = \frac{1+D(M)}{2}$. Consequently the whole game is equivalent to the following one-dimensional optimization:

Reduction. LB chooses a composition of 1 into $m \leq n+1$ positive parts (his $m-1 \leq n$ marks). XY then refines this composition with at most n further cuts (each of XY's marks lies in the interior of one current piece and splits it in two; conversely every such refinement is achievable, with all marks distinct, since cut points are interior to pieces). LB's resulting share is $\frac{1+D(M)}{2}$, where M is the final multiset of pieces.

We therefore must compute $D^* = \max_{\text{LB}} \min_{\text{XY}} D(M)$; the answer will be $c = \frac{1+D^*}{2}$.

2. Two representations of D

Lemma 2.1 (odd-set identity). *For a finite multiset M of positive reals and $t \geq 0$ let $N(t) = \#\{x \in M : x > t\}$. Then*

$$D(M) = |\{t \geq 0 : N(t) \text{ is odd}\}|,$$

the Lebesgue measure of the “odd set”.

Proof. Sort $M = \{p_1 \geq \dots \geq p_m\}$ and put $p_{m+1} = 0$. For $t \in [p_{i+1}, p_i)$ we have $N(t) = i$ (ties give empty intervals, which do not affect the measure). Hence $|\{t : N(t) \text{ odd}\}| = \sum_{i \text{ odd}} (p_i - p_{i+1}) = D(M)$. ■

A *matching* μ of M is a collection of disjoint unordered pairs $\{x, y\}$ of elements of M (elements may be left unpaired); its *cost* is

$$\text{cost}(\mu) = \sum_{\{x, y\} \in \mu} |x - y| + \sum_{z \text{ unmatched}} z.$$

Lemma 2.2 (matching formula). $D(M) = \min_{\mu} \text{cost}(\mu)$, the minimum over all matchings of M .

Proof. For $t \geq 0$ and a matching μ let $c_{\mu}(t)$ be the number of pairs of μ with exactly one element $> t$, plus the number of unmatched elements $> t$. Since $|x - y| = \int_0^{\infty} \mathbf{1}[\text{exactly one of } x, y > t] dt$ and $z = \int_0^{\infty} \mathbf{1}[z > t] dt$, we have $\text{cost}(\mu) = \int_0^{\infty} c_{\mu}(t) dt$. If $b(t)$ denotes the number of pairs with both elements $> t$, then $N(t) = 2b(t) + c_{\mu}(t)$, so $c_{\mu}(t) \equiv N(t) \pmod{2}$ and $c_{\mu}(t) \geq 0$; in particular $c_{\mu}(t) \geq \mathbf{1}[N(t) \text{ odd}]$ for every t . Integrating and using Lemma 2.1 gives $\text{cost}(\mu) \geq D(M)$ for every

μ . Equality is attained by the *adjacent* matching: sorting $M = \{p_1 \geq \dots \geq p_m\}$ and pairing $(p_1, p_2), (p_3, p_4), \dots$ gives cost $\sum_{i \text{ odd}} (p_i - p_{i+1}) = D(M)$. ■

Corollary 2.3 (parity cancellation). *If M contains two equal elements x, x , then $D(M) = D(M \setminus \{x, x\})$. Consequently, if every value in M occurs with even multiplicity, then $D(M) = 0$.*

Proof. In the decreasing ordering of M the two copies of x occupy two adjacent positions $j, j+1$ and contribute $(-1)^{j+1}x + (-1)^{j+2}x = 0$; removing them shifts all later elements by two positions, preserving the sign of each term. ■

3. Part A: the geometric split (lower bound)

Put $T_n = \frac{1}{2^{n+1} - 1}$. LB marks n points dividing the stick into $n+1$ pieces of lengths

$$a_i = 2^{i-1}T_n \quad (i = 1, \dots, n+1), \quad \sum_{i=1}^{n+1} a_i = (2^{n+1} - 1)T_n = 1.$$

Definition. Positive reals $a_1 \leq a_2 \leq \dots \leq a_m$ are *super-increasing with margin $\delta > 0$* if $a_1 \geq \delta$ and $a_i \geq \delta + \sum_{j < i} a_j$ for every $i \geq 2$.

The family above is super-increasing with margin $\delta = T_n$: indeed $a_1 = T_n$ and $\sum_{j < i} a_j = (2^{i-1} - 1)T_n = a_i - T_n$, so $a_i = \delta + \sum_{j < i} a_j$.

Theorem 3.1. *Let $a_1 \leq \dots \leq a_m$ be super-increasing with margin δ , and let M be obtained from these pieces by at most $m-1$ cuts (each cut splits one current piece into two positive pieces). Then $D(M) \geq \delta$.*

Proof. By Lemma 2.2 it suffices to show that **every** matching μ of M has $\text{cost}(\mu) \geq \delta$. Fix μ . Each element of M is a fragment of exactly one piece a_i . Build a multigraph G on the vertex set $\{1, \dots, m\}$: - for each pair $\{x, y\} \in \mu$ whose fragments come from two **different** pieces $i \neq j$, add an edge $\{i, j\}$; - for each pair $\{x, y\} \in \mu$ whose two fragments come from the **same** piece i , add a **loop** at i .

Unmatched fragments create no edges. Write e for the total number of edges (loops included) and u for the number of unmatched fragments. Counting fragments in two ways, with $c \leq m-1$ the number of cuts:

$$2e + u = \#\{\text{fragments}\} = m + c \leq 2m - 1,$$

so $e \leq m-1$ (as e is an integer and $u \geq 0$). Let C range over the connected components of G (connectivity ignores loops; loops still count as edges of their component), and let e_C, v_C be the numbers of edges and vertices of C . If $e_C \geq v_C$ held for **every** component, summing would give $e \geq m$, contradicting $e \leq m-1$. Hence some component C has $e_C \leq v_C - 1$. A connected multigraph on v_C vertices has at least $v_C - 1$ edges, with equality only if it is a tree (a loop or a parallel edge would force $e_C \geq v_C$); so C is a tree. In particular C has **no loops**, i.e. no self-pair of fragments occurs inside C : every pair of μ within C is a cross-pair between two different pieces of C , and these cross-pairs are exactly the $v_C - 1$ edges of the tree.

Tree inequality. A tree is bipartite: choose signs $\epsilon_i \in \{\pm 1\}$ for $i \in C$ so that $\epsilon_i = -\epsilon_j$ whenever $\{i, j\}$ is an edge. We claim

$$\left| \sum_{i \in C} \epsilon_i a_i \right| \leq \text{cost}(C), \quad \text{cost}(C) := \sum_{\substack{\{x, y\} \in \mu \\ \text{within } C}} |x - y| + \sum_{\substack{z \text{ unmatched} \\ \text{in } C}} z.$$

Indeed, expand each a_i as the sum of its fragments and group the signed sum $\sum_{i \in C} \epsilon_i a_i$ according to μ . A cross-pair $\{x, y\}$ with x from i and y from j contributes $\epsilon_i x + \epsilon_j y = \epsilon_i(x - y)$, of absolute value $|x - y|$; an unmatched fragment z of piece i contributes $\epsilon_i z$, of absolute value z . By the triangle inequality $|\sum_{i \in C} \epsilon_i a_i| \leq \sum_{\text{cross-pairs}} |x - y| + \sum_{\text{unmatched}} z = \text{cost}(C)$.

Margin. Let k be the largest index in C . Then, using the reverse triangle inequality and the margin property,

$$\left| \sum_{i \in C} \epsilon_i a_i \right| \geq a_k - \sum_{i \in C \setminus \{k\}} a_i \geq \delta,$$

because $C \setminus \{k\} \subseteq \{1, \dots, k-1\}$: if $k \geq 2$ then $a_k \geq \delta + \sum_{j < k} a_j \geq \delta + \sum_{i \in C \setminus \{k\}} a_i$, and if $k = 1$ then $C = \{1\}$ and $a_1 \geq \delta$.

Combining, $\text{cost}(\mu) \geq \text{cost}(C) \geq |\sum_{i \in C} \epsilon_i a_i| \geq \delta$. Since μ was arbitrary, Lemma 2.2 gives $D(M) \geq \delta$. ■

Corollary 3.2 (lower bound). *With the geometric split, whatever XY does, $D(M) \geq T_n$, so LB guarantees a share of at least*

$$\frac{1 + T_n}{2} = \frac{1}{2} \cdot \frac{2^{n+1}}{2^{n+1} - 1} = \frac{2^n}{2^{n+1} - 1}.$$

Proof. The final multiset M is a refinement of the geometric family (which is super-increasing with margin T_n) by at most $n = m - 1$ cuts (with $m = n + 1$); Theorem 3.1 applies. ■

4. Part B: Xiang Yu's counterstrategy (upper bound)

Now fix **any** composition of 1 into $m \leq n + 1$ positive pieces $a_1 \leq \dots \leq a_m$ chosen by LB. We exhibit an XY strategy ($\leq n$ cuts) forcing $D \leq T_n$.

Case $m \leq n$. XY bisects every piece: $m \leq n$ cuts. Each piece x becomes two pieces $x/2, x/2$, so every value occurs with even multiplicity in the final multiset; by Corollary 2.3, $D = 0$.

Case $m = n + 1$. This rests on two lemmas about *chains*. For a nonempty finite multiset $B = \{b_1 \geq b_2 \geq \dots \geq b_r\}$ of positive reals define its *chain value* as the nested absolute difference

$$v(B) = |\dots ||b_1 - b_2| - b_3| - \dots - b_r|.$$

Lemma 4.1 (realizability lemma). *Let P, N be disjoint finite multisets of positive reals, with sums $\Sigma P, \Sigma N$, and $P \cup N \neq \emptyset$. - (a) If $\Sigma P = \Sigma N$, then some ordering of $P \cup N$ has nested absolute difference 0. - (b) If $v := \Sigma P - \Sigma N > 0$ and no subset of $P \cup N$ has sum strictly between ΣN and ΣP , then some ordering of $P \cup N$ has nested absolute difference v .*

Proof. We construct an ordering $e_1, \dots, e_r$ of $P \cup N$ by a greedy process. Set $E_0 = 0$. At step k , while elements remain: if $E_{k-1} > 0$, append any remaining element $e_k \in N$ and put $E_k = E_{k-1} - e_k$; if $E_{k-1} < 0$, append any remaining element $e_k \in P$ and put $E_k = E_{k-1} + e_k$; if $E_{k-1} = 0$, append any remaining element from either (nonempty) side, with the corresponding sign.

We claim this process never gets stuck, i.e. the side it must draw from is never empty while the other side is nonempty. Indeed, suppose the process halts with a remaining part $P' \subseteq P$ and $N' \subseteq N$, not both empty, and let $I \subseteq P$ and $J \subseteq N$ be the parts already processed, so the current signed sum is $E = \Sigma I - \Sigma J$.

- *Halt with $E > 0$* : this requires N exhausted ($J = N$) and $P' \neq \emptyset$. Then $E = \Sigma I - \Sigma N > 0$, so $\Sigma I > \Sigma N$, while $\Sigma I = \Sigma P - \Sigma P' < \Sigma P$ (as $P' \neq \emptyset$ and all elements are positive). Hence ΣI lies strictly between ΣN and ΣP : impossible — in case (b) by the hypothesis, and in case (a) because $\Sigma I \leq \Sigma P = \Sigma N$ would give $E \leq 0$.
- *Halt with $E < 0$* : this requires P exhausted ($I = P$) and $N' \neq \emptyset$. Then the full signed sum is $\Sigma P - \Sigma N = E - \Sigma N' < 0$, contradicting $\Sigma P - \Sigma N = v > 0$ in case (b) and $\Sigma P - \Sigma N = 0$ in case (a).
- *Halt with $E = 0$* : then both sides are empty (at $E = 0$ either nonempty side may be drawn from), so nothing remains — the process is finished, not stuck.

Hence the process consumes all of $P \cup N$ and ends at $E_r = \Sigma P - \Sigma N$.

Finally we check that the nested absolute difference of the ordering $e_1, \dots, e_r$ equals $|E_r|$. Define $R_1 = e_1$ and $R_k = |R_{k-1} - e_k|$; we claim $R_k = |E_k|$ for all k . This holds for $k = 1$ since $E_1 = \pm e_1$. Inductively: if $E_{k-1} \geq 0$ and $e_k \in N$, then $E_k = E_{k-1} - e_k$ and $|E_k| = ||E_{k-1}| - e_k| = |R_{k-1} - e_k| = R_k$; if $E_{k-1} < 0$ and $e_k \in P$, then $E_k = -|E_{k-1}| + e_k$ and again $|E_k| = ||E_{k-1}| - e_k| = R_k$; if $E_{k-1} = 0$ then $|E_k| = e_k = |0 - e_k| = R_k$. Thus the chain value is $R_r = |E_r| = |\Sigma P - \Sigma N|$, which is 0 in case (a) and v in case (b). ■

Theorem 4.2 (chain theorem). *Let $a_1, \dots, a_m$ be positive reals with sum S . Then some nonempty sub-multiset $B \subseteq \{a_1, \dots, a_m\}$ has chain value $v(B) \leq \frac{S}{2^m - 1}$.*

Proof. Consider the 2^m subset sums $s(I) = \sum_{i \in I} a_i$ ($I \subseteq \{1, \dots, m\}$); they lie in $[0, S]$. Sort them as $0 = t_0 \leq t_1 \leq \dots \leq t_{2^m-1} = S$ (with multiplicity). The $2^m - 1$ consecutive gaps sum to $t_{2^m-1} - t_0 \leq S$, so some consecutive gap satisfies

$$g := t_{j+1} - t_j \leq \frac{S}{2^m - 1}.$$

By definition of the sorted list, **no subset sum lies strictly between t_j and t_{j+1}** . Pick subsets A, B with $s(A) = t_{j+1}$, $s(B) = t_j$, and set $P = \{a_i : i \in A \setminus B\}$, $N = \{a_i : i \in B \setminus A\}$ (as multisets). Then $P \cup N \neq \emptyset$ and $\Sigma P - \Sigma N = s(A) - s(B) = g \geq 0$.

If $g = 0$, Lemma 4.1(a) gives an ordering of $P \cup N$ with chain value $0 \leq \frac{S}{2^m-1}$.

If $g > 0$: any subset J of $P \cup N$ with $\Sigma N < \Sigma J < \Sigma P$ would lift to the subset $J \cup (A \cap B)$ of $\{a_1, \dots, a_m\}$ whose sum lies strictly between $s(B)$ and $s(A)$, a contradiction. So the hypothesis of Lemma 4.1(b) holds, and some ordering of $P \cup N$ has chain value $g \leq \frac{S}{2^m-1}$. ■

(The bound is sharp: for $a_i = 2^{i-1}$ the subset sums are exactly $0, 1, \dots, 2^m - 1$, every gap is $1 = \frac{S}{2^m-1}$, and every nonempty chain value is a nonzero integer, hence ≥ 1 .)

XY's strategy from a chain. Let $B = \{b_1 \geq \dots \geq b_r\}$ be a nonempty subset of LB's pieces with chain value $v = v(B) \leq T_n$ (Theorem 4.2 with $S = 1$, $m = n + 1$). XY plays as follows. Maintain a *running value* v_{run} (initially b_1 , represented by one physical piece) and process $b_2, \dots, b_r$ in turn. To process b_k with current running value $v = v_{\text{run}}$: - if $v = 0$: declare b_k the new running piece ($v_{\text{run}} := b_k$); no cut; - if $v = b_k > 0$: leave both pieces (they are equal and will cancel); $v_{\text{run}} := 0$; no cut; - if $v \neq b_k$ and both positive: cut the **larger** of the two pieces $\{v_{\text{run}}, b_k\}$ at distance $\min(v, b_k)$ from one end, producing pieces of sizes $\min(v, b_k)$ and $|v - b_k|$. The piece of size $\min(v, b_k)$ together with the smaller of the two original pieces forms an **equal pair** (both pieces remain in play; they will cancel in D by Corollary 2.3). The piece of size $|v - b_k|$ becomes the new running piece; $v_{\text{run}} := |v - b_k|$. One cut.

After processing all of B , XY bisects every piece outside B ($n + 1 - r$ cuts). The total number of cuts is at most $(r - 1) + (n + 1 - r) = n$. All cut points are distinct (each cut is made inside a piece never cut before: the running piece is created by the previous step's cut, and each b_k is used at most once) and each lies in the interior of one of LB 's original pieces; hence XY can mark all of them simultaneously, and they differ from LB 's marks.

In the final multiset, every value occurs with even multiplicity except possibly the final running value $v_{\text{run}} = v(B)$: bisected pieces contribute pairs of halves, each equal pair formed above contributes two equal pieces, each tie leaves two equal pieces, and only the last running piece (if $v(B) > 0$) is left over. Removing equal pairs (Corollary 2.3) leaves $\{v(B)\}$ or $\emptyset$, whence

$$D(M) = v(B) \leq T_n, \quad \text{so } LB\text{'s share is at most } \frac{1 + T_n}{2} = \frac{2^n}{2^{n+1} - 1}.$$

(If $v(B) = 0$ every value has even multiplicity and $D = 0$.)

5. Conclusion

Combining Corollary 3.2 (LB guarantees $\frac{1+T_n}{2}$ via the geometric split) with Section 4 (XY always holds LB to at most $\frac{1+T_n}{2}$), the largest value LB can guarantee is

$$c = \frac{1 + T_n}{2} = \frac{2^n}{2^{n+1} - 1}.$$

Sanity checks: $n = 1$: $c = \frac{2}{3}$ (LB marks at $\frac{1}{3}$; any cut of the long piece leaves $D \geq \frac{1}{3}$); $n = 2$: $c = \frac{4}{7}$; $n = 3$: $c = \frac{8}{15}$ — all matching exhaustive grid computation.