

An Unofficial Compendium of Worked Solutions to the Problems in *The Method of Coordinates*

by I.M. Gelfand, E. G. Glagoleva & A. A. Kirillov

Deepak Venkatesh

September 19, 2026

Abstract

This book is an independent and unofficial collection of worked solutions to problems inspired by the book *The Method of Coordinates* by I. M. Gelfand, E. G. Glagoleva and A. A. Kirillov, first published in 1967. The original text presents a wide range of problems in elementary algebra, some accompanied by solutions and others left as exercises for the reader.

The aim of this book is to provide clear, logically consistent solutions to all problems addressed, written in a style accessible to motivated students and self learners. Proofs have been verified primarily by hand, and in some cases supported by small computational checks written in Scheme (a dialect of Lisp) or in Python. Large language models were used only as auxiliary tools, primarily for generating figures in L^AT_EX and for reviewing exposition.

This work is not affiliated with, endorsed by, or approved by the original authors or publisher. Any errors, omissions, or inconsistencies are entirely my own. Issues and corrections may be reported at this repository.

Copyright and Disclaimer

Unofficial companion.

This document is an **independent, unofficial collection of worked solutions** to problems from *The Method of Coordinates* by I. M. Gelfand , E. G. Glagoleva and A. A. Kirillov (first published 1967).

This work is **not affiliated with, endorsed by, or approved by** the original authors or the publisher. The names of the authors and the title of the book are used only for identification and reference.

The original problem statements are not reproduced here. All exposition and solutions are written independently by the author of this document.

Copyright © 2026 Deepak Venkatesh

License:

This work is licensed under a Creative Commons “Attribution-NonCommercial-ShareAlike 4.0 International” license.



Project repository:

<https://github.com/deepak-venkatesh/gelfand-method-coordinates>

For
My Daughter and Son

Contents

I	Part 1	7
1	Chapter 1: The Coordinates of Points on a Line	7
1.1	The Number Axis	7
1.2	The Absolute Value of a Number	10
1.3	The Distance Between Two Points	10
2	Chapter 2: The Coordinates of Points in the Plane	10
2.1	The Coordinate Plane	10
2.2	Relations Connecting Coordinates	10
2.3	The Distance Between Two Points	10
2.4	Defining Figures	10
2.5	We Begin to Solve Problems	10
2.6	Other System of Coordinates	10
3	Chapter 3: The Coordinates of a Point in Space	10
3.1	Coordinate Axes and Planes	10
3.2	Defining Figures in Space	10
II	Part 2	10
4	Chapter 1: Introduction	10
4.1	Some General Considerations	10
4.2	Geometry as an Aid in Calculation	10
4.3	The Need for Introducing Four-Dimensional Space	10
4.4	The Peculiarities of Four-Dimensional Space	10
4.5	Some Physics	10
5	Chapter 2: Four-Dimensional Space	10
5.1	Coordinate Axes and Planes	10
5.2	Some Problems	10

6 Chapter 3: The Four-Dimensional Cube	10
6.1 The Definition of the Sphere and the Cube	10
6.2 The Structure of the Four-Dimensional Cube	10
6.3 Problems on the Cube	10

Working Notes

Challenging Problems for Middle Schoolers:

Problem Numbers:

Typographical errors in:

Problem Numbers:

Part I

Part 1


1 Chapter 1: The Coordinates of Points on a Line

1.1 The Number Axis

Exercise 1


(a)

The points are marked below. Note that $\frac{13}{3}$ lies somewhere between 4 and 5.



(b)

The two points are 3 units to the right and 3 units to the left of M as shown in the line below.



Exercise 2

(a)

Going by the convention that the right side of the number line indicates larger number, in this exercise $a > b$.

(b)

On the right are the larger numbers

- $A(-3)$
- $B(4)$
- $B(4)$
- $A(3)$

Exercise 3

It is easy to say that $A(a)$ is on the right of $B(-a)$ but this assumes that a is a positive number. So this could be true but the other case is that if a were a negative number then $A(a)$ would be a negative number and $B(-a)$ would be a positive number, thus $B(-a)$ would be on the right.

The third case is that $a = 0$ then both A and B would be 0 and they would be at the same point on the number line.

Exercise 4

1.2 The Absolute Value of a Number

1.3 The Distance Between Two Points

2 Chapter 2: The Coordinates of Points in the Plane

2.1 The Coordinate Plane

2.2 Relations Connecting Coordinates

2.3 The Distance Between Two Points

2.4 Defining Figures

2.5 We Begin to Solve Problems

2.6 Other System of Coordinates

3 Chapter 3: The Coordinates of a Point in Space

3.1 Coordinate Axes and Planes

3.2 Defining Figures in Space

Part II

Part 2

4 Chapter 1: Introduction

4.1 Some General Considerations

4.2 Geometry as an Aid in Calculation

4.3 The Need for Introducing Four-Dimensional Space

4.4 The Peculiarities of Four-Dimensional Space

4.5 Some Physics

GitHub repository: <https://github.com/deepak-venkatesh/gelfand-method-coordinates> 10

5 Chapter 2: Four-Dimensional Space

5.1 Coordinate Axes and Planes

5.2 Some Problems