NABL & ISO APPROVED



 Patient Name
 : Mr JAI RAM RAJANI
 Bill Date
 : Jul 21, 2022, 03:47 PM

 DOB/Age/Gender
 : 63Yrs/Male
 Sample Collected : Jul 21, 2022, 03:11 PM

 Patient ID
 : 1085133
 Sample Received : Jul 21, 2022, 03:48 PM

 Referred by
 : Dr.
 Report Date
 : Jul 21, 2022, 05:31 PM

Sample Type : Whole blood EDTA BarcodeNo : H167638
Client : STANDARD KANPUR Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range

HEMATOLOGY REPORT Lifecare Health Gold Plus Package Complete Blood Count (CBC)

Complete Blood Count (CBC)				
RBC PARAMETERS	<u> </u>	(<i>020)</i>		
Hemoglobin Method : colorimetric	10.7	g/dL	13.0 - 17.0	
RBC Count Method : Electrical impedance	5.50	10^6/μl	4.5 - 5.5	
PCV Method : Calculated	36.2	%	40 - 50	
MCV Method : Calculated	65.5	fl	83 - 101	
MCH Method : Calculated	19.4	pg	27 - 32	
MCHC Method : Calculated	29.6	g/dL	31.5 - 34.5	
RDW (CV) Method : Calculated	17.5	%	11.6 - 14.0	
RDW-SD Method : Calculated	40.8	fl	35.1 - 43.9	
WBC PARAMETERS				
TLC Method : Electrical impedance and microscopy DIFFERENTIAL LEUCOCYTE COUNT	11	10^3/µl	4 - 10	
Neutrophils	53	%	40 - 80	
Lymphocytes	36	%	20 - 40	
Monocytes	5	%	2 - 10	
Eosinophils	6	%	1 - 6	
Basophils	0	%	<2.0	
Absolute leukocyte counts Method : calculated				
Neutrophils*	5.83			
Lymphocytes*	3.96			
Monocytes*	0.55			
Eosinophils*	0.66	103/uL	0.02 - 0.44	
Basophils*	0			
PLATELET PARAMETERS				
Platelet Count Method : Electrical impedance and microscopy	217	10^3/μl	150 - 410	
Mean Platelet Volume (MPV) Method : Calculated	11.7	fL	9.3 - 12.1	









NABL & ISO APPROVED



 Patient Name
 : Mr JAI RAM RAJANI
 Bill Date
 : Jul 21, 2022, 03:47 PM

 DOB/Age/Gender
 : 63Yrs/Male
 Sample Collected : Jul 21, 2022, 03:11 PM

 Patient ID
 : 1085133
 Sample Received : Jul 21, 2022, 03:48 PM

 Referred by
 : Dr.
 Report Date
 : Jul 21, 2022, 05:31 PM

Sample Type : Whole blood EDTA BarcodeNo : H167638

Client : STANDARD KANPUR Report Status : Final Report

Test Description	Value(s)	Unit(s)	Reference Range
PCT Method : Calculated	0.3	%	0.17 - 0.32
PDW Method : Calculated	15.6	fL	8.3 - 25.0
P-LCR Method : Calculated	40.4	%	18 - 50
P-LCC Method : Calculated	88.0	%	44 - 140
Mentzer Index	11.91	%	

Interpretation:

CBC provides information about red cells, white cells and platelets. Results are useful in the diagnosis of anemia, infections, leukemias, clotting disorders and many other medical conditions.









NABL & ISO APPROVED



 Patient Name
 : Mr JAI RAM RAJANI
 Bill Date
 : Jul 21, 2022, 03:47 PM

 DOB/Age/Gender
 : 63Yrs/Male
 Sample Collected : Jul 21, 2022, 03:11 PM

 Patient ID
 : 1085133
 Sample Received : Jul 21, 2022, 03:48 PM

 Referred by
 : Dr.
 Report Date
 : Jul 21, 2022, 07:23 PM

Sample Type : Whole blood EDTA BarcodeNo : H167638

Client : STANDARD KANPUR Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range

HEMATOLOGY REPORT Lifecare Health Gold Plus Package HbA1C (Glycosylated Hemoglobin)

GLYCOSYLATED HEMOGLOBIN (HbA1c) 7.3 % < 5.7 ESTIMATED AVERAGE GLUCOSE 162.81 mg/dL -

Interpretation:

Interpretation For HbA1c% As per American Diabetes Association (ADA)

Reference Group	HbA1c in %
Non diabetic adults >=18 years	<5.7
At risk (Prediabetes)	5.7 - 6.4
Diagnosing Diabetes	>= 6.5
Therapeutic goals for glycemic control	Age > 19 years Goal of therapy: < 7.0 Age < 19 years Goal of therapy: <7.5

Note:

10

240

Comments :

HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations ADA criteria for correlation between HbA1c & Mean plasma glucose levels.

1	HbA1c(%)	Mean Plasma Glucose (mg/dL)	HbA1c(%)	Mean Plasma Glucose (mg/dL)
	6	126	12	298
Ì	8	183	14	355

16





723-000-0806





413

^{1.} Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled. 2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

NABL & ISO APPROVED



 Patient Name
 : Mr JAI RAM RAJANI
 Bill Date
 : Jul 21, 2022, 03:47 PM

 DOB/Age/Gender
 : 63Yrs/Male
 Sample Collected : Jul 21, 2022, 03:11 PM

 Patient ID
 : 1085133
 Sample Received : Jul 21, 2022, 03:48 PM

 Referred by
 : Dr.
 Report Date
 : Jul 21, 2022, 07:24 PM

Sample Type : FLUORIDE F BarcodeNo : BC451833
Client : STANDARD KANPUR Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range

BIOCHEMISTRY REPORT
Lifecare Health Gold Plus Package
Glucose Fasting (BSF)

GLUCOSE FASTING **108.6** mg/dL 70 - 100

Method : Hexokinase

Note:- Kindly Correlate Clinically.



© 723-000-0806



are@lifecarehealthlabs.com



NABL & ISO APPROVED



 Patient Name
 : Mr JAI RAM RAJANI
 Bill Date
 : Jul 21, 2022, 03:47 PM

 DOB/Age/Gender
 : 63Yrs/Male
 Sample Collected : Jul 21, 2022, 03:11 PM

 Patient ID
 : 1085133
 Sample Received : Jul 21, 2022, 03:48 PM

 Referred by
 : Dr.
 Report Date
 : Jul 21, 2022, 06:43 PM

Sample Type : Serum BarcodeNo : BC451834
Client : STANDARD KANPUR Report Status : Final Report

Test Description	Value(s)	Unit(s)	Reference Range	
------------------	----------	---------	-----------------	--

BIOCHEMISTRY REPORT Lifecare Health Gold Plus Package

Liver Function Test (LFT)

	Liver Function T	est (LFT)	
BILIRUBIN TOTAL Method : Photometric	0.2	mg/dL	0.2 - 1.2
BILIRUBIN DIRECT Method : Diazo Reaction	0.1	mg/dL	0.0 - 0.5
BILIRUBIN INDIRECT Method : Calculation (T Bil - D Bil)	0.1	mg/dL	0.1 - 1.0
SGOT/AST Method : IFCC without P5P	34.0	U/L	5 - 34
SGPT/ALT Method : IFCC without P5P	22.1	U/L	0 to 55
SGOT/SGPT Ratio	1.54	-	-
ALKALINE PHOSPHATASE Method: IFCC	67.0	U/L	40 - 150
TOTAL PROTEIN Method : Biuret	6.6	g/dL	6.4 - 8.3
ALBUMIN Method : BCG	3.9	gm/dL	3.8 - 5.0
GLOBULIN Method : Calculation (T.P - Albumin)	2.7	g/dL	2.3 - 3.5
ALBUMIN : GLOBULIN RATIO Method : Calculation (Albumin/Globulin)	1.44	-	1.0 - 2.1
GAMMA GLUTAMYL TRANSFERASE (GGT) Method : Photometric	34.3	U/L	12 - 64

vietnoa : Pnotometrio

Interpretation:

The liver filters and processes blood as it circulates through the body. It metabolizes nutrients, detoxifies harmful substances, makes blood clotting proteins, and performs many other vital functions. The cells in the liver contain proteins called enzymes that drive these chemical reactions. When liver cells are damaged or destroyed, the enzymes in the cells leak out into the blood, where they can be measured by blood tests Liver tests check the blood for two main liver enzymes. Aspartate aminotransferase (AST),SGOT: The AST enzyme is also found in muscles and many other tissues besides the liver. Alanine aminotransferase (ALT), SGPT: ALT is almost exclusively found in the liver. If ALT and AST are found together in elevated amounts in the blood, liver damage is most likely present. Alkaline Phosphatase and GGT: Another of the liver's key functions is the production of bile, which helps digest fat. Bile flows through the liver in a system of small tubes (ducts), and is eventually stored in the gallbladder, under the liver. When bile flow is slow or blocked, blood levels of certain liver enzymes rise: Alkaline phosphatase Gamma-utamyl transpeptidase (GGT) Liver tests may check for any or all of these enzymes in the blood. Alkaline phosphatase is by far the most commonly tested of the three. If alkaline phosphatase and GGT are elevated, a problem with bile flow is most likely present. Bile flow problems can be due to a problem in the liver, the gallbladder, or the tubes connecting them. Proteins are important building blocks of all cells and tissues. Proteins are necessary for your body's growth, development, and health. Blood contains two classes of protein, albumin and globulin. Albumin proteins keep fluid from leaking out of blood vessels. Globulin proteins play an important role in your immune system. Low total protein may indicate: 1.bleeding 2.liver disorder 3.malnutrition 4.agammaglobulinemia High Protein levels 'Hyperproteinemia: May be seen in dehydration due to inadequate water intake or to excessive w



Dr.S.P.Mittal
MBBS. MD PATHOLOGY

© 723-000-0806

are@lifecarehealthlabs.com

mww.lifecarehealthlabs.com

NABL & ISO APPROVED



 Patient Name
 : Mr JAI RAM RAJANI
 Bill Date
 : Jul 21, 2022, 03:47 PM

 DOB/Age/Gender
 : 63Yrs/Male
 Sample Collected : Jul 21, 2022, 03:11 PM

 Patient ID
 : 1085133
 Sample Received : Jul 21, 2022, 03:48 PM

 Referred by
 : Dr.
 Report Date
 : Jul 21, 2022, 07:26 PM

Sample Type : Serum BarcodeNo : BC451834
Client : STANDARD KANPUR Report Status : Final Report

Test Description	Value(s)	Unit(s)	Reference Range	
------------------	----------	---------	-----------------	--

BIOCHEMISTRY REPORT Lifecare Health Gold Plus Package

	Kidney Function Tes	st (KFT)	
BLOOD UREA Method : Urease	21.0	mg/dL	18 - 55
CREATININE Method : Photometric	0.9	mg/dL	0.72 - 1.25
BUN Method : Urease	9.81	mg/dL	8.4 - 25.7
BUN/CREATININE RATIO	10.9		
URIC ACID Method : Uricase	7.1	mg/dL	3.5 - 7.2
CALCIUM Serum Method : Arsenazo III	10.0	mg/dL	8.8 - 10.0
PHOSPHORUS Method : Photometric	4.4	mg/dL	2.3 - 4.7
SODIUM Method : Potentiometric	136.2	mmol/L	136 - 145
POTASSIUM Method : Potentiometric	4.21	mmol/L	3.5 - 5.1
Chloride Method : Photometric	102.4	mmol/L	98 - 107

Interpretation:

SUMMARY:-Kidney function tests is a collective term for a variety of individual tests and proceduresthat can be done toevaluate how well the kidneys are functioning. Many conditions can affect the ability of the kidneys to carryout their vital functions. Somelead to a rapid (acute) decline in kidney functionothers lead to a gradual (chronic) declineinfunction. Both result in a buildup of toxic waste subst done on urine samples, as well as on blood samples. A number of symptoms may indicate a problem with your kidneys. These include: high blood pressure, blood in urine frequent urges to urinate, difficulty beginning urination, painful urination, swelling in the hands and feet due to a buildup of fluids in the body. A single symptom may not mean something serious. However, when occurring simultaneously, these symptoms suggest that your kidneys are not working properly. Kidney function tests can help determine the reason. Electrolytes (sodium, potassium, and chloride) are present in the human body and the balancing act of the electrolytes in our bodies is essential for normal function of our cells and organs. There has to be a balance. Ionized calcium this test if you have signs of kidney or parathyroid disease. The test may also be done to monitor progress and treatment of these diseases.





© 723-000-0806

□ care@lifecarehealthlabs.com



NABL & ISO APPROVED



Patient Name : Mr JAI RAM RAJANI Bill Date : Jul 21, 2022, 03:47 PM DOB/Age/Gender: 63Yrs/Male Sample Collected: Jul 21, 2022, 03:11 PM Patient ID : 1085133 Sample Received : Jul 21, 2022, 03:48 PM Referred by : Dr. Report Date : Jul 21, 2022, 06:43 PM

: BC451834 Sample Type : Serum BarcodeNo Client : STANDARD KANPUR Report Status : Final Report

Test Description Unit(s) Value(s) Reference Range

BIOCHEMISTRY REPORT Lifecare Health Gold Plus Package

Linial Drafile

	<u>Lipid Profil</u>	<u>e</u>	
TOTAL CHOLESTEROL Method : Enzymatic - Cholesterol Oxidase	141.0	mg/dL	Desirable : <200 Borderline : 200-239 High : >240
TRIGLYCERIDES Method : Colorimetric - Lip/Glycerol Kinase	163.2	mg/dL	Normal : <150 Borderline : 150-199 High : 200-499 Very high : >500
HDL CHOLESTEROL Method : Accelerator Selective Detergent	37.3	mg/dL	>40
NON HDL CHOLESTEROL Method : Calculated	103.7	mg/dL	<130
LDL CHOLESTEROL Method : Calculated	71.06	mg/dL	Optimal <100 Near optimal/above optimal 100-129 Borderline high 130-159 High 160-189 Very high >190
V.L.D.L CHOLESTEROL Method : Calculated	32.64	mg/dL	< 30
CHOL/HDL Ratio Method : Calculated	3.78	-	3.5 - 5.0
HDL/ LDL RATIO Method : Calculated	0.52	-	Desirable : 0.5 - 3.0
			Borderline : 3.1 - 6.0
1.01.4101.00.00	4.04		High : > 6.0
LDL/HDL Ratio	1.91	-	

Method: Calculated

Interpretation:

Lipid level assessments must be made following 9 to 12 hours of fasting, otherwise assay results might lead to erroneous interpretation. NCEP recommends of 3 different samples to be drawn at intervals of 1 week

NATIONAL LIPID ASSOCIATION RECOMMENDATIONS (NLA-2014)	TOTAL CHOLESTEROL in mg/dL	TRIGLYCERIDE in mg/dL	LDL CHOLESTEROL in mg/dL	NON HDL CHOLESTEROL in mg/dL
Optimal	<200	<150	<100	<130
Above Optimal			100-129	130 - 159
Borderline High	200-239	150-199	130-159	160 - 189
High	>=240	200-499	160-189	190 - 219
Very High	-	>=500	>=190	>=220





NABL & ISO APPROVED



 Patient Name
 : Mr JAI RAM RAJANI
 Bill Date
 : Jul 21, 2022, 03:47 PM

 DOB/Age/Gender
 : 63Yrs/Male
 Sample Collected : Jul 21, 2022, 03:11 PM

 Patient ID
 : 1085133
 Sample Received : Jul 21, 2022, 03:48 PM

 Referred by
 : Dr.
 Report Date
 : Jul 21, 2022, 06:43 PM

Sample Type : Serum BarcodeNo : BC451834
Client : STANDARD KANPUR Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range

BIOCHEMISTRY REPORT Lifecare Health Gold Plus Package

Iron Studies

IRON Method : Ferene	129.2	μg/dL	65 - 175
TIBC Method : Calculated	336.7	μg/dL	250 - 450
UIBC Method : Ferene	207.5	μg/dL	69 - 240
TRANSFERRIN SATURATION	38.37	%	-

Method: Method: Derived from IRON and TIBC values

Interpretation:

Increased levels due to iron ingestion or ineffective erythropoiesis. Decreased levels due to infection, inflammation, malignancy, menstruation and Fe deficiency. Needs to be taken into consideration with TIBC. Transferrin Saturation: Low level Transferrin Saturation can indicate iron deficiency, erythropoiesis, infection, or inflammation. High level Transferrin Saturation can indicate recent ingestion of dietary iron, ineffective erythropoiesis, haemochromatosis or liver disease. High TIBC, UIBC, or transferrin usually indicates iron deficiency, but they are also increased in pregnancy and with the use of oral contraceptives. Low TIBC, UIBC, or transferrin may occur if someone has: Hemochromatosis, Certain types of anemia due to accumulated iron, Malnutrition, kidney disease that causes a loss of protein in urine.









NABL & ISO APPROVED



 Patient Name
 : Mr JAI RAM RAJANI
 Bill Date
 : Jul 21, 2022, 03:47 PM

 DOB/Age/Gender
 : 63Yrs/Male
 Sample Collected : Jul 21, 2022, 03:11 PM

 Patient ID
 : 1085133
 Sample Received : Jul 21, 2022, 03:48 PM

 Referred by
 : Dr.
 Report Date
 : Jul 21, 2022, 05:50 PM

Sample Type : Serum BarcodeNo : BC451834
Client : STANDARD KANPUR Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range

BIOCHEMISTRY REPORT Lifecare Health Gold Plus Package Vitamin B12

 Vitamin - B12
 295.5
 pg/mL
 187 - 883

 Method : CMIA









NABL & ISO APPROVED



 Patient Name
 : Mr JAI RAM RAJANI
 Bill Date
 : Jul 21, 2022, 03:47 PM

 DOB/Age/Gender
 : 63Yrs/Male
 Sample Collected : Jul 21, 2022, 03:11 PM

 Patient ID
 : 1085133
 Sample Received : Jul 21, 2022, 03:48 PM

 Referred by
 : Dr.
 Report Date
 : Jul 21, 2022, 05:50 PM

Sample Type : Serum BarcodeNo : BC451834
Client : STANDARD KANPUR Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range

BIOCHEMISTRY REPORT Lifecare Health Gold Plus Package Vitamin D 25 - Hydroxy

Vitamin D 25 - Hydroxy 35.2 ng/mL Deficiency : <30 ng/mL

Method : CMIA



© 723-000-0806



are@lifecarehealthlabs.com



NABL & ISO APPROVED



 Patient Name
 : Mr JAI RAM RAJANI
 Bill Date
 : Jul 21, 2022, 03:47 PM

 DOB/Age/Gender
 : 63Yrs/Male
 Sample Collected : Jul 21, 2022, 03:11 PM

 Patient ID
 : 1085133
 Sample Received : Jul 21, 2022, 03:48 PM

 Referred by
 : Dr.
 Report Date
 : Jul 21, 2022, 05:50 PM

Sample Type : Serum BarcodeNo : BC451834
Client : STANDARD KANPUR Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range

BIOCHEMISTRY REPORT Lifecare Health Gold Plus Package

Thyroid Profile Total

TRIIODOTHYRONINE (T3) Method: CMIA	1.2	ng/ml	0.71 - 2.01
TOTAL THYROXINE (T4) Method: CMIA	8.1	μg/dL	4.87 - 11.2
THYROID STIMULATING HORMONE (TSH)	2.5	mIU/L	0.35 - 4.94

Method : CMIA

Interpretation:

Primary malfunction of the thyroid gland may result in excessive (hyper) or below normal (hypo) release of T3 or T4. In addition as TSH directly affects thyroid function, malfunction of the pituitary or the hypo - thalamus influences the thyroid gland activity. Disease in any portion of the thyroid-pitutary-hypothala- mus system may influence the levels of T3 and T4 in the blood. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels may be low. In addition, in the Euthyroid Sick Syndrome, multiple alterations in serum thyroid function test findings have been recognized in patients with a wide variety of non-thyroidal illnesses (NTI) without evidence of preexisting thyroid or hypothalami c-pitutary diseases. Thyroid Binding Globulin (TBG) concentrations remain relatively constant in healthy individuals. However, pregnancy, excess estrogen's, androgen's, antibiotic steroids and glucocorticoids are known to alter TBG levels and may cause false thyroid values for Total T3 and T4 tests.

TSH	T4	T3	INTERPRETATION	
High	Normal	Normal	Mild (subclinical) hypothyroidism	
High	Low	Low or normal	Hypothyroidism	
Low	Normal	Normal	Mild (subclinical) hyperthyroidism	
Low	High or normal	High or normal	Hyperthyroidism	
Low	Low or normal	Low or normal	Nonthyroidal illness; pituitary (secondary) hypothyroidism	
Normal	High	High	Thyroid hormone resistance syndrome (a mutation in the thyroid hormone receptor decreases thyroid hormone function)	









NABL & ISO APPROVED



 Patient Name
 : Mr JAI RAM RAJANI
 Bill Date
 : Jul 21, 2022, 03:47 PM

 DOB/Age/Gender
 : 63Yrs/Male
 Sample Collected : Jul 21, 2022, 03:11 PM

 Patient ID
 : 1085133
 Sample Received : Jul 21, 2022, 03:48 PM

 Referred by
 : Dr.
 Report Date
 : Jul 21, 2022, 11:04 PM

Sample Type : Spot Urine BarcodeNo : CP163878
Client : STANDARD KANPUR Report Status : Final Report

Test Description Value(s) Unit(s) Reference Range

CLINICAL PATHOLOGY REPORT Lifecare Health Gold Plus Package Urine Routine & Microscopic Examination

PHYSICAL EXAMINATON

Volume	20	mL			
Colour	Pale yellow				
Transparency	Slightly Hazy		Clear		
Deposit	Absent				
CHEMICAL EXAMINATION					
Reaction (pH) Method : Double Indicator	6.0		5.5-8.0		
Specific Gravity Method : Ion Exchange	1.015	-	1.010 - 1.030		
Urine Glucose (sugar)	Negative		Negative		
Urine Protein (Albumin)	Negative	-	Negative		
Urine Ketones (Acetone)	Negative	-	Negative		
Blood	Negative		Negative		
Leucocyte esterase	Negative	-	Negative		
Bilirubin Urine	Negative		Negative		
Nitrite Method : Griless Test	Negative		Negative		
Urobilinogen Method : Ehrlichs Test	Normal		Normal		
MICROSCOPIC EXAMINATION					
Pus Cells (WBCs)	2-4	/hpf	0 - 5		
Epithelial Cells	1-2	/hpf	0 - 4		
Red blood Cells	Absent	/hpf	Absent		
Crystals	Absent		Absent		
Cast	Absent		Absent		
Yeast Cells	Absent		Absent		
Amorphous deposits	Present		Absent		
Bacteria	Absent		Absent		
Protozoa	Absent		Absent		









CONDITIONS OF REPORTING

- 1. It is Presumed that specimen belongs to patient named or identified, such verification being carried out at the point of generation of said specimen
- 2. A test might not be performed due to following reason:
- Specimen Quantity not sufficient (Inadequate collection/spillage during transit)
- Specimen Quality not acceptable (Hemolysis/clotted/lipemic.)
- Incorrect sample type
- Test cancelled either on request of patient or doctor

In any of the above case a fresh specimen will be required for testing and reporting

- 3. The results of the tests may vary from lab to lab; time to time for the same patient
- 4. The reported results are dependent on individual assay methods, equipment, method sensitivity, specificity and quality of the specimen received
- 5. Partial representation of report is not allowed
- 6. The reported tests are for the notification of the referring doctor, only to assist him/her in the diagnosis and management of the patient
- 7. If Sample collection date is not stated on test requisition form, the current date will be printed by default as the date of collection.
- 8. Report with status "Preliminary" means one or more test are yet to be reported
- 9. This report is not valid for Medico Legal Purpose
- 10. Applicable Jurisdiction will be of "Delhi" for any dispute/claim concerning the test(s) & results of the test (s)