

Started on	Sunday, 12 February 2023, 8:28 PM
State	Finished
Completed on	Sunday, 12 February 2023, 8:28 PM
Time taken	10 secs
Marks	0.00/205.00
Grade	0.00 out of 100.00



Question 1

Not answered

Marked out of 5.00

Psychological mechanisms

Identify one central mechanism involved for each of the following situations;

Tom is a seven-year-old boy who is told to finish his homework before he can go and play with his mates

Choose... ▾

Miss. Y is a 23-year-old woman who has a history of spider phobia. She is put into a relaxed state by soothing music, suggestion and visualizing pleasant images. While she is in a relaxed state, she is shown a picture of a spider

Choose... ▾

Rosy is a 6-year-old girl who usually plays with her dog without fear. One day the dog accidentally bites her. She then becomes fearful of approaching the pet.

Choose... ▾

Lucy is a 7-year-old girl who is seeing a speech therapist for mutism. She only speaks in the presence of the therapist but not others

Choose... ▾

Sam is a 3-year-old boy, who is learning to write. His mother appreciates him for holding his pencil correctly, then for scribbling something on the paper and later for writing something, which resembles like a letter of the alphabets.

Choose... ▾

Your answer is incorrect.

Explanation:

- 1) This is Premack principle, which states that high-frequency behaviour can be used to reinforce a low-frequency behaviour. This is also called as Grandma's rule - eat the spinach to get the dessert. It is nothing but positive reinforcement but instead of a material reinforcer, an already existing behaviour itself (eating dessert) is linked to desired response (eating spinach).
- 2) The answer is reciprocal inhibition, where a stimulus with desired response and stimulus with the undesired response are presented together repeatedly; then incompatibility leads to reduced undesired response.
- 3) It is classical conditioning, which is a type of associative learning, and it is a relatively rapid process. In discriminative learning- a behaviour that is reinforced only in certain situations leads to selective expression of the behaviour in only those selected situations.

4) Discrimination is a process where learned responses are made only to specific stimuli and not to other similar stimuli.

5) It is shaping that is a form of operant conditioning in which a desirable behaviour pattern is learnt by the successive reinforcement of behaviours closer to the desired one.

The correct answer is: Tom is a seven-year-old boy who is told to finish his homework before he can go and play with his mates → Premack principle, Miss. Y is a 23-year-old woman who has a history of spider phobia. She is put into a relaxed state by soothing music, suggestion and visualizing pleasant images. While she is in a relaxed state, she is shown a picture of a spider → Reciprocal inhibition, Rosy is a 6-year-old girl who usually plays with her dog without fear. One day the dog accidentally bites her. She then becomes fearful of approaching the pet. → Classical conditioning, Lucy is a 7-year-old girl who is seeing a speech therapist for mutism. She only speaks in the presence of the therapist but not others → Discriminative learning, Sam is a 3-year-old boy, who is learning to write. His mother appreciates him for holding his pencil correctly, then for scribbling something on the paper and later for writing something, which resembles like a letter of the alphabets. → Shaping



Question 2

Not answered

Marked out of 5.00

Psychological Interventions

Choose one appropriate intervention from the given list for each situation below:

An 8-year-old girl suffers from bed-wetting. She does not have any depressive features or anatomical lesions. Her cystourethrogram has been normal. Her parents are not keen on pharmacotherapy.

Choose... ▾

A 50-year-old man with longstanding paranoid schizophrenia is living in a warden controlled community dwelling. His brother who has been his primary carer has suddenly died of a heart attack three weeks ago and since then the patient is withdrawn, tearful, and sleepless and has been talking about death.

Choose... ▾

A 32-year-old recently married woman working as a schoolteacher has had a miscarriage. She feels upset and worried about her husband's disappointment.

Choose... ▾

A 45-year-old patient with schizophrenia has been on the depot for last two years, but this has to be stopped as he is developing Parkinsonian symptoms. He has a history of poor adherence with oral medications in the past, and his mental health team is reluctant to change his prescription to oral therapy again.

Choose... ▾

A 39-year-old woman with a history of low mood, low self-confidence and poor self-esteem. She is not keen to take any medications.

Choose... ▾

Your answer is incorrect.

Explanation:

- 1) Behavioural therapy using urine alarm or awarding stars (reinforcement) can help enuresis.
- 2) For a well-established severe mental illness currently facing a crisis due to internal or external factors, crisis intervention can be timely.
- 3) Counselling may be useful where there is no serious identified mental illness, but psychological support is necessary nevertheless.
- 4) Compliance therapy may be helpful as a rehabilitation measure in those with long-standing schizophrenia and poor adherence.

5) For mild to moderate depression, CBT should be considered as the first-line treatment of choice.

The correct answer is: An 8-year-old girl suffers from bed-wetting. She does not have any depressive features or anatomical lesions. Her cystourethrogram has been normal. Her parents are not keen on pharmacotherapy. → Behavioural therapy, A 50-year-old man with longstanding paranoid schizophrenia is living in a warden controlled community dwelling. His brother who has been his primary carer has suddenly died of a heart attack three weeks ago and since then the patient is withdrawn, tearful, and sleepless and has been talking about death. → Crisis intervention, A 32-year-old recently married woman working as a schoolteacher has had a miscarriage. She feels upset and worried about her husband's disappointment. → Counselling, A 45-year-old patient with schizophrenia has been on the depot for last two years, but this has to be stopped as he is developing Parkinsonian symptoms. He has a history of poor adherence with oral medications in the past, and his mental health team is reluctant to change his prescription to oral therapy again. → Compliance therapy, A 39-year-old woman with a history of low mood, low self-confidence and poor self-esteem. She is not keen to take any medications. → Cognitive behavioural therapy



Question 3

Not answered

Marked out of 5.00

Terminologies in psychotherapy

For each of the following statements, describe one of the options listed below;

I don't get upset with my patients usually. I am treating this patient who makes me furious, and I can't understand why I feel so irritated with her

Choose...

There are only 2 Options for my problems. I am either going to be on my own and isolated, or I get involved in a relationship and feel overwhelmed by it. I am confused, and I don't think I can win either way.

Choose...

I want to tell my wife that I am bisexual and can stop hiding important parts of my life with her. But I am worried that the shock might kill her and so I can't.

Choose...

I always give in to the whims of others to avoid any guilt feelings. But this also makes me very angry and then the guilt factor sets in. I end up finally feeling so guilty that I become even more flexible and accommodating to others, which upsets me again.

Choose...

This term refers to the range of tasks that children cannot yet accomplish on their own but can do with the help of adults or other children

Choose...

Your answer is incorrect.

Explanation:

- 1) Countertransference refers to the therapist's attitudes towards the patient. Any strong emotions a therapist feels towards a patient can originate from the therapist's past, his/her conflicts or unresolved problems and may be inappropriately transferred onto the patient.
- 2) Dilemmas, Snags and Traps are terms used in Interpersonal therapy. When a person acts as though available action or possible roles were limited to polarized alternatives, this is referred to as a dilemma.
- 3) An individual is said to be in a snag when appropriate roles are abandoned because the person makes an assumption that others would oppose them or because they are perceived as forbidden or dangerous.

4) Traps are repetitive cycles where consequences of the behaviour feedback into its perpetuation. Here negative assumptions generate acts that produce consequences that reinforce the assumptions.

5) The zone of proximal development was proposed by Vygotsky. It refers to the range of tasks that children cannot yet accomplish on their own but can do with the help of adults or other children. Zone of proximal development refers to functions that are not yet fully achieved but are in the process 'pipeline'. While actual achievement measures development after completion retrospectively, ZPD has a prospective sense as to what the child will be achieving in near future.

The correct answer is: I don't get upset with my patients usually. I am treating this patient who makes me furious, and I can't understand why I feel so irritated with her → Countertransference, There are only 2 Options for my problems. I am either going to be on my own and isolated, or I get involved in a relationship and feel overwhelmed by it. I am confused, and I don't think I can win either way. → Dilemma, I want to tell my wife that I am bisexual and can stop hiding important parts of my life with her. But I am worried that the shock might kill her and so I can't. → Snag, I always give in to the whims of others to avoid any guilt feelings. But this also makes me very angry and then the guilt factor sets in. I end up finally feeling so guilty that I become even more flexible and accommodating to others, which upsets me again. → Trap, This term refers to the range of tasks that children cannot yet accomplish on their own but can do with the help of adults or other children → Zone of proximal development



Question 4

Not answered

Marked out of 5.00

Pioneers in therapy models (1)

Match each therapy below with its pioneer using the list of options:

Cognitive Behavioural Therapy	Choose... ▾
Cognitive analytical therapy	Choose... ▾
Dialectical behavioural therapy	Choose... ▾
Interpersonal therapy.	Choose... ▾
Mentalization based treatment	Choose... ▾

Your answer is incorrect.

Explanation:

- 1) The Proponent of CBT is Aaron Beck.
- 2) Cognitive analytical therapy, which contains aspects of both cognitive therapy and psychoanalytic psychotherapy, was developed by Anthony Ryle
- 3) The proponent of DBT is Marsha Linehan who invented it to reduce self-harm in borderline patients.
- 4) The proponents of Interpersonal therapy are - Klerman & Weissman, 1993.
- 5) Mentalization based treatment for Borderline PD started with the psychoanalytically oriented partial hospitalization programme with the standard psychiatric care that is now fully materialized as a mentalization-based treatment (Bateman & Fonagy, 2004).

The correct answer is: Cognitive Behavioural Therapy → Aaron Beck, Cognitive analytical therapy → Anthony Ryle, Dialectical behavioural therapy → Marsha Linehan, Interpersonal therapy. → Klerrmann and Weissmann, Mentalization based treatment → Bateman & Fonagy

Question 5

Not answered

Marked out of 5.00

Pioneers in therapy models (2)

Match each pioneer below with its therapy above;

Franz Alexander & Thomas French

Choose...



Miller & Rollnick

Choose...



Shapiro

Choose...



Albert Ellis

Choose...



Wolpe

Choose...



Your answer is incorrect.

Explanation:

- 1) In 1946, Franz Alexander and Thomas French identified the essential characteristics of brief psychodynamic psychotherapy. Since then, many workers such as David Malan in England, Peter Sifneos in the United States, and Habib Davanloo in Canada have influenced the field.
- 2) Motivational Interviewing (Miller & Rollnick, 1991) is often used in managing alcohol and substance misuse. In line with Roger's client centred therapy, Miller and Rollnick did extensive work with substance users and developed motivational interviewing.
- 3) Eye movement desensitisation and reprocessing were serendipitously discovered by a psychologist called Shapiro when she first applied it to herself. It is based on the hypothesis that bilateral stimulation, in the form of eye movements, allows the processing of traumatic memories. During the therapy, the patient focuses on particular images, negative sensations and associated cognitions. Bilateral stimulation, in the form of eye movements, is applied to desensitise the individual to these memories. More positive feelings and cognitions are introduced.
- 4) Albert Ellis developed Rational Emotive Therapy in 1955. It is a reality-based, short-term therapy aimed at cognitive restructuring with many shared features with contemporaneous CBT.

5) The proponent of systematic desensitization is Wolpe. The therapy is based on the behavioral principle of counterconditioning (i.e. gradual approach of feared situation in a psychophysiological state that inhibits anxiety leads to a reduction of anxiety response.)

The correct answer is: Franz Alexander & Thomas French → Brief psychodynamic psychotherapy, Miller & Rollnick → Motivational interviewing, Shapiro → Eye movement desensitization reprocessing, Albert Ellis → Rational-emotive therapy, Wolpe → Systematic desensitization



Question 6

Not answered

Marked out of 5.00

Family therapy

For the definitions given below, identify the terms commonly used in family therapy

A pattern of relationship in which a third person is recruited to bypass the conflict between two people thus stabilizing the link between the former two.

Loss of autonomy due to a blurring of psychological boundaries and becoming emotionally over-involved

Interdependence of action and reaction (X leads to Y leads to X), as opposed to linear causality that implies X leads to Y

The therapist makes preliminary attempts to explain the patient's symptoms in terms of the contexts within which they occur

Blaming a convenient but innocent person for the problems among other members of the family

Your answer is incorrect.

Explanation:

- 1) Triangulation is a pattern of relationship in which a third person is recruited to bypass the conflict between 2 people thus stabilizing the link between the original two people.
- 2) Enmeshment is the loss of autonomy due to a blurring of psychological boundaries: emotionally too close or over involved. On the contrary, Disengagement is the term used to describe emotionally uninvolved or too distant state leading to psychological isolation.
- 3) Circularity is the term used to define interdependence of action and reaction (X leads to Y leads to X), as opposed to linear causality that implies X leads to Y.
- 4) The hypothesis is an initial supposition about family functioning that can be tested out in the session and is the starting point for further investigation.
- 5) Scapegoating is blaming a convenient but innocent person for problems and conflicts among other members of the family.

The correct answer is: A pattern of relationship in which a third person is recruited to bypass the conflict between two people thus stabilizing the link between the former two. → Triangulation, Loss of autonomy due to a blurring of psychological boundaries and becoming emotionally over-involved → Enmeshment, Interdependence of action and reaction (X leads to Y leads to X), as opposed to linear causality that implies X leads to Y → Circularity, The therapist makes preliminary attempts to explain the patient's symptoms in terms of the contexts within which they occur → Hypothesis, Blaming a convenient but innocent person for the problems among other members of the family → Scapegoating



Question 7

Not answered

Marked out of 5.00

Principles of Psychotherapy

Using the descriptions given below, identify the type of psychotherapy

Focuses mainly on the present functioning of the family rather than its past experiences. Strategies include employing paradoxical injunctions.

Choose...

Focuses on a set of unspoken rules that organize ways in which family members relate to one another

Choose...

This therapy is based on the principle that current problems in the family originate from past experiences of its family members

Choose...

This therapy uses cognitive therapy techniques with a psychodynamic framework

Choose...

The focus areas in this therapy are transitions, disputes, grief and sensitivity

Choose...

Your answer is incorrect.

Explanation:

1) The systemic family therapy focuses on the present functioning of the family rather than past experiences. It uses paradoxical injunctions where the therapist makes the patient intentionally engage in the unwanted behaviour e.g. avoid a phobic object or perform a compulsive ritual. This counterintuitive approach can provide new insights for some patients. Other terms associated with this therapy includes circularity, hypothesizing, neutrality, paradox and counter paradox.

2) The structural model (Minuchin) views family as a structure built of interpersonal relationships. Such a structure will have a hierarchy of power (parents in charge of children), firm boundaries and set of unspoken rules. Families function particularly well when these family structures are intact. Disrupted structure leads to disrupted communication.

3) Psychodynamic therapy is based on the principle that current problems in the family originate from past experiences of its family members.

4) The cognitive analytical therapy contained aspects of both cognitive therapy, psychoanalytic psychotherapy and developed by Anthony Ryle.

5) In interpersonal therapy, emotional problems are best understood by studying the interpersonal context in which they arise. Four interpersonal problem areas are addressed: grief, interpersonal role disputes, role transitions and interpersonal deficits.

The correct answer is: Focuses mainly on the present functioning of the family rather than its past experiences. Strategies include employing paradoxical injunctions. → Systemic family therapy, Focuses on a set of unspoken rules that organize ways in which family members relate to one another → Structural family therapy, This therapy is based on the principle that current problems in the family originate from past experiences of its family members → Psychodynamic therapy, This therapy uses cognitive therapy techniques with a psychodynamic framework → Cognitive analytical therapy, The focus areas in this therapy are transitions, disputes, grief and sensitivity → Interpersonal therapy



Question 8

Not answered

Marked out of 5.00

Developments in psychotherapy

With the descriptions given below, identify the proponents;

Expanded drive theory to include non-sexual drives

Choose... ▾

Introduced methods of play therapy

Choose... ▾

Proposed that gratification is obtained through relationships as well as through satisfaction of one's desires

Choose... ▾

Examination of games and scripts that characterize relationships

Choose... ▾

Every individual has the motivation and ability to change, and he or she is the best person to decide on the direction of change.

Choose... ▾

Your answer is incorrect.

Explanation:

- 1) Jung expanded the drive theory to include drives other than libido. He also introduced ideas of 'collective unconscious' and personality archetypes. Jung is also associated with concepts such as extroversion and introversion.
- 2) Melanie Klein introduced methods of play therapy. She is also associated with theories of childhood development including primitive defense mechanisms such as 'splitting'.
- 3) Donald Winnicott was one of the proponents of the object relations theory. He argued that gratification could be obtained through relationships as well as through satisfaction of desires.
- 4) Eric Berne is associated with the transactional analysis. He emphasized the importance of therapeutic attributes of genuineness, accurate empathy and unconditional positive regard.

5) Client-centered psychotherapy is associated with Carl Rogers. This therapy is based on the principle that every individual has the motivation and ability to change, and he or she is the best person to decide on the direction of change. Client-centered therapy is nondirective as the therapists avoid directing the therapeutic process. It emphasizes the importance of 'therapeutic attributes of genuineness', 'unconditional positive regard' and 'accurate empathy'.

The correct answer is: Expanded drive theory to include non-sexual drives → Jung, Introduced methods of play therapy → Klein, Proposed that gratification is obtained through relationships as well as through satisfaction of one's desires → Winnicott, Examination of games and scripts that characterize relationships → Berne, Every individual has the motivation and ability to change, and he or she is the best person to decide on the direction of change. → Carl Rogers



Question 9

Not answered

Marked out of 5.00

Proponents of analytical models (1)

For each of the description below, select the most appropriate historical proponent from the list:

Basic assumptions	Choose... ▾
Collective unconscious	Choose... ▾
Formulation Matrix	Choose... ▾
Good enough mother	Choose... ▾
Paranoid/schizoid position	Choose... ▾

Your answer is incorrect.

Explanation:

- 1) Bion postulated that whenever a group gets derailed from its task, it deteriorates into one of three primary states, that are also called basic assumptions. It includes dependency, pairing, or fight-flight. These basic assumptions must be tackled to allow group work to proceed.
- 2) Jung founded analytic psychology and coined the term collective unconscious to describe all of the mankind's collective symbolic past.
- 3) The formulation matrix in group therapy was described by Foulkes. It refers to a hypothetical web of communication and relationship in a given group. It is the common shared ground, which ultimately determines the meaning and significance of all events and upon which all communication and interpretations, verbal and nonverbal rest.
- 4) 'Good enough mother' is a concept proposed by Winnicott. It refers to the fact that a mother need not be perfect - but good enough to provide a growth-sustaining environment (holding).
- 5) Melanie Klein described two positions in her object relations theory, namely paranoid-schizoid and depressive position. The paranoid-schizoid position occurs during first three months of life. It is the result of an infant's attempt to master its fear and anxiety immediately after birth. This position concerns an inability to perceive objects as a whole object but splitting them into their good and bad parts. Infant

experiences mother as two separate objects; good object which feeds, nurtures and a bad object which is withholding, harsh when not available to him This is associated with the use of 'splitting' and 'projection' as defence mechanisms.

The correct answer is: Basic assumptions → Bion, Collective unconscious → Jung, Formulation Matrix → Foulkes, Good enough mother → Winnicott, Paranoid/schizoid position → Klein



Question 10

Not answered

Marked out of 4.00

Psychological treatments (1)

Choose one most appropriate treatment for each description below:

A woman who was previously treated for anorexia now presents with binge eating and induced vomiting.

Choose...



An Iraq war veteran is plagued by recurrent nightmares and not able to sleep well. He avoids watching news channels that report war incidents.

Choose...



A 24-year-old lady who self-harms repeatedly requires intensive community team input.

Choose...



A 21-year-old woman with fear of spiders shows a tendency to avoid any objects that are even remotely connected to spiders

Choose...



Your answer is incorrect.

Explanation:

- 1) This woman has bulimia, which is not uncommon in those who had anorexia nervosa in the past. Interpersonal therapy (IPT) has been established as an effective treatment for bulimia. The IPT does not focus on bulimic symptoms but focuses on interpersonal problems that become the focus of treatment.
- 2) The diagnosis is post-traumatic stress disorder. Shapiro developed EMDR or Eye Movement Desensitization and Reprocessing in 1989.
- 3) The diagnosis is borderline personality disorder. Dialectic behaviour therapy was introduced to reduce self-harm in borderline patients by Marsha Linehan. The four modes of treatment in DBT are as follows: group skills training, individual therapy, phone consultations, and consultation team.
- 4) The diagnosis is spider phobia, which is a form of simple phobia. The treatment of choice is systematic desensitisation, which is based on the principles of counter conditioning. It consists of 3 steps; relaxation training, hierarchy construction and desensitisation of the stimulus. The patient is exposed to a graded hierarchy of situations in a step-wise fashion.

The correct answer is: A woman who was previously treated for anorexia now presents with binge eating and induced vomiting. → Interpersonal therapy, An Iraq war veteran is plagued by recurrent nightmares and not able to sleep well. He avoids watching news channels that report war incidents. → Eye Movement Desensitization and Reprocessing, A 24-year-old lady who self-harms repeatedly requires intensive community team input. → Dialectic behaviour therapy, A 21-year-old woman with fear of spiders shows a tendency to avoid any objects that are even remotely connected to spiders → Systematic desensitisation



Question 11

Not answered

Marked out of 5.00

Group therapy (2)

Identify the types of groups that are particularly useful for each of the following conditions

Carers of people with Alzheimer's dementia	Choose... ▾
Drug dependency	Choose... ▾
Phobia Therapy	Choose... ▾
Alcohol dependence	Choose... ▾
Sexual deviancy	Choose... ▾

Your answer is incorrect.

Explanation:

Support groups are aimed at providing peer support for carers and family of those with a learning disability, chronic illness such as dementia.

Problem-focused groups are helpful for client groups facing a specific behavioural problem such as alcohol dependence, drug dependency, sexual deviancy.

Behavioural groups are useful in treating phobia.

Others groups include activity groups, which are used for patients who are unsuitable for other group activities. Activities may include vocations such as art, computing and gardening. It is mostly used in learning disability, chronic psychosis, and other disorders with functional impairment. The aim of psychodynamic groups is lasting change through exploratory therapy.

The correct answer is: Carers of people with Alzheimer's dementia → Support groups, Drug dependency → Problem focused groups, Phobia Therapy → Behavioural groups, Alcohol dependence → Problem focused groups, Sexual deviancy → Problem focused groups

Question 12

Not answered

Marked out of 4.00

Psychotherapeutic concepts (4)

For each of the description below, identify the psychoanalytical concept from the list below

Displacement of feelings and attitudes derived from past relationships and experiences onto the therapist

Choose...

This refers to the expression of therapist's understanding of the meaning of feelings, attitudes, defense mechanisms and behaviours currently exhibited during therapy.

Choose...

A particular reaction that occurs in the analytic setting when the analyst expresses to the patient that he or she is making significant progress in treatment, and there appears to be a marked increase in the patient's symptoms

Choose...

This refers to any hindrance noted in the course of therapeutic progress during psychotherapy

Choose...

Your answer is incorrect.

Explanation:

1) Transference refers to a displacement of feelings and attitudes derived from past relationships and experiences onto the therapist. It is bidimensional - includes replaying past experiences + seeking a new relationship with the therapist. It may involve idealizing, erotic or highly negative and denigrating approaches. Countertransference can be regarded as the transference from an analyst on to the patient due to the past experiences of the analyst. Initially viewed as a resistance or obstacle to therapy, this is now regarded to be due to both analysts' past experiences and also the reaction of an analyst to what the patient projects onto him/her. Hence analyzing countertransference can provide insight into a patient's psychic state in the same sense as analyzing transference.

2) The term interpretation refers to the expression of therapist's understanding of the meaning of feelings, attitudes, defense mechanisms and behaviours currently exhibited during therapy.

3) The NTR - negative therapeutic reaction (Freud 1923) describes a particular reaction that occurs in the analytic setting when the analyst expresses to the patient that he or she is making significant progress in therapy. When such a comment is made, the patient's defenses are mobilized to protect him/her from any positive therapeutic gain, and there appears to be a marked increase in the patient's symptoms; i.e.

rather than the patient benefiting from the analyst's expressed satisfaction, the patient's condition worsens.

4) Resistance refers to any hindrance noted in the course of therapeutic progress during psychotherapy. Repression resistance refers to patient's difficulty in gaining access to certain ideas and emotions. Transference resistance refers to patient's unconscious wish to keep therapeutic relationship similar to past relationships.

The correct answer is: Displacement of feelings and attitudes derived from past relationships and experiences onto the therapist → Transference, This refers to the expression of therapist's understanding of the meaning of feelings, attitudes, defense mechanisms and behaviours currently exhibited during therapy. → Interpretation, A particular reaction that occurs in the analytic setting when the analyst expresses to the patient that he or she is making significant progress in treatment, and there appears to be a marked increase in the patient's symptoms → Negative therapeutic reaction, This refers to any hindrance noted in the course of therapeutic progress during psychotherapy → Resistance



Question 13

Not answered

Marked out of 5.00

Cognitive elements & disorders

For each of the cognitive elements given below, identify the most relevant psychiatric disorder associated with it from list below;

Negative view of self, world and future

Choose... ▾

Catastrophic misinterpretation of physiological experiences

Choose... ▾

Alternate action or substitute action can undo or compensate for another thought or action

Choose... ▾

Negative global external attributional bias

Choose... ▾

Guilt and self-blame with feelings of loss of control

Choose... ▾

Your answer is incorrect.

Explanations:

- 1) The cognitive element in depressive disorder includes a negative view of self/past (worthlessness), the world/present (helplessness) and future (hopelessness).
- 2) In panic disorder, catastrophic misinterpretation of normal physiological experiences can occur.
- 3) The cognitive elements related to OCD include the notion that thoughts are as powerful as actions so that an alternative or substitute action can undo or compensate for another thought or action (Undoing).
- 4) Negative global external attributional bias is the cognitive element seen in paranoid personality disorder that forms the basis of ideas of reference and persecution.
- 5) In PTSD, there is severe guilt and self-blame with feelings of loss of control.

The correct answer is: Negative view of self, world and future → Depression, Catastrophic misinterpretation of physiological experiences → Panic disorder, Alternate action or substitute action can undo or compensate for another thought or action → Obsessive compulsive

disorder, Negative global external attributional bias → Paranoid personality disorder, Guilt and self-blame with feelings of loss of control → Posttraumatic stress disorder



Question 14

Not answered

Marked out of 5.00

Cognitive profile of psychiatric disorders

For each of the cognitive content described below, identify the associated psychiatric disorder from the list

Hopelessness and cognitive constriction (one-way exit)	Choose... ▾
Fear of physical and psychological danger	Choose... ▾
Inflated view of self, life experience and future	Choose... ▾
Attribution of serious medical disorder	Choose... ▾
Misinterpretation of social threat	Choose... ▾

Your answer is incorrect.

Explanation:

- 1) The cognitive elements in suicidal behaviour often include hopelessness and cognitive constriction (one-way exit).
- 2) In generalized anxiety disorder, there is fear of physical and psychological danger with the misinterpretation of any situation as threatening.
- 3) In hypomania and mania, there is an inflated view of self, life experience and future.
- 4) In hypochondriasis, there is a non-catastrophic misinterpretation of any bodily sensations during information processing leading to attribution of serious medical disorders.
- 5) A catastrophic misinterpretation is seen in panic attacks. There is a misinterpretation of social threat in social phobia.

The correct answer is: Hopelessness and cognitive constriction (one-way exit) → Suicidal behaviour, Fear of physical and psychological danger → Generalised anxiety disorders, Inflated view of self, life experience and future → Hypomanic episode, Attribution of serious medical disorder → Hypochondriasis, Misinterpretation of social threat → Social Phobia



Question 15

Not answered

Marked out of 5.00

Cognitive errors

With the descriptions given below, identify the type of cognitive error the patient is harbouring:

A depressed schoolteacher thinks, "One student in the class is sleeping so the entire class is bored of my lecture."

Choose... ▾

A dysthymic student can talk only about his minor failures and withdrawals from college competitions, and chooses to ignore the A grades and accolades that he has received so far

Choose... ▾

A woman with PTSD says " Its my fault; I shouldn't have played that music when my husband was driving."

Choose... ▾

A woman with panic attacks can sense mild palpitation after some housework, and thinks she is going to have a heart attack and immediate death.

Choose... ▾

A depressed University student whose exam is scheduled for last week of March this year says "I did my exam on the last week of March last year too and failed, I am sure it is going to happen again".

Choose... ▾

Your answer is incorrect.

Explanation:

- 1) In this situation, the depressed schoolteacher is generalizing a small element to the whole (Overgeneralizing).
- 2) This dysthymic student selectively picks failures and ignores positive features (selective abstraction).
- 3) The woman with PTSD is assuming responsibility for all that has gone bad (personalization).
- 4) The woman with panic attacks is predicting the worst case scenario to happen, which is a massive heart attack and immediate death (catastrophic thinking).
- 5) The university student is predicting his future outcomes without sufficient evidence (arbitrary inference).

The correct answer is: A depressed schoolteacher thinks, "One student in the class is sleeping so the entire class is bored of my lecture." → Overgeneralization, A dysthymic student can talk only about his minor failures and withdrawals from college competitions, and chooses to

ignore the A grades and accolades that he has received so far → Selective abstraction, A woman with PTSD says "Its my fault; I shouldn't have played that music when my husband was driving." → Personalisation, A woman with panic attacks can sense mild palpitation after some housework, and thinks she is going to have a heart attack and immediate death. → Catastrophic thinking, A depressed University student whose exam is scheduled for last week of March this year says "I did my exam on the last week of March last year too and failed, I am sure it is going to happen again". → Arbitrary inference



Question 16

Not answered

Marked out of 4.00

Negative automatic thought

With the descriptions given below, identify the type of automatic negative thoughts:

A woman complains that her husband never takes her with him on business trips. She concludes that he is concealing an affair and ignores other possibilities

A junior psychiatric trainee goes for an interview for a senior training post. The interview panel was warm and friendly but despite this the trainee came away, and all they could remember was that one of the board members frowned. The trainee considers that this means she will not be successful.

A woman finds her son smoking. She is devastated as she believes this means he will be doing street drugs as well.

A trainee mentioned after the MRCPsych exam- I will either pass the MRCPsych exam, get the forensic ST4 job or I will fail the exam, have no training job forever."

Your answer is incorrect.

Explanation:

- 1) Arbitrary inference- Predicting without sufficient evidence.
- 2) Selective abstraction- the trainee selectively picks and focussed on the one negative and ignored the positives.
- 3) Overgeneralization - In this case, the mother described in the question has made a general conclusion based on a single event. Generalizing a small element to the whole is called overgeneralisation.
- 4) The trainee has dichotomous thinking or black and white thinking

The correct answer is: A woman complains that her husband never takes her with him on business trips. She concludes that he is concealing an affair and ignores other possibilities → Arbitrary inference, A junior psychiatric trainee goes for an interview for a senior training post. The interview panel was warm and friendly but despite this the trainee came away, and all they could remember was that one of the board members frowned. The trainee considers that this means she will not be successful. → Selective abstraction, A woman finds her son

smoking. She is devastated as she believes this means he will be doing street drugs as well. → Overgeneralization, A trainee mentioned after the MRCPsych exam- I will either pass the MRCPsych exam, get the forensic ST4 job or I will fail the exam, have no training job forever." → All or nothing thinking



Question 17

Not answered

Marked out of 3.00

Psychotherapy treatments (2)

Identify the treatment options that can be used in the following situations:

A 23-year-old man dependent on alcohol has cut down his drinking for the last three weeks and wanted to rebuild his life. He is in a new stable relationship.

Choose... ▾

A 20-year-old woman is living alone, reports traumatic childhood and significant self-harming behaviour.

Choose... ▾

A 25 years old lady, who is the primary carer for her mother who has dementia, reports mild depressive features.

Choose... ▾

Your answer is incorrect.

Explanation:

1) The current recommendation concerning psychological approaches in treating alcohol dependence is to offer cognitive behavioural therapies, behavioural therapies or social network and environment-based therapies. They are focused specifically on alcohol-related cognitions, behaviour, problems and social networks. Behavioural couples therapy is particularly recommended if service users have a regular partner who is willing to participate in treatment and are aiming for abstinence (as described in this question).

2) NICE CG133 recommends that for self-harm, providers must consider offering 3 to 12 sessions of a psychological intervention that is specifically structured with the aim of reducing self-harm. Also the intervention should be tailored to individual need, and could include cognitive behavioural, psychodynamic or problem-solving elements. Given the presence of childhood trauma, psychodynamic therapy is appropriate here.

3) For carers of dementia patients who have a negative psychological impact from caring, supportive psychotherapy is recommended initially (usually in group settings).

The correct answer is: A 23-year-old man dependent on alcohol has cut down his drinking for the last three weeks and wanted to rebuild his life. He is in a new stable relationship. → Behavioural couples therapy, A 20-year-old woman is living alone, reports traumatic childhood and significant self-harming behaviour. → Psychodynamic, A 25 years old lady, who is the primary carer for her mother who has dementia, reports mild depressive features. → Supportive psychotherapy

Question 18

Not answered

Marked out of 3.00

Defence mechanisms

For each of the definitions given below, identify the most likely defence mechanisms from the list below.

Involuntary but complete exclusion of painful reality from awareness

Choose...



Exclusion from conscious awareness of anxiety provoking ideas and or feelings so they become unconscious

Choose...



Reversion to the modes of psychological functioning that are characteristic of earlier life stages

Choose...



Your answer is incorrect.

Explanation:

1) Denial: It is the apparent refusal to acknowledge a threatening reality. It may persist despite all of the relevant facts being consistently pointed out. It is not same as conscious avoidance of painful topics or thoughts.

2) Repression: This refers to withholding an idea or feeling from consciousness. Primary repression is the withholding of ideas and feelings before they have attained consciousness: secondary repression is withholding from awareness the ideas/feelings that were previously experienced at a conscious level. Note that this differs from suppression - suppression is mere postponement not the loss of thoughts from conscious perception. Repression is the primary defense. Other defences reinforce it.

3) Regression: Moving back into childish or earlier developmental phase to avoid confronting a conflict. Regression is also considered an essential concomitant of the creative process.

The correct answer is: Involuntary but complete exclusion of painful reality from awareness → Denial, Exclusion from conscious awareness of anxiety provoking ideas and or feelings so they become unconscious → Repression, Reversion to the modes of psychological functioning that are characteristic of earlier life stages → Regression



Question 19

Not answered

Marked out of 3.00

Psychodynamic psychotherapy

Choose ONE of the above terms best suited for each description below:

During psychodynamic therapy, a patient and her therapist identify the theme of inappropriately strict and sometimes abusive upbringing by the patient's father as relevant to her current condition. But despite repeated attempts of gentle probing from the therapist, the patient refuses to talk about her father directly.

Choose...

Following some sessions with the above patient who reminds the therapist of his difficult teenage daughter, the therapist says "be extra careful with your new boyfriend". During the middle phase of a year-long planned psychotherapy, a patient starts losing the considerable therapeutic gains made during initial part of the therapy without any apparent stressor in her life.

Choose...

A particular reaction that occurs in the analytic setting when the analyst expresses to the patient that he or she is making significant progress in therapy, and there appears to be a marked increase in the patient's symptoms

Choose...

Your answer is incorrect.

Explanation:

- 1) Resistance is a term used to describe the various ways in which patients in psychotherapy oppose the process of change.
- 2) The broad conceptualization of counter-transference is intended to include all the psychological and emotional reactions of the therapist toward the patient. Some individuals in psychoanalytic treatment manifest negative therapeutic responses, in which insights that ought to lead to symptomatic improvement produce worsening instead.
- 3) The NTR - negative therapeutic reaction (Freud 1923) describes a particular reaction that occurs in the analytic setting when the analyst expresses to the patient that he or she is making significant progress in therapy. When such a comment is made, the patient's defenses are mobilized to protect him/her from any positive therapeutic gain, and there appears to be a marked increase in the patient's symptoms; i.e. rather than the patient benefiting from the analyst's expressed satisfaction, the patient's condition worsens

The correct answer is: During psychodynamic therapy, a patient and her therapist identify the theme of inappropriately strict and sometimes abusive upbringing by the patient's father as relevant to her current condition. But despite repeated attempts of gentle probing from the therapist, the patient refuses to talk about her father directly. → Resistance, Following some sessions with the above patient who reminds the therapist of his difficult teenage daughter, the therapist says "be extra careful with your new boyfriend". During the middle phase of a year-long planned psychotherapy, a patient starts losing the considerable therapeutic gains made during initial part of the therapy without any apparent stressor in her life. → Countertransference, A particular reaction that occurs in the analytic setting when the analyst expresses to the patient that he or she is making significant progress in therapy, and there appears to be a marked increase in the patient's symptoms → Negative therapeutic reaction



Question 20

Not answered

Marked out of 4.00

Couples therapy

Identify the type of treatment using the descriptions given below;

This therapy employs communication training, reciprocity negotiation, structural moves requiring disagreement, reversed role-play and the use of paradox.

Choose...

The aim of this therapy is to help each partner understand his/her emotional needs and how they relate to those of the other.

Choose...

Tom is alcohol dependent and is seen for this therapy with his wife, Sally. They arrange a daily 'sobriety contract' in which Tom states his intent is not to drink, and Sally expresses her support for Tom's effort to remain abstinent.

Choose...

This therapy focuses on hidden rules and on disagreements about who makes these rules and on inconsistencies between these two levels of interaction.

Choose...

Your answer is incorrect.

Explanation:

- 1) Behavioural systems couple therapy was developed by Crowe. It has two sets of components. The behavioural components are reciprocity negotiating and training in communication. The systems components are structural moves (include requiring disagreement, reversed role play & sculpting, timetables & tasks) and the use of paradox. Treatments usually last for 5-10 sessions over 3-6 months.
- 2) Psychodynamic couples therapy: The aim of this treatment is to help each partner understand his/her emotional needs and how they relate to those of the other. It could be done in many ways. One therapist may see the couple together, two therapists may see them together, or 2 therapists may see the patients separately but meet regularly to coordinate their treatment.
- 3) Behavioural couples therapy is a brief, but highly structured therapy based on the principles of operant conditioning. The therapist tries to identify ways in which undesired behaviour between the couple is being reinforced unwittingly by one of its consequences. Each partner is asked to say what alternative actions they desire in the other, described in concrete terms. Each partner agrees a way of rewarding the other when the desired behaviour is carried out.

4) In systems approach to couple therapy the focus of treatment is on hidden rules that govern the behaviour of the couple towards one another, on disagreements about who makes these rules and on inconsistencies between these two levels of interaction. Important concepts include enmeshment, which is excessive involvement with the other person and the idea of a cycle of 'cause and effect' such that neither person is whole to blame. These ideas are discussed in conflicts arising in the everyday life of the couple, and it is hoped to arrive at a more balanced and cooperative relationship.

(Ref: The Shorter Oxford Textbook of Psychiatry 6th ed Pg 754-756)

The correct answer is: This therapy employs communication training, reciprocity negotiation, structural moves requiring disagreement, reversed role-play and the use of paradox. → Behavioural systems couple therapy, The aim of this therapy is to help each partner understand his/her emotional needs and how they relate to those of the other. → Psychodynamic couple therapy, Tom is alcohol dependent and is seen for this therapy with his wife, Sally. They arrange a daily 'sobriety contract' in which Tom states his intent is not to drink, and Sally expresses her support for Tom's effort to remain abstinent. → Behavioural couples therapy, This therapy focuses on hidden rules and on disagreements about who makes these rules and on inconsistencies between these two levels of interaction. → Systems approach to couple therapy



Question 21

Not answered

Marked out of 4.00

Cognitive elements of CBT

For each of the cognitive elements given below, identify the most relevant psychiatric disorder associated with it from list below:

Catastrophic misinterpretation of physiological experiences

Choose... ▾

Alternate action or substitute action can undo or compensate for another thought or action

Choose... ▾

Negative global external attributional bias

Choose... ▾

Misinterpretation of any situation as overly threatening

Choose... ▾

Your answer is incorrect.

Explanations:

In panic disorder and somatization, catastrophic misinterpretation of normal bodily symptoms can occur.

The cognitive elements related to OCD include the notion that thoughts are as powerful as actions so that an alternative or substitute action can undo or compensate for another thought or action (Undoing).

Negative global external attributional bias refers to the tendency to repeatedly attribute all negative events to external factors or other people. This is seen in paranoid personality disorder.

In anxiety, the self is seen as vulnerable, the world dangerous and the future uncertain. Schemas associated with vulnerability are seen to be of evolutionary usefulness in energising the organism and preparing for fight or flight from danger (Beck 1995). In anxiety disorders, these schemas are activated inappropriately in non-threatening situations. Any danger signals in the internal or external environment become predominant and reinforce the fear.

The correct answer is: Catastrophic misinterpretation of physiological experiences → Panic disorder, Alternate action or substitute action can undo or compensate for another thought or action → Obsessive compulsive disorder, Negative global external attributional bias →

Paranoid personality disorder, Misinterpretation of any situation as overly threatening → Generalised anxiety disorder



Question 22

Not answered

Marked out of 4.00

Therapy options

For each patient, select the one most appropriate therapeutic option.

A young woman develops anxiety, recurrent flashbacks and incapacitating fear of driving after having a dreadful accident.

Choose...

A 25-year-old woman presents with high-frequency self-harm behaviour. She says she cannot trust people anymore as none of her previous therapists offered her anything useful. They transferred her care back to her community psychiatrist whenever her self-harming behaviour appeared after quiescence.

Choose...

A 16-year-old girl starts drinking heavily and binges. She is also found to be skipping school. These set of symptoms begin at the same time as her parents started having frequent conflicts threatening marital separation.

Choose...

A 22-year-old woman describes an irresistible urge and tension that can only be relieved by pulling her hair. Sometimes she experiences a pleasurable sensation as the hair is pulled.

Choose...

Your answer is incorrect.

Explanation:

1) Trauma-focused cognitive-behavioral methods, such as stimulus confrontation and cognitive restructuring, have proved very helpful and were often the treatments of choice for individuals with PTSD. [Seidler GH, Wagner FE. Comparing the efficacy of EMDR and trauma-focused cognitive-behavioral therapy in the treatment of PTSD: a meta-analytic study. Psychol Med. 2006 Nov;36(11):1515-22.]

2) The DBT model assumes that individuals with Borderline personality disorder lack essential interpersonal, self-regulation, and distress tolerance skills. Those personal and environmental factors may frequently block and inhibit the use of behavioural skills or reinforce maladaptive actions. [Feigenbaum J. Dialectical behaviour therapy. Psychiatry. Volume 7, Issue 3, Pages 112-116 (March 2008)].

3) Family therapy concentrates on altering the interactions between or among family members. It seeks to improve the functioning of the family as a unit, or its subsystems, and the functioning of individual members of the family. [Kaplan & Sadock's Synopsis of Psychiatry: Behavioral Sciences/Clinical Psychiatry, 10th Edition Pg: 940].

4) Habit reversal therapy is a cognitive behavioral therapy technique for trichotillomania and Tourette's syndrome that includes four steps - self-monitoring, awareness training, stimulus control and stimulus-response intervention or competing response intervention. [Bloch MH. Systematic Review: Pharmacological and Behavioral Treatment for Trichotillomania. BIOL PSYCHIATRY 2007]

The correct answer is: A young woman develops anxiety, recurrent flashbacks and incapacitating fear of driving after having a dreadful accident. → Trauma focussed CBT, A 25-year-old woman presents with high-frequency self-harm behaviour. She says she cannot trust people anymore as none of her previous therapists offered her anything useful. They transferred her care back to her community psychiatrist whenever her self-harming behaviour appeared after quiescence. → Dialectic behavioural therapy, A 16-year-old girl starts drinking heavily and binges. She is also found to be skipping school. These set of symptoms begin at the same time as her parents started having frequent conflicts threatening marital separation. → Family therapy, A 22-year-old woman describes an irresistible urge and tension that can only be relieved by pulling her hair. Sometimes she experiences a pleasurable sensation as the hair is pulled. → Habit reversal training



Question 23

Not answered

Marked out of 4.00

Psychotherapy

Choose one appropriate therapy for each of the following:

Graded exposure

Choose...

Traps, dilemmas, snags and reformulation letters

Choose...

Rolling with resistance

Choose...

Zen Buddhist principles

Choose...

Your answer is incorrect.

Explanation:

1) Behavioural therapy for phobias is based on the principle that avoidance of fearful object feeds to the fear by reinforcing phobia each time the object is avoided. In BT, therapists work with the client to help the client face their fears. The difference with graded exposure is it is carried out much more slowly, taking little steps forward each time.

2) Traps, dilemmas and snags are terms associated with Anthony Ryle's CAT. Traps are things we cannot escape. Certain kinds of thinking and acting end in a vicious circle, when, however hard we try, things seem to get worse instead of better. Dilemmas are false choices and narrow options. Snags (Subtle Negative Aspects of Goals) are what is happening when we say "I want to have a better life, or I want to change my behaviour but.....".

3) Resistance can take several forms. It includes negating, excusing, blaming, minimizing, arguing, ignoring challenging and interrupting. In motivational interviewing one does not directly oppose resistance but, rather, rolls or flows with it.

4) Dialectical Behavioral Therapy (DBT) for borderline personality and Mindfulness Based Cognitive Behavioral Therapy (CBT) for depression are psychotherapy programs that have become popular interventions utilizing Zen mindfulness philosophy.

The correct answer is: Graded exposure → Behavioural therapy, Traps, dilemmas, snags and reformulation letters → Cognitive analytical therapy, Rolling with resistance → Motivational interviewing, Zen Buddhist principles → Mindfulness based CBT

Question 24

Not answered

Marked out of 4.00

Cognitive distortions

Identify the type of cognitive distortions for each of the following examples

Tom is a perfectionist. When he doesn't get something right, he feels as if he is a 'total failure'.

Choose...



Sam feels overwhelmed by the amount of work he has to do and thinks there is no point in doing any of it.

Choose...



Sally was shopping when she had some palpitations. She felt very anxious and expected to collapse with a heart attack.

Choose...



George saw her girlfriend talking to another man on the street and concluded that she was cheating on him

Choose...



Your answer is incorrect.

Explanation:

- 1) Overgeneralizing is generalizing a small element to the whole: e.g. 'If something doesn't work out once, assume it never will'.
- 2) People with all or nothing thinking view things in extreme categories. For example, an individual may see the tasks allocated to him/her in a job as either all feasible or altogether impossible, with no 'grey-area' between the two.
- 3) Catastrophisation is predicting the worst-case scenario to develop even from seemingly trivial occurrences.
- 4) Arbitrary inference refers to drawing conclusions without sufficient evidence

The correct answer is: Tom is a perfectionist. When he doesn't get something right, he feels as if he is a 'total failure'. → Overgeneralizing, Sam feels overwhelmed by the amount of work he has to do and thinks there is no point in doing any of it. → All or nothing thinking, Sally was shopping when she had some palpitations. She felt very anxious and expected to collapse with a heart attack. → Catastrophisation, George saw her girlfriend talking to another man on the street and concluded that she was cheating on him → Arbitrary inference

Question 25

Not answered

Marked out of 4.00

Psychotherapy variations

Using the descriptions given below, identify the type of psychotherapy

Focus on a set of unspoken rules that organize ways in which family members relate to one another. Such a structure will have the hierarchy of power (parents in charge of children) and firm boundaries.

Choose...

This approach focus on how the family members communicate with each other and concentrate on improving communication skills

Choose...

Focuses mainly on the present functioning of the family rather than past experiences and uses paradoxical injunctions

Choose...

This approach emphasises more active interventions and specific solutions for clearly defined family problem

Choose...

Your answer is incorrect.

Explanation:

- 1) Structural model (Minuchin) views family as a structure built of interpersonal relationships. Such a structure will have the hierarchy of power (parents in charge of children), firm boundaries and set of unspoken rules. Families function particularly well when these family structures are intact. Disrupted structure leads to disrupted communication.
- 2) In Eclectic family therapy, the focus is on the present situation of the family, how the members communicate with each other and concentrate on improving communication skills. Communication theory is also very helpful in family therapy, and much of the efforts to assist the family can be directed towards helping them develop better communication skills. During the family therapy session, any topic or interview content could be discussed; what is crucial is the feeling tones, relationship, and communication skills exercised.
- 3) Systemic family therapy (Milan) It gives great emphasis on circular and reflexive questioning. In a circular fashion, each family member is asked to comment and reflect on each other's response. Focuses mainly on the present functioning of the family rather than past experiences and uses paradoxical injunctions.

4) Strategic therapy by Haley. It emphasises more active interventions and specific solutions for a clearly defined family problem. Believes that symptoms are maintained by behaviors that seek to suppress them at first

The correct answer is: Focus on a set of unspoken rules that organize ways in which family members relate to one another. Such a structure will have the hierarchy of power (parents in charge of children) and firm boundaries. → Structural family therapy, This approach focus on how the family members communicate with each other and concentrate on improving communication skills → Eclectic family therapy, Focuses mainly on the present functioning of the family rather than past experiences and uses paradoxical injunctions → Systemic family therapy, This approach emphasises more active interventions and specific solutions for clearly defined family problem → Strategic family therapy



Question 26

Not answered

Marked out of 4.00

Psychotherapy

Using the descriptions given below, identify the type of psychotherapy

This therapy is based on the principle that current problems in the family originate from past experience and becomes long standing

Choose...

This therapy focuses on reciprocal role repertoire, recognition of maladaptive behaviour and reformulation of this behaviour

Choose...

This therapy focuses on Emotion modulation skills and distress tolerance skills in personality disorder patients

Choose...

The techniques used are based on learning theory that are used to extinguish maladaptive behaviours and substitute more adaptive ones

Choose...

Your answer is incorrect.

Explanation:

- 1) Psychodynamic therapy is based on the principle that current problems in the family originate from past experiences of its family members.
- 2) Cognitive analytical therapy includes recognition of maladaptive behaviour and reformulation of this behaviour and revision.
- 3) DBT techniques include mindfulness skills, interpersonal effectiveness training, Emotion modulation skills and distress tolerance skills.
- 4) Behavioural therapy is based on learning theory that are used to extinguish maladaptive behaviours and substitute more adaptive ones

The correct answer is: This therapy is based on the principle that current problems in the family originate from past experience and becomes long standing → Psychodynamic psychotherapy, This therapy focuses on reciprocal role repertoire, recognition of maladaptive behaviour and reformulation of this behaviour → Cognitive analytical therapy, This therapy focuses on Emotion modulation skills and distress tolerance skills in personality disorder patients → Dialectical behavioural therapy, The techniques used are based on learning theory that are used to extinguish maladaptive behaviours and substitute more adaptive ones → Behavioural therapy



Question 27

Not answered

Marked out of 6.00

Therapeutic factors in Group Therapy

With the definitions given below, identify the therapeutic factors in Group Psychotherapy from the list of options given:

The expression of ideas, thoughts, and suppressed material that is accompanied by an emotional response. It produces a state of relief in the patient.

Choose...



Putting another person's need before one's own and learning that there is value in giving to others.

Choose...



The sense that the group is working together toward a common goal

Choose...



Confirmation of reality by comparing one's conceptualisations with those of other group members

Choose...



The awareness in a patient that he or she is not alone in having problems; others share similar complaints or difficulties in life..

Choose...



It is a process by which repressed material, particularly a painful experience or conflict, is brought back to consciousness.

Choose...



Your answer is incorrect.

Explanation:

- 1) Catharsis is the expression of ideas, thoughts, and suppressed material that is accompanied by an emotional response. It produces a state of relief in the patient.
- 2) Altruism is putting another person's need before one's own and learning that there is value in giving to others.
- 3) Group Cohesion is the sense that the group is working together toward a common goal.
- 4) Consensual validation is Confirmation of reality by comparing one's conceptualisations with those of other group members.

5) Universalisation is the awareness of the patient that he or she is not alone in having problems; others share similar complaints or difficulties in learning; the patient is not unique. 6) Abreaction is the process by which repressed material, particularly a painful experience or conflict, is brought back to consciousness

The correct answer is: The expression of ideas, thoughts, and suppressed material that is accompanied by an emotional response. It produces a state of relief in the patient. → Catharsis, Putting another person's need before one's own and learning that there is value in giving to others. → Altruism, The sense that the group is working together toward a common goal → Cohesion, Confirmation of reality by comparing one's conceptualisations with those of other group members → Consensual validation, The awareness in a patient that he or she is not alone in having problems; others share similar complaints or difficulties in life.. → Universalisation, It is a process by which repressed material, particularly a painful experience or conflict, is brought back to consciousness. → Abreaction



Question 28

Not answered

Marked out of 5.00

Cognitive profile of psychiatric disorders (2)

Identify the type of cognitive distortion for each of the definitions given below;

Always predicting the worst-case scenario to happen	Choose... ▾
Predicting without sufficient evidence	Choose... ▾
Picks failures and errors and ignores positive features	Choose... ▾
Exaggerates negative events	Choose... ▾
Black or white thinking	Choose... ▾

Your answer is incorrect.

Explanation:

- 1) Always predicting the worst-case scenario to happen suggests catastrophization. e.g. A woman with panic attacks who can sense mild palpitation after some housework thinks she is going to have a massive heart attack and immediate death.
- 2) Predicting without sufficient evidence is arbitrary inference.
- 3) In selective abstraction, the patient selectively picks failures and errors and ignores positive features. - e.g. A dysthymic student can talk only about his minor failures and withdrawals from college competitions, and chooses to ignore the A grades and accolades that he has received so far.
- 4) In magnification, the person exaggerates negative events, and you tend to filter out or disqualify more positive information. In minimization, the person minimizes positive outcomes.
- 5) Black or white thinking is also called as dichotomous thinking where the subject can see problems and challenges to have only two outcomes - missing all the alternative options in between the highly favourable and highly unfavourable choices.

The correct answer is: Always predicting the worst-case scenario to happen → Catastrophisation, Predicting without sufficient evidence → Arbitrary inference, Picks failures and errors and ignores positive features → Selective abstraction, Exaggerates negative events → Magnification, Black or white thinking → Dichotomization



Question 29

Not answered

Marked out of 5.00

Terminologies in psychotherapy (2)

Identify the type of therapy using the terms given below;

Setting of tasks for the family before sessions	Choose... ▾
Circular questioning and problem-solving	Choose... ▾
Reframing and unbalancing	Choose... ▾
Roll with resistance and support self-efficacy	Choose... ▾
Thought monitoring and behavioural experiments	Choose... ▾

Your answer is incorrect.

Explanation:

- 1) Strategic therapy involves using active interventions to fit the particular problem and involves setting of tasks before sessions for the family.
- 2) The techniques involved in systemic family therapy include circular questioning, problem solving, hypothesizing, neutrality, paradox and counter-paradox.
- 3) The structural therapy involves hierarchies, redesigning, reframing the family systems so that it approximates a normative model of care but open boundaries between generational subsystems.
- 4) There are five general principles of motivational interviewing: express empathy, develop discrepancy, avoid argumentation, roll with resistance, and support self-efficacy. In CBT, a common strategy used is thought monitoring (use of self-monitoring and self-help assignments between session).
- 5) A behavioural experiment in CBT often involves setting up an experiment or task to confirm or test the conclusions of the examination of the thoughts.

The correct answer is: Setting of tasks for the family before sessions → Strategic family therapy, Circular questioning and problem-solving → Systemic family therapy, Reframing and unbalancing → Structural family therapy, Roll with resistance and support self-efficacy → Motivational interviewing, Thought monitoring and behavioural experiments → Cognitive behavioural therapy



Question 30

Not answered

Marked out of 4.00

History of Psychotherapy

For each of the description/intervention below chose one historical figure associated with the concept of the given list

Coined the term Therapeutic Community

Choose... ▾

Described permissiveness and democratisation in group settings.

Choose... ▾

Described curative factors that help in the process of recovery in group therapies

Choose... ▾

Credited with the establishment of therapeutic community model

Choose... ▾

Your answer is incorrect.

Explanation:

Tom Main coined the term therapeutic community.

Maxwell Jones established the therapeutic community, and

Rapoport explained the key features of Therapeutic communities that include communalism, democracy, permissiveness and reality confrontation.

Yalom cited 11 'curative' factors responsible for a change in groups. The curative factors include the instillation of hope, universality, imparting information (feedback), altruism, corrective recapitulation, socialisation techniques, imitative behaviour, interpersonal learning, group cohesiveness, catharsis and existential factors. Of these, cohesiveness, and learning from feedback are valued positively though other factors may also be important.

The correct answer is: Coined the term Therapeutic Community → Main, Described permissiveness and democratisation in group settings. → Rapoport, Described curative factors that help in the process of recovery in group therapies → Yalom, Credited with the establishment of therapeutic community model → Jones

Question 31

Not answered

Marked out of 5.00

History of Psychiatry

Identify the pioneers/proponents for each of the following concepts/treatments

CBT for eating disorders

Choose...



Play Therapy

Choose...



Analytical psychology

Choose...



Cognitive therapy for depression

Choose...



CBT for delusions and hallucinations

Choose...



Your answer is incorrect.

Explanation:

- 1) A standard form of CBT that is used to treat eating disorders is called CBT-Enhanced (CBT-E). It was developed by Christopher G. Fairburn, originally intended for bulimia nervosa specifically; it was eventually extended to all eating disorders.
- 2) Melanie Klein initially introduced the technique of using play as a means of analyzing children under the age of six. She believed that observing a child's play is a tool similar to the free association in adults and provides access to the child's unconscious.
- 3) Carl Gustav Jung is the founder of the school of Analytical Psychology.
- 4) Cognitive therapy (CT) is a type of psychotherapy developed by American psychiatrist Aaron Beck.
- 5) Following the pioneering work of Kingdon & Turkington in 1994, there has been a growing interest in the application of cognitive techniques in psychosis for the treatment of delusions and hallucinations.

The correct answer is: CBT for eating disorders → Christopher Fairburn, Play Therapy → Melanie Klein, Analytical psychology → Jung, Cognitive therapy for depression → Beck, CBT for delusions and hallucinations → Turkington

Question 32

Not answered

Marked out of 4.00

Proponents & Therapies II

Match each of the proponents below with their therapy/school listed below

Jacob Moreno	Choose... 
Masters and Johnson	Choose... 
Joseph Wolpe	Choose... 
Carl Rogers	Choose... 

Your answer is incorrect.

Explanation:

Carl Rogers is best known for his contributions to a humanistic form of therapy initially called as non-directive therapy. He felt that the client should direct the progress of the therapy. The therapist should not lead the client, but rather be there for the client. It is also known as client centred or Rogerian therapy.

Jacob Levy Moreno, the originator of psychodrama, was a Viennese doctor who practiced with storytelling in children's groups, and it is followed by having the children act out the stories. He later used this method with adults that formed the basis for psychodrama.

Moreno entered psychodrama through the practice of sociometry. It is an observational charting of how people interact in groups.

The correct answer is: Jacob Moreno → Psychodrama, Masters and Johnson → Dual sex therapy, Joseph Wolpe → Reciprocal inhibition, Carl Rogers → Client-centered psychotherapy

Question 33

Not answered

Marked out of 5.00

Proponents & Therapies III

Match each of the proponents below with their therapy/school listed below

Jacobson	Choose... 
Frederick Perls	Choose... 
Otto Rank	Choose... 
Arthur Janov	Choose... 
Albert Ellis	Choose... 

Your answer is incorrect.

Explanation:

Progressive muscular relaxation was advocated by Hacobson.

Gestalt therapy was founded by Frederick (Fritz) and Laura Perls in the 1940s. It teaches therapists and patients the phenomenological method of awareness. The goal is for clients to become aware of what they are doing, how they are doing it, and how the clients can change themselves, and at the same time, to learn to accept and value themselves.

Primal therapy is a form of regressive psychotherapy based on trauma. Arthur Janov, who reinstated Freud's views that neurosis is caused by the repressed pain of childhood trauma, originally described this form of treatment.

Ellis described RET - Rational Emotive Therapy.

Will Therapy of Otto Rank stressed choice, responsibility, and action. He argued that neurotics were failed artists. They could regain their will through analysis, in a process of self-creation or rebirth.

The correct answer is: Jacobson → Progressive muscular relaxation, Frederich Perls → Gestalt therapy, Otto Rank → Will Therapy, Arthur Janov → Primal therapy, Albert Ellis → Rational-emotive therapy



Question **34**

Not answered

Marked out of 4.00

Proponents & Therapies IV

Match each of the proponents below with their therapy/school listed below

Ayllon and Azrin	Choose... ▾
Klerman and Weissmann	Choose... ▾
Marsha Linehan	Choose... ▾
Victor E Frankl	Choose... ▾

Your answer is incorrect.

Explanation:

Interpersonal Psychotherapy of Depression (IPT) was developed in the New Haven-Boston Collaborative Depression Research Project by Gerald Klerman and Myrna Weissman for the treatment of ambulatory depressed, nonpsychotic, nonbipolar patients.

The token economy was pioneered by Ayllon, Azrin, Paul & Krasner (Camarillo State Hospital, Calif.). The full range of self-care, social, and work behaviors were thought to be modifiable through systematic and preplanned use of antecedents and consequences of these actions. The "psychopathological" behavior of the mentally ill was conceptualized as being subject to the same "laws of learning" that influenced normal behavior. The tokens could be conveniently dispensed to patients contingent on their exhibiting improvements in their behavior. (Am J Psychiatry 157:9, September 2000).

Logotherapy according to Frankl (1978) is a 'therapy through meaning'. It significantly contradicts more traditional conceptualisations of psychotherapy, where the individual would be provided with meaning through therapy. Logotherapy is based on the belief that the meaning one attaches to circumstances, rather than the circumstances themselves, determines their impact.

The correct answer is: Ayllon and Azrin → Token economy, Klerman and Weissmann → Interpersonal therapy, Marsha Linehan → Dialectical behavioural therapy, Victor E Frankl → Existential Logotherapy

Question 35

Not answered

Marked out of 8.00

Group therapy (1)

Choose two Options for the questions below from the list below:

1. Bion's factors which affect group therapy.

    

2. Interpersonal "Curative factors" in a group.

    

3. Factors seen facilitating communication in a psychodynamic group

    

4. Basic techniques used in psychodrama

    **Explanation:**

1) There are three factors that affect group therapy initially described by Bion. They are pairing, dependency, and fight or flight.

2) Yalom cited 11 'curative' factors responsible for a therapeutic change in groups. The positive curative factors include cohesiveness and universality. Other factors include the instillation of hope, imparting information (feedback), altruism, corrective recapitulation, socialisation techniques, imitative behaviour, interpersonal learning, catharsis and existential factors.

3) The factors that facilitate communication in a psychodynamic group include free-floating discussion and mirroring. Exchange, Resonance and Translation are also identified as factors affecting communication in a group setting.

4) Role playing and mirroring are techniques used in psychodrama. Role reversal/playing refers to the exchange of the patient's role for the role of a significant person. Mirror technique refers to imitating the patient and speaking in the proxy. A therapeutic dramatization of emotional problems is the main principle employed.

Correct Answer is : 1. Dependency, Fight or flight 2. Cohesiveness, Universality 3. Free floating discussion, Mirroring 4. Mirroring , Role playing



Question **36**

Not answered

Marked out of 8.00

Psychotherapeutic concepts (1)

Identify TWO concepts associated with each of the following psychotherapies

1. Interpersonal psychotherapy

 ✖ ✖

2. Cognitive analytical therapy

 ✖ ✖

3. Dialectical behavioural therapy

 ✖ ✖

4. Eye movement desensitisation reprocessing

 ✖ ✖

Explanation:

- 1) In interpersonal therapy, 4 problem areas are explored - interpersonal role disputes (conflicts with family member, colleague, or friend), role transitions (change from one role to other - includes graduation, beginning or end of career, migration, medical ill health etc.), grief (loss of a loved object/relation) and interpersonal deficits (social impoverishment, inadequate interpersonal relationships).
- 2) The two theoretical bases in cognitive analytical therapy are - (PSM) procedural sequence model and role repertoires. Faulty revisions lead to maladaptive sequences that contribute to the pathology. Ryle described three main kinds of faulty procedures: Traps, Snags and Dilemmas.
- 3) The important core techniques used in Dialectical Behaviour Therapy includes distress tolerance, Interpersonal effectiveness training, Core mindfulness training and Emotion regulation skills.
- 4) Body scan is one of the eight stages of eye movement desensitisation & reprocessing. The Subjective Units of Distress Scale (SUDS) is a scale of 0 to 10. It is helpful for measuring the subjective intensity of disturbance or distress currently experienced by an individual. It has been used in EMDR and other trauma-focussed therapies for anxiety disorders.
- Correct Answer is : 1. Role transitions, Role disputes 2. Role repertoires, Maladaptive procedural sequences 3. Interpersonal effectiveness, Core mindfulness 4. Subjective units of distress scale, Body scan



Question 37

Not answered

Marked out of 8.00

Psychotherapeutic concepts (2)

Identify TWO concepts associated with each of the following psychotherapies

1. Behavioural couple therapy (Choose TWO)

 ✖ ✖

2. Family therapy (Choose TWO)

 ✖ ✖

3. Systematic desensitisation (Choose TWO)

 ✖ ✖

4. Rational emotional therapy (Choose TWO)

 ✖ ✖

Explanation:

1) Behavioural Couple Therapy is a specific intervention for alcoholism. It works directly to increase relationship factors conducive to abstinence. The therapist arranges a daily "sobriety contract". According to the contract, the patient states his or her intent not to drink or use drugs that day (one day at a time). The spouse of the patient expresses support for the patient's efforts to stay abstinent. Relapse prevention is the final activity of BCT.

2) Paradoxical Injunction and emotional triangle are terms associated with family therapy. An emotional triangle is a three-party system where the closeness of two members (in either positive or negative sense) tends to exclude the third. This hot triangle leads to symptom formation. The term 'paradoxical injunction' was formerly associated with Milan school of family therapy.

3) Systematic Desensitization (Wolpe) is based on the behavioural principle of counterconditioning (i.e., gradual approach of feared situation in a psychophysiological state that inhibits anxiety leads to the reduction of anxiety response). Systematic desensitization consists of three steps a) relaxation training b) hierarchy construction, and c) desensitization of the stimulus.

4) Rational emotional therapy was developed by Albert Ellis. It is a reality-based, short-term therapy, aimed at 'cognitive re-structuring' with many similarities to the later developed CBT. The therapist aims to encourage the patient to examine their basic beliefs; encourage the replacement of 'irrational thinking' with rational thinking and to examine the resultant change in their emotional response.

Correct Answer is : 1. Sobriety contract, Relapse prevention 2. Paradoxical Injunction, Emotional triangle 3. Hierarchy construction, Counter conditioning 4. Cognitive restructuring, Irrational thinking



Question 38

Not answered

Marked out of 8.00

Psychotherapeutic concepts (3)

Identify TWO concepts associated with each of the following psychotherapies

1. Psychoanalytical Psychotherapy (Choose TWO)

 ✖ ✖

2. Group therapy (Choose TWO)

 ✖ ✖

3. Cognitive behavioural therapy (Choose TWO)

 ✖ ✖

4. Therapeutic community (Choose TWO)

 ✖ ✖

Explanation:

1) Resistance refers to any hindrance noted in the course of therapeutic progress during analytical psychotherapy. Interpretation refers to the expression of therapist's understanding of the meaning of feelings, attitudes, defense mechanisms and behaviours currently exhibited during therapy. Interpretation is usually based on the specific psychoanalytical theory practiced by the therapist.

2) Group cohesion refers to the sense that the group is working together towards a common goal. Altruism, which refers to helping and supporting others, is also one of Yalom's 11 curative factors.

3) CBT identifies three levels of cognitive dysfunctions: negative automatic thoughts, conditional assumptions and fundamental or core beliefs (or schemas). Schemas are one's appraisal of oneself, and they usually begin with - 'I am' e.g. 'I am useless', 'I am a loser', etc. Automatic thoughts are cognitions that automatically arise in certain situations or as a reflex response to certain behaviours. Negative automatic thoughts predominate depressive thinking.

4) Permissiveness and reality confrontation are 2 of the four components of the therapeutic community (Rappaport / Henderson Hospital Model). Permissiveness is tolerating each other, and realizing unpredictable behaviour can happen within the community. Reality confrontation is self-deception or distortions from reality are dealt with honestly and openly by all members without formalities. The other two principles are communalism and democratization.

Correct Answer is : 1. Resistance, Interpretation 2. Altruism, Cohesion 3. Schemas, Automatic thoughts 4. Reality confrontation, Permissiveness



Question 39

Not answered

Marked out of 7.00

Safety behaviours

Identify the safety behaviours commonly used in various disorders

1. Obsessive compulsive disorder ✖
2. Panic disorder ✖
3. Post-traumatic stress disorder ✖
4. Social phobia (Choose TWO) ✖ ✖
5. Hypochondriasis (Choose TWO) ✖ ✖

Explanation:

Active suppression is safety behaviour tried in PTSD as well as OCD - but this increases the probability that the intrusive thoughts and flashbacks will occur.

Hyperventilation is a safety behaviour often seen in response to feelings of choking or respiratory arrest during a panic attack. But this, unfortunately, produces a flurry of symptoms due to physiological acid-base changes.

Rigorous self-monitoring, avoiding eye contact and monitoring of speech may be safety behaviour in social phobia- this can feed to the core symptoms.

The safety behaviours involved in hypochondriasis are reassurance seeking and checking for symptoms.

Correct Answer is : 1. Active suppression 2. Hyperventilation 3. Active suppression 4. Rigorous self-monitoring, Avoiding eye contact 5. Reassurance seeking, Checking for symptoms

Question 40

Not answered

Marked out of 6.00

Psychotherapy terms

Identify 2 concepts associated with each of the following psychotherapies?

1. Psychoanalytical Psychotherapy

 ✖ ✖

2. Solution focussed therapy

 ✖ ✖

3. Cognitive analytical therapy

 ✖ ✖

Explanation:

1) Resistance refers to any hindrance noted in the course of therapeutic progress during analytical psychotherapy. Interpretation refers to the expression of therapist's understanding of the meaning of feelings, attitudes, defense mechanisms and behaviours currently exhibited during therapy. Interpretation is usually based on psychoanalytical theory practiced by the therapist.

2) Solution focussed therapy aims to empower patients to recognise and make use of their strengths, It is a brief intervention that may be delivered in a single session. The techniques include problem free talk, preferred future, exception finding and scales.

3) The three Rs of cognitive analytical therapy are recognition of maladaptive behaviour and beliefs, reformulation of these ideas and revision.

Correct Answer is : 1. Resistance, Interpretation 2. Problem free talk, Preferred future 3. Recognition of maladaptive behaviour, Reformulation



Question 41

Not answered

Marked out of 8.00

Group Psychotherapy

For each of the following descriptions given below, select the two therapeutic concepts from the list below;

1. These are characteristic features of analytical group psychotherapy



2. These are recognised unconscious strategies by which group members avoid working in the group and are called basic assumptions



3. These are reasons for the effectiveness of group psychotherapy and are cited as reasons why interpersonal groups have a therapeutic effect



4. These are basic techniques used in psychodrama



Explanation: 1) The characteristic features in analytical group psychotherapy are free-floating discussion and interpretation. Interpretation refers to the expression of therapist's understanding of the meaning of feelings, attitudes, defense mechanisms and behaviours currently exhibited during therapy. Interpretation is usually based on psychoanalytical theory practiced by the therapist. 2) There are three factors affecting group therapy described by Bion (Basic assumptions). They are pairing, dependency, fight or flight. 3) Yalom cited 11 'curative' factors responsible for a change in groups. The curative factors include cohesiveness and universality. The other factors include the instillation of hope, imparting information (feedback), altruism, corrective recapitulation, socialisation techniques, imitative behaviour, interpersonal learning, catharsis and existential factors. 4) Role playing and mirroring are techniques used in psychodrama. Role reversal/playing refers to the exchange of the patient's role for the role of a significant person. Mirror technique refers to imitating the patient and speaking in the proxy. Jacob Moreno founded psychodrama. A therapeutic dramatization of emotional problems is the main principle employed.

Correct Answer is : 1. Free floating discussion, Interpretation 2. Dependence, Pairing 3. Cohesiveness, Universality 4. Role playing, Mirroring



© 2023 SPMM Course Limited, UK.

[Refund Policy](#) | [Terms & Conditions](#)

