

Started on	Sunday, 12 February 2023, 8:29 PM
State	Finished
Completed on	Sunday, 12 February 2023, 8:29 PM
Time taken	12 secs
Marks	0/40
Grade	0 out of 100



Question 1

Not answered

Marked out of 1

Pritchard criteria include all of the following except

Select one:

- ☐ Ability to instruct legal counsel
- ☐ Ability to follow proceedings in the court
- ☐ Ability to understand charges
- ☐ Ability to understand the difference between diminished responsibility vs automatisms
- ☐ Ability to challenge any of the jurors to whom they might object

Your answer is incorrect.

Explanation:

R v Pritchard criteria are often cited when discussing the 'test' for fitness to plead. An individual is unfit to plead if they are unable to:

- Understand the course of proceedings on trial
- To comprehend the details of evidence
- To know they might challenge any of the jurors to whom they might object
- To instruct legal counsel

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 774.

The correct answer is: Ability to understand the difference between diminished responsibility vs automatisms

Question 2

Not answered

Marked out of 1

Which of the following defendants is most likely to be fit to plead in a court trial?

Select one:

- ☐ A defendant who cannot understand the nature of the charges
- ☐ A defendant with amnesia for the offence
- ☐ A defendant who does not understand trial procedures
- ☐ A defendant with lack of ability to challenge the jurors
- ☐ A defendant who is catatonic, mute and not able to communicate

Your answer is incorrect.

Explanation:

R v Pritchard criteria are often cited when discussing the 'test' for fitness to plead. An individual is unfit to plead if they are unable to:

- Understand the course of proceedings on trial
- To comprehend the details of evidence
- To know they might challenge any of the jurors to whom they might object
- To instruct legal counsel

These factors are relevant at the time of trial only (not for arrest or laying charges). Amnesia for the offence does not interfere with fitness to plead, but short-term memory failure due to organic impairments may be such as to make following proceedings in court impossible.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 774-775.

The correct answer is: A defendant with amnesia for the offence

Question **3**

Not answered

Marked out of 1

What percentage of people in police custody has a mental illness at the time of arrest?

Select one:

- ☐ 25-50%
- ☐ 2-5%
- ☐ 5-10%
- ☐ 20-25%
- ☐ 15-20%

Your answer is incorrect.

2-5% of people held in custody by the police suffer from mental disorder. About 1-2% suffer from severe mental illness (Oxford Handbook of Psychiatry 2nd ed - Pg650)

The correct answer is: 2-5%



Question 4

Not answered

Marked out of 1

The prevalence of psychotic disorders in prisons compared to the general population is

Select one:

- ☐ 10 times
- ☐ 50 times
- ☐ 5 times
- ☐ 20 times
- ☐ 40 times

Your answer is incorrect.

A rate higher by ten times (in the UK, Ref: Brugha et al., 2005) to twenty times (elsewhere), giving a prevalence rate - 2-15% has been noted in various studies. Fazel and Danesh conducted a meta-analysis of 62 surveys from 12 countries including 22 790 prisoners wherein 3-7% of men had psychotic illnesses, 10% major depression, and 65% a personality disorder, including 47% with antisocial personality disorder. Prisoners were about ten times more likely to have antisocial personality disorder than the general population. Fazel, S and Danesh, J (2002) Serious mental disorders in 23 000 prisoners: a systematic review of 62 surveys. The Lancet 359, 545-550.

The correct answer is: 10 times

Question 5

Not answered

Marked out of 1

Five patients are being discharged from a psychiatric inpatient unit. Considering their diagnostic profile, choose the one who has the highest risk of being violent in next six months.

Select one:

- ☐ 35 year old man with depression
- ☐ 33 year old man with bipolar disorder
- ☐ 34 year old man with schizophrenia
- ☐ 30 year old man with mixed personality disorder
- ☐ 32 year old man with a substance use disorder

Your answer is incorrect.

Explanation:

Alcohol and drug-related problems are more strongly linked to offending and violence than any other mental disorders. A number of aspects of alcohol and substance misuse may be relevant - direct effects of intoxication or withdrawal; funding the habit; personal and social consequences of dependence; the neuropsychiatric sequelae of prolonged misuse; and the social context (peer group, socio-economic deprivation, childhood mistreatment). Also, personal characteristics (impulsivity and sensation seeking), which may lead to substance misuse, may also be associated with offending.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 747.

The correct answer is: 32 year old man with a substance use disorder

Question 6

Not answered

Marked out of 1

Most offenders charged with exhibitionism belong to the age group

Select one:

- ☐ 55-65 years
- ☐ 45-55 years
- ☐ 15-25 years
- ☐ 35-45 years
- ☐ 25-35 years

Your answer is incorrect.

Most exhibitionists are male and their victims are usually female adults or children. Most males arrested for exhibitionism are in the age group 16 to 25. In general, exhibitionists have the highest sex offence recidivism rates compared to other offender groups though recidivism reduces significantly when charged/sentenced once.

The correct answer is: 15-25 years



Question 7

Not answered

Marked out of 1

The male:female conviction rate for any offence in the UK is approximately

Select one:

- ☐ 5 to 1
- ☐ 2 to 1
- ☐ 20 to 1
- ☐ 1 to 1
- ☐ 10 to 1

Your answer is incorrect.

Explanation:

Females comprise under 20% of offenders. Peak rates occur at age 18-20 years for males, and 2-3 years earlier for females.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 728.

The correct answer is: 5 to 1



Question 8

Not answered

Marked out of 1

When women with psychosis commit homicide, their victims are most often aged

Select one:

- ☐ >75 years
- ☐ 36-59 years
- ☐ 16-35 years
- ☐ <16 years
- ☐ 60-75 years

Your answer is incorrect.

Explanation:

When women with psychosis kill another person, the victim is most commonly a family member <18 years old. Most child victims of homicide are killed by a parent or stepparent and in the majority of cases, mental illness is not a feature of filicide. Homicide of a child is more likely to be committed by a male. Women perpetrators are more likely to have a history of a mental disorder and symptoms at the time of the offence (most often an affective disorder, followed by a primary psychotic disorder).

Ref: Ferranti J et al. Characteristics of female homicide offenders found not guilty by reason of insanity. Journal of the American Academy of Psychiatry and the Law. 2013 Dec 1.

Flynn SM et al. Filicide: mental illness in those who kill their children. PloS one. 2013 Apr 4;8(4):e58981.

The correct answer is: <16 years

Question 9

Not answered

Marked out of 1

The presence of mental illness increases the risk of being victimised with violent assault by

Select one:

- ☐ 10 times
- ☐ 12 times
- ☐ 20 times
- ☐ 2 times
- ☐ 5 times

Your answer is incorrect.

In National Crime Survey (Khalifeh et al., 2015), 40% of patients with a severe mental illness were noted to be victims of crime in the past year, compared to 14% of general population (odds ratio = 2.8); the risk ratio was much higher when violent assaults were considered (19% patients v. 3% of controls (OR = 5.3)).

The correct answer is: 5 times



Question 10

Not answered

Marked out of 1

To be defined as a murder-suicide event, the suicide of a homicide perpetrator must occur within a period of

Select one:

- ☐ 1 year after the murder
- ☐ 1 day after the murder
- ☐ 1 hour after the murder
- ☐ 1 week after the murder
- ☐ 1 month after the murder

Your answer is incorrect.

Murder-suicide: It is defined as a murder that is followed by the suicide of the perpetrator within one week of the homicide. The incidence is 0.2 –0.3/100,000 per year (Eliason, 2009).

The correct answer is: 1 week after the murder



Question 11

Not answered

Marked out of 1

The proportion of stalking incidents that are reported to the police is

Select one:

- ☐ 1 in 5
- ☐ 1 in 4
- ☐ 1 in 3
- ☐ 2 in 3
- ☐ 1 in 2

Your answer is incorrect.

Of all the different types of interpersonal violence, stalking is the most commonly reported violence in terms of police records. But even for this, in only one in three (31% women last year; 30% men) cases did the police come to know. British Crime Survey 2001

The correct answer is: 1 in 3



Question 12

Not answered

Marked out of 1

The proportion of those discharged from medium secure units that are convicted for a violent offence within five years of discharge is

Select one:

- ☐ 1 in 3
- ☐ 1 in 6
- ☐ 1 in 20
- ☐ 1 in 4
- ☐ 1 in 2

Your answer is incorrect.

Follow-up studies of Medium Secure discharges suggest that nearly 25% will be readmitted within the 12 months after discharge. The reconviction rate for any offence in 2 years is 15%, but for violent offences this is 6%. This rate nearly doubles to 1 in 3 reoffending in 5 years, but only 1 in 6 will be convicted for a violent offence. From high secure units, at least 20% of those discharged will never live in the community in the ten years after discharge: they get re-institutionalised elsewhere. In this time at least 15% would have been convicted for a serious further offence.

The correct answer is: 1 in 6



Question 13

Not answered

Marked out of 1

Aversion therapy, as historically used in the treatment of several types of paraphilias, is what kind of treatment?

Select one:

- ☐ Behavioural
- ☐ Psychodynamic
- ☐ Psychoeducational
- ☐ Skills training
- ☐ Psychoanalytic

Your answer is incorrect.

Explanation:

Initial treatments addressing paraphilic behaviours were generated in the late 1800s. Both Charcot & Magnan and von Schrenck-Notzing outlined methods for resolving 'unusual' sexual interests. However, little was done after these early attempts until a series of single case reports emerged in the 1950s, mostly conducted by staff at the Institute of Psychiatry of the University of London. Psychiatrists at this Institute reported the effectiveness of behavioural conditioning-based treatments (such as electric aversion therapy) for several types of paraphilias. However, it was not long before the limits of these Pavlovian treatment approaches became apparent.

Ref: Marshall WL. A brief history of psychological theory, research, and treatment with adult male sex offenders. Current Psychiatry Reports. 2018 Aug;20(8):1-8.

The correct answer is: Behavioural

Question **14**

Not answered

Marked out of 1

The risk of stalking by a patient or patient's relative over the course of a psychiatrist's career is;

Select one:

- ☐ 20%
- ☐ 10%
- ☐ 30%
- ☐ 40%
- ☐ 50%

Your answer is incorrect.

Data from the UK suggests that 20% psychiatrists report being stalked at some point during one's career.

<http://isp.sagepub.com/content/54/4/350.abstract>

The correct answer is: 20%



Question 15

Not answered

Marked out of 1

What proportion of elderly offenders has a mental illness?

Select one:

- ☐ 20%
- ☐ 40%
- ☐ 30%
- ☐ 10%
- ☐ 50%

Your answer is incorrect.

Older offenders have more chronic health issues and less family and peer support. 20% of them are reported to have a mental illness.

The correct answer is: 20%



Question **16**

Not answered

Marked out of 1

The proportion of chronic domestic violence that gets reported by women in the UK to the police is

Select one:

- ☐ 50%
- ☐ 20%
- ☐ 30%
- ☐ 10%
- ☐ 40%

Your answer is incorrect.

Sadly many incidents of domestic violence are unreported, not recorded or not prosecuted; therefore domestic violence is often described as a 'hidden crime'. The police were told of 12% of incidents overall, and 22% of those against female chronic victims. They offered advice or support in 60% of incidents, which was found helpful in 40% of cases they were told about. British Crime Survey 1996

The correct answer is: 20%



Question 17

Not answered

Marked out of 1

The proportion of all domestic violence victims that are male is

Select one:

- ☐ 40%
- ☐ 1%
- ☐ 10%
- ☐ 60%
- ☐ 5%

Your answer is incorrect.

Data from Home Office statistical bulletins and the British Crime Survey show that men made up about 40% of domestic violence victims each year between 2004-05 and 2008-09, the last year for which figures are available. In 2006-07, men made up 43.4% of all those who had suffered partner abuse in the previous year, which rose to 45.5% in 2007-08 but fell to 37.7% in 2008-09. British Crime Survey 2010

The correct answer is: 40%



Question 18

Not answered

Marked out of 1

Approximately what percentage of people who commit homicides have a mental illness?

Select one:

- ☐ 5%
- ☐ 0.2%
- ☐ 10%
- ☐ 1%
- ☐ 7%

Your answer is incorrect.

Explanation:

Homicide is the killing of a person by another. A minority of offenders are mentally disordered. From 1993-2019 there were 1540 cases of people being killed by people with mental health conditions in the UK. 11% of people convicted of homicide in the UK from 2005-2015 were mental health patients, most diagnosed with schizophrenia. On average there are around 120 victims of homicide by patients or people who were mentally ill at the time of offence in the UK every year. Many of the victims are children, many are elderly, and many are patients themselves. Victims of homicides by mentally disordered offenders are more often acquaintances (especially a member of the perpetrator's family), rather than strangers.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 732-733.

<https://blogs.bmj.com/bmj/2019/06/28/sarah-markham-we-have-to-learn-from-homicides-committed-by-mental-health-patients>

The correct answer is: 10%

Question 19

Not answered

Marked out of 1

The percentage of children with conduct disorder who go on to have antisocial personality disorder is

Select one:

- ☐ 10%
- ☐ 25%
- ☐ 90%
- ☐ 5%
- ☐ 50%

Your answer is incorrect.

Explanation:

Around half of children with conduct disorder will receive a diagnosis of antisocial personality disorder as adults. Substance misuse is also commonly seen in adult life. There is an increased risk of early death, often by violent and sudden means.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 665.

The correct answer is: 50%



Question **20**

Not answered

Marked out of 1

The percentage of new offenders among those charged of indecent exposure is

Select one:

- ☐ 50%
- ☐ 10%
- ☐ 40%
- ☐ 80%
- ☐ 5%

Your answer is incorrect.

With indecent exposures, the offender usually stops reoffending after a conviction; serious progressive crimes are very rare. Nearly 20-30% reoffend and if such reoffence leads to a conviction, the prognosis is very poor in terms of high recidivism.

The correct answer is: 80%



Question **21**

Not answered

Marked out of 1

The percentage of violence in the population that can be ascribed to psychosis is

Select one:

- ☐ 5%
- ☐ 2%
- ☐ 1%
- ☐ 20%
- ☐ 10%

Your answer is incorrect.

Explanation:

Patients with psychosis commit approximately 1 in 20 violent crimes (population attributable risk of 5%). When people with psychosis are violent, the victim is more likely to be known to them (particularly relatives) than when violence is committed by non-psychotic individuals.

Ref: Fazel S, Grann M. The population impact of severe mental illness on violent crime. American journal of psychiatry. 2006 Aug;163(8):1397-403.

Sample D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 746.

The correct answer is: 5%

Question **22**

Not answered

Marked out of 1

A man drinks excessively and then commits a sexual offence. Which of the following is true?

Select one:

- ☐ Voluntary intoxication is a sane automatism
- ☐ Voluntary intoxication is an insane automatism
- ☐ There is a fixed punishment for alcohol-related sexual offences
- ☐ Drunken behaviour cannot be punished legally
- ☐ Voluntary intoxication does not constitute a plea for insanity

Your answer is incorrect.

Explanation:

Voluntary intoxication is never a defence to a crime of basic intent, such as sexual assault, and cannot be used as the basis for an insanity plea.

Ref: <https://www.cps.gov.uk/legal-guidance/intoxication>

The correct answer is: Voluntary intoxication does not constitute a plea for insanity

Question 23

Not answered

Marked out of 1

Which of the following factors contributes most to the risk of suicide in prison?

Select one:

- ☐ Young age at index crime
- ☐ Divorced marital status
- ☐ Absence of previous contact with mental health services
- ☐ Being a member of a minority ethnic group
- ☐ Occupying a single cell

Your answer is incorrect.

Explanation:

Prison suicide and self-harm has been rising in the UK. The most common method of suicide in prison is hanging. The majority of suicides occur within 6 months of imprisonment; life-sentence prisoners are over-represented. **Occupying a single cell/solitary confinement** is a particularly strong risk factor for both self-harm and suicide. Remand prisoners, young offenders, and those with a history of substance misuse and violent offences are also at higher risk of suicide.

Ref: Favril L et al. Risk factors for self-harm in prison: a systematic review and meta-analysis. The Lancet Psychiatry. 2020 Aug 1;7(8):682-91.

Burki T. Crisis in UK prison mental health. The Lancet Psychiatry. 2017 Dec 1;4(12):904.

The correct answer is: Occupying a single cell

Question **24**

Not answered

Marked out of 1

With respect to arson, which of the following is correct?

Select one:

- ☐ Males offend more than females
- ☐ Alcohol misuse is very common
- ☐ Women receive disproportionately more psychiatric disposals
- ☐ There is an association with low intellectual functioning
- ☐ All of the listed options

Your answer is incorrect.

Explanation:

Arson is a criminal act in which one willfully and maliciously sets fire to or aids in setting fire to a structure, dwelling, or property of another. Most arsonists are male, aged late adolescence-early adulthood, unmarried, poorly educated, living alone, and unemployed or unskilled labourers. Psychiatric disorders are over-represented in arsonists compared to the general population and other offender groups. Substance use disorders (especially alcohol) and personality disorders (particularly antisocial) are most common. Others include psychotic disorders and intellectual disability/low intellectual functioning.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 744.

Burton P et al. Firesetting, arson, pyromania, and the forensic mental health expert. Journal of the American Academy of Psychiatry and the Law Online. 2012 Sep 1;40(3):355-65.

The correct answer is: All of the listed options

Question 25

Not answered

Marked out of 1

Which of the following is an accurate reflection of genetic factors associated with criminal behaviour?

Select one:

- ☐ Higher MAO-A levels are associated with increased crime rates
- ☐ Callous traits are the most heritable in conduct disorder/antisocial individuals
- ☐ Criminal behaviour is entirely environmental
- ☐ Antisocial traits show 90% heritability
- ☐ Monozygotic twins are less concordant than dizygotes for antisocial behaviour

Your answer is incorrect.

Explanation:

In individuals with conduct disorder/antisocial personality, callous-unemotional traits show substantial genetic influences (heritability) and more modest environmental influences.

Ref: Henry J, Pingault JB, Boivin M, Rijdsdijk F, Viding E. Genetic and environmental aetiology of the dimensions of Callous-Unemotional traits. Psychological Medicine. 2016 Jan;46(2):405-14.

The correct answer is: Callous traits are the most heritable in conduct disorder/antisocial individuals

Question **26**

Not answered

Marked out of 1

A 33-year-old woman is repeatedly charged with shoplifting and is subsequently diagnosed with kleptomania. Kleptomania is classified as which of the following in ICD-11?

Select one:

- ☐ Impulse control disorder
- ☐ Malingering
- ☐ Paraphilic disorder
- ☐ Anxiety or fear-related disorder
- ☐ Factitious disorder

Your answer is incorrect.

Explanation:

In ICD-11, kleptomania is classified as an impulse control disorder. Kleptomania is characterised by a recurrent failure to control strong impulses to steal objects in the absence of an apparent motive (e.g., objects are not acquired for personal use or monetary gain). There is an increasing sense of tension or affective arousal before instances of theft and a sense of pleasure, excitement, relief, or gratification during and immediately after the act of stealing. After stealing items, many individuals with kleptomania experience guilt or shame for having committed a theft, but these feelings do not prevent recurrence of the behaviour. Women are significantly more likely to be diagnosed with kleptomania than men.

Ref: International Statistical Classification of Diseases and Related Health Problems (11th ed. ICD-11; World Health Organization, 2020)

The correct answer is: Impulse control disorder

Question **27**

Not answered

Marked out of 1

Which of the following is true concerning paraphilic disorders?

Select one:

- ☐ Isolated paraphilic disorders warrant detention under the Mental Health Act 1983
- ☐ They involve persistent, intense patterns of atypical sexual arousal
- ☐ They are more common in women than men
- ☐ They are defined in ICD-11, but not DSM-5
- ☐ They are never associated with dangerous behaviours

Your answer is incorrect.

Explanation:

Paraphilic disorders are characterised by persistent and intense patterns of atypical sexual arousal, manifested by sexual thoughts, fantasies, urges, or behaviours, the focus of which involves others whose age or status renders them unwilling or unable to consent and on which the person has acted or by which he or she is markedly distressed. They are defined in both DSM-5 and ICD-11. Paraphilic disorders are more common in males than females. There is overlap between these disorders, sex offending, and inappropriate sexual behaviour.

Ref: International Statistical Classification of Diseases and Related Health Problems (11th ed. ICD-11; World Health Organization, 2020)

Sample D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 504-505.

The correct answer is: They involve persistent, intense patterns of atypical sexual arousal

Question **28**

Not answered

Marked out of 1

A death resulting from intentional act to kill, but because of provocation, is termed

Select one:

- ☐ The McNaughton rule
- ☐ Actus reus
- ☐ Involuntary culpable homicide
- ☐ Automatism
- ☐ Voluntary culpable homicide

Your answer is incorrect.

Explanation:

Manslaughter/culpable homicide may be voluntary or involuntary. Voluntary culpable homicide is defined as killing with intent to murder, but a partial defence applies (e.g. suicide pact, severe provocation). Involuntary culpable homicide applies when the conduct was grossly negligent, given the risk of death, and did kill; or when the conduct was an unlawful act involving a danger of harm and resulted in death.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 732.

The correct answer is: Voluntary culpable homicide

Question 29

Not answered

Marked out of 1

Which of the following syndromes is associated with military prisoners?

Select one:

- ☐ Werewolf syndrome
- ☐ Battered baby syndrome
- ☐ Ganser syndrome
- ☐ Munchausen syndrome
- ☐ Threat control override syndrome

Your answer is incorrect.

Explanation:

In 1898, Ganser described four criminals showing several common features, including clouding of consciousness with disorientation, auditory and visual hallucinations (or pseudo-hallucinations), amnesia for the period during which the symptoms manifest, conversion symptoms and recent head injury, infection, and severe emotional stress. Approximate answers suggest that the patient understands questions but appears to be deliberately avoiding the correct answer. Ganser believed this to be a hysterical condition with the unconscious production of symptoms to avoid a court appearance. Many now believe that Ganser syndrome is indicative of either an organic or a psychotic state, rather than hysteria as originally believed. A similar condition of approximate answers is found in those consciously feigning illness - this should be called malingering or factitious disorder according to the nature of the gain. Ganser syndrome and malingering/factitious disorder are often confused despite the conscious basis for the latter.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. pp. 68-69.

The correct answer is: Ganser syndrome

Question 30

Not answered

Marked out of 1

Males are usually more antisocial than females. Which of the following is true in this regard?

Select one:

- ☐ Male excess is less pronounced for more violent crimes
- ☐ Psychopathy is more common in females
- ☐ Before puberty girls are more antisocial than boys
- ☐ Around age 15, drug and alcohol related offences are equally common in boys and girls
- ☐ Females are exposed to more risk factors for antisocial behaviour than males

Your answer is incorrect.

In mid-adolescence, the antisocial behaviours have equal sex distribution. At all other ages, males predominate. This may be due to higher exposure to risk factors in males. The divide between the genders is much more pronounced for violent crimes than petty offences.

The correct answer is: Around age 15, drug and alcohol related offences are equally common in boys and girls

Question **31**

Not answered

Marked out of 1

A 12-year-old boy is brought to a Child Assessment Clinic following incidents of shoplifting. If he is diagnosed to have a psychiatric disorder, it is most likely to be

Select one:

- ☐ Oppositional defiant disorder
- ☐ ADHD
- ☐ Conduct-dissocial disorder
- ☐ Bipolar disorder
- ☐ Depression

Your answer is incorrect.

Explanation:

In ICD-11, conduct-dissocial disorder is characterised by a repetitive and persistent pattern of behaviour in which the basic rights of others or major age-appropriate societal norms, rules, or laws are violated such as aggression towards people or animals; destruction of property; deceitfulness or theft; and serious violations of rules. Deceitfulness or theft may include stealing items of value (e.g., shoplifting, forgery), lying to obtain goods or favours or to avoid obligations, and/or breaking into someone's house, building, or car.

Ref: International Statistical Classification of Diseases and Related Health Problems (11th ed. ICD-11; World Health Organization, 2020)

The correct answer is: Conduct-dissocial disorder

Question **32**

Not answered

Marked out of 1

Which of the following offences is almost exclusively committed by women?

Select one:

- ☐ Child abuse
- ☐ Arson
- ☐ Combined Suicide-Homicide
- ☐ Fraud
- ☐ Child-stealing

Your answer is incorrect.

Child stealing is exclusively reported in female offenders. The act can occur in previously delinquent young women who needs a child as a comfort, or to manipulate a relationship.

The correct answer is: Child-stealing



Question **33**

Not answered

Marked out of 1

Sex offences are:

Select one:

- ☐ Only committed by adults.
- ☐ Much the same across all sections of the society
- ☐ Never committed by people with mental illness
- ☐ More commonly committed by people with learning difficulties
- ☐ Committed more by women than men

Your answer is incorrect.

Most sex offenders do not have a major mental illness (Grubin and Gunn, 1991). However, people with schizophrenia or related psychoses may commit sex offences or show abnormal sexual behaviour. Sexual offences constitute 1% of all offences: almost exclusively men offend in this regard. The offenders are widely heterogeneous and no consistent or absolute correlation is seen between a sexual offence and sexual disorder.

The correct answer is: More commonly committed by people with learning difficulties

Question **34**

Not answered

Marked out of 1

Which of the following is a core feature of borderline personality disorder?

Select one:

- ☐ Aggression
- ☐ Lack of empathy
- ☐ Magical thinking
- ☐ Efforts to avoid real or imagined abandonment
- ☐ Depression

Your answer is incorrect.

Explanation:

Features of borderline personality disorder include unstable self-image; unstable personal relationships; impulsive behaviour in areas such as personal safety and substance misuse; self-harm, suicidal threats, or gestures; affective instability due to marked reactivity of mood; episodes of micro-psychosis; chronic feelings of emptiness; frantic efforts of avoid real or imagined abandonment; and inappropriate, intense anger.

Ref: American Psychiatric Association. Diagnostic and statistical manual of mental disorders (DSM-5®). American Psychiatric Pub; 2013 May 22.

The correct answer is: Efforts to avoid real or imagined abandonment

Question **35**

Not answered

Marked out of 1

According to large follow-up studies, the most important mediator of violence in patients with mental illness is

Select one:

- ☐ Discharge from hospital
- ☐ Cognitive impairment
- ☐ Presence of strangers in the immediate neighbourhood
- ☐ Lack of formal education
- ☐ Substance abuse

Your answer is incorrect.

Explanation:

Substance use is common in mentally disordered offenders. Acute intoxication is associated with disinhibition and disorganisation. Alcohol and drug use hugely inflates the risk of violence among mentally disordered offenders. Alcohol and drug-related problems are more strongly linked to offending and violence than any other mental disorders.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 747.

Varshney M et al. Violence and mental illness: what is the true story?. J Epidemiol Community Health. 2016 Mar 1;70(3):223-5.

The correct answer is: Substance abuse

Question 36

Not answered

Marked out of 1

Insane automatisms due to sleepwalking may be offered as a defence in court. Which of the following features makes a diagnosis of sleepwalking more likely?

Select one:

- ☐ Dream-like recollection of the offence
- ☐ Offence occurs within 2 hours of going to sleep
- ☐ Awakening with intense feelings of anxiety and fear
- ☐ Attempts are made to conceal the offence
- ☐ Narrative-like recollection of the offence

Your answer is incorrect.

Explanation:

If an individual commits an offence when their body is not under the control of their mind, (e.g. when asleep), they are not guilty of the offence. Legally, this is called an automatism. Insane automatisms are due to an intrinsic cause (e.g. sleepwalking, brain tumours, epilepsy) and result in acquittal on the grounds of insanity. Sleepwalking involves complex, automatic behaviours (automatisms) such as wandering, attempting to dress or undress, carrying objects, eating, urinating in unusual places, and rarely driving a car or sexual behaviour. Behaviours usually occur 15-120 minutes following sleep onset. The activity never appears intentional or planned, and only rarely does aggressive behaviour occur. The person usually has no recollection of the episode. If awakened during the episode, they are usually confused and disoriented.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 460, 778.

The correct answer is: Offence occurs within 2 hours of going to sleep

Question **37**

Not answered

Marked out of 1

There is a sound evidence base for the treatment of sex offenders with

Select one:

- ☐ Lithium
- ☐ Androgen replacement
- ☐ Counselling
- ☐ Cognitive-behavioural therapy
- ☐ Dynamic psychotherapy

Your answer is incorrect.

Explanation:



Risk Management and Treatment of Sex Offenders

Key monitoring techniques

- Talking to the offender
- Talking to people who know the offender
- Overt or covert surveillance
- Use of closed-circuit television
- Urine drug testing
- Alcohol breath testing
- Polygraph interviews
- Checking the offender's use of social media/internet
- Checking the offender's association with others

Supervision strategies

- Electronic monitoring
- Exclusion zones
- Geographical restrictions
- Preventing contact with potential victims
- Notification of whereabouts and other personal details
- Preventing contact with sex offenders or previous victims
- Preventing use of electronics
- Detention in institutions
- Limiting housing options
- Mandatory assessment
- Limiting alcohol or drug use

Specific treatments

Psychological

- Group CBT (treatment of choice)
- Risk, Needs, Responsivity approach (deliver treatment to riskier individuals, target factors associated with recidivism, and match learning style of offenders)
- Good Lives model (achieve needs/goals in prosocial ways)

Pharmacological

- Antiliberals (e.g. cyproterone acetate, gonadotropin-releasing hormone agonists)
- SSRIs, oestrogens

Adapted from Russell and Darjee, 2013

SPMM COURSE



Ref: Russell K, Darjee R. Practical assessment and management of risk in sexual offenders. Advances in psychiatric treatment. 2013 Jan;19(1):56-66.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 741.

Khan O et al. Pharmacological interventions for those who have sexually offended or are at risk of offending. Cochrane Database of Systematic Reviews. 2015(2).

The correct answer is: Cognitive-behavioural therapy

Question 38

Not answered

Marked out of 1

Of those listed, the greatest risk factor for suicide in a prisoner is

Select one:

- ☐ Female sex
- ☐ Elderly age
- ☐ History of substance misuse
- ☐ History of arson
- ☐ History of sexual offences

Your answer is incorrect.

Explanation

Prison suicide and self-harm has been rising in the UK. The most common method of suicide in prison is hanging. The majority of suicides occur within 6 months of imprisonment; life-sentence prisoners are over-represented. Occupying a single cell/solitary confinement is a particularly strong risk factor for both self-harm and suicide. Remand prisoners, young offenders, and those with a history of substance misuse and violent offences are also at higher risk of suicide.

Ref: Favril L et al. Risk factors for self-harm in prison: a systematic review and meta-analysis. The Lancet Psychiatry. 2020 Aug 1;7(8):682-91.

Burki T. Crisis in UK prison mental health. The Lancet Psychiatry. 2017 Dec 1;4(12):904.

The correct answer is: History of substance misuse

Question 39

Not answered

Marked out of 1

Which of the following is classified under sane automatism?

Select one:

- ☐ Epilepsy
- ☐ Sleepwalking
- ☐ Sleep terrors
- ☐ Brain tumours
- ☐ All of the listed options

Your answer is incorrect.

Explanation:

If an individual commits an offence when their body is not under the control of their mind, (e.g. when asleep), they are not guilty of the offence. Legally, this is called an automatism. In England and Wales, two legal types of automatism are recognised: insane and sane (automatism simpliciter). The distinction is based on whether the behaviour is likely to recur. Insane automatisms are due to an intrinsic cause (e.g. sleepwalking, brain tumours, epilepsy) and result in acquittal on the grounds of insanity. Sane automatisms are due to an extrinsic cause (e.g., confusional states, concussion, reflex actions after bee stings, dissociative states, sleep terrors, and hypoglycaemia) and result in complete acquittal.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 778.

The correct answer is: Sleep terrors

Question 40

Not answered

Marked out of 1

The following is true regarding offending behaviour in patients with learning disabilities

Select one:

- ☐ Recidivism is lower compared to offenders without LD.
- ☐ People with LD are more likely to commit violent offences
- ☐ Fire setting is the most common offense
- ☐ Prevalence of learning disability in prisons is nearly 40%
- ☐ Offending rates are higher in profound LD.

Your answer is incorrect.

The most common offences are property offences. The element of severe violence is rare in LD offenders. Prison prevalence rates vary from 2-40%. Most offenders have mild or borderline LD. Recidivism is comparable to those without LD, and may be related to difficulties in understanding cautions and probations.

The correct answer is: Prevalence of learning disability in prisons is nearly 40%



© 2023 SPMM Course Limited, UK.

[Refund Policy](#) | [Terms & Conditions](#)

