

Started on	Sunday, 12 February 2023, 8:30 PM
State	Finished
Completed on	Sunday, 12 February 2023, 8:30 PM
Time taken	10 secs
Marks	0/36
Grade	0 out of 100

Question 1

Not answered

Marked out of 1

Studies concerning supervised discharge and related arrangements agree on which of the following?

Select one:

- ☐ Frequent contact with services is unnecessary for an efficient supervised discharge
- ☐ No reduction occurs in hospitalisation rates after a supervised discharge
- ☐ Supervised discharge is most effective for those with non-affective psychosis
- ☐ No reduction occurs in rates of violence after a supervised discharge
- ☐ Use of existing powers of supervised discharge is widespread and uniform

Your answer is incorrect.

Supervised discharges are most effective in schizophrenia patients, not personality disorders. This reduces hospitalisations and violence rate but frequent contacts are necessary.

The correct answer is: Supervised discharge is most effective for those with non-affective psychosis

Question 2

Not answered

Marked out of 1

A 24-year-old man admits having fantasies of being beaten with a sharp spiny weapon while having sexual arousal. He has asked some of his sexual partners to kick him with shoe heels to achieve sexual pleasure. This is consistent with

Select one:

- ☐ Voyeurism
- ☐ Frotteurism
- ☐ Fetishism
- ☐ Sadism
- ☐ Masochism

Your answer is incorrect.

Masochism is a paraphilia where the subject achieves sexual gratification through fantasies or actual acts involving humiliation or infliction of pain to self.

The correct answer is: Masochism



Question 3

Not answered

Marked out of 1

Which of the following is an actuarial tool for violence risk assessment?

Select one:

- ☐ Hare Psychopathy Checklist
- ☐ HCR 20
- ☐ Violence Risk Appraisal Guide
- ☐ Health of Nations Outcome Scale
- ☐ None of the listed options

Your answer is incorrect.

Explanation:

The Violence Risk Appraisal Guide (VRAG) is an actuarial instrument for violence risk assessment. An actuarial instrument is a statistical approach based on multivariate analyses of factors in samples of forensic patients or prisoners to determine which predict further violence. It is reliant on historical factors and considered by some to be inflexible and unsuitable to inform risk management.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 748.

The correct answer is: Violence Risk Appraisal Guide

Question 4

Not answered

Marked out of 1

Which of the following reduces an offender's responsibility for a crime?

Select one:

- ☐ Experiencing PTSD after the offence
- ☐ Committing the offence in response to a delusion
- ☐ Having amnesia for the offence
- ☐ Experiencing numbing and dissociation before the offence
- ☐ Having a history of epilepsy

Your answer is incorrect.

Symptoms preventing the formation of mental thought processes required for mens rea include but not limited to specific delusions relevant to the situation, an abnormal mood state (severe depression or severe mania) or associated psychomotor changes, or impaired concentration and severe cognitive impairment.

The correct answer is: Committing the offence in response to a delusion

Question 5

Not answered

Marked out of 1

The most frequently used assessment tool for personality disorders seen in forensic practice is

Select one:

- ☐ SORAG
- ☐ PCL-R
- ☐ HCR-20
- ☐ VRAG
- ☐ SARA

Your answer is incorrect.

PCL-R (Hare) is a scale to diagnose Psychopathy, informs risk assessment and treatment decisions. 0 - 40 score range; 0-2 for each item; 20 items in total. Cut off of 25 used to diagnose psychopathy. In strictest sense, PCL was not designed to be an actuarial tool for risk assessment on its own

The correct answer is: PCL-R



Question 6

Not answered

Marked out of 1

All of the following are risk factors for expression of antisocial behaviour except

Select one:

- ☐ Both parents being full-time employed
- ☐ High social deprivation
- ☐ Low social cohesion in the neighbourhood
- ☐ Multiple caregivers during development
- ☐ Low parental affection

Your answer is incorrect.

Parental unemployment, rather than full-time employment, predicts antisocial behaviour.

The correct answer is: Both parents being full-time employed



Question 7

Not answered

Marked out of 1

What is the best predictor of future sexual offences?

Select one:

- ☐ Mental illness
- ☐ Previous non-sexual offences
- ☐ History of illicit drug use
- ☐ Previous sexual offences
- ☐ Lack of victim empathy

Your answer is incorrect.

Marshall's follow-up study (1994) of nearly 13 000 offenders of all types released from prison in England and Wales in 1987 found that men with a history of sex offences were more likely to be convicted of a sex offence over the next four years than those with no history of sexual offending. 'Rapists tend to be more criminally versatile than child molesters and are more likely to reoffend non-sexually than sexually. Grubin, D. and Wingate, S. (1996) 'Sexual offence recidivism: Prediction versus understanding.' Criminal Behaviour and Mental Health 6, 349-359.

The correct answer is: Previous sexual offences

Question 8

Not answered

Marked out of 1

What is the most reliable predictor to identify whether mothers will murder their children?

Select one:

- ☐ Family income
- ☐ History of substance misuse
- ☐ Previous history of sexual abuse in the mother
- ☐ Maternal age
- ☐ There are no reliable predictors

Your answer is incorrect.

Explanation:

Though teenage mothers and mothers with affective disorders may be at higher risk, there are no reliable predictors to identify whether mothers will murder their children.

Flynn SM et al. Filicide: mental illness in those who kill their children. PloS one. 2013 Apr 4;8(4):e58981.

The correct answer is: There are no reliable predictors



Question 9

Not answered

Marked out of 1

What form of domestic violence is most prevalent?

Select one:

- ☐ Homosexual couples -males
- ☐ Homosexual couples -females
- ☐ Heterosexual females assaulting children
- ☐ Heterosexual females assaulting males
- ☐ Heterosexual males assaulting females

Your answer is incorrect.

Males assaulting females is most prevalent. Surveys, which include sexual violence and a broader range of abuse, indicate that domestic violence affects 1 in 4 women at some time in their lives. Women are victims of 70% of domestic violence.

The correct answer is: Heterosexual males assaulting females



Question 10

Not answered

Marked out of 1

Which of the following is true about sane automatism?

Select one:

- ☐ It includes epilepsy
- ☐ The defendant had control over their mind during the offence
- ☐ Intent to harm need not be excluded
- ☐ It is a very common defence in psychiatric patients
- ☐ It results in complete acquittal if successful

Your answer is incorrect.

Explanation:

If an individual commits an offence when their body is not under the control of their mind, (e.g. when asleep), they are not guilty of the offence. Legally, this is called an automatism. This defence has given rise to considerable problems, stemming mainly from the acquittal which results if it succeeds. As a result, courts have restricted the scope and availability of automatism, making it a rare plea. In England and Wales, two legal types of automatism are recognised: insane and sane (automatism simpliciter). The distinction is based on whether the behaviour is likely to recur. Insane automatisms are due to an intrinsic cause (e.g. sleepwalking, brain tumours, epilepsy) and result in acquittal on the grounds of insanity. Sane automatisms are due to an extrinsic cause (e.g., confusional states, concussion, reflex actions after bee stings, dissociative states, sleep terrors, and hypoglycaemia) and result in complete acquittal.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 778.

Mackay RD. An anatomy of automatism. Medicine, Science and the Law. 2015 Jul;55(3):150-5.

The correct answer is: It results in complete acquittal if successful

Question 11

Not answered

Marked out of 1

A 31-year-old man stalks his ex-wife with the hope that a reconciliation of their broken relationship will occur. He is most appropriately categorised as what type of stalker?

Select one:

- ☐ Incompetent suitors
- ☐ Rejected stalkers
- ☐ Resentful stalkers
- ☐ Predatory stalkers
- ☐ Intimacy seekers

Your answer is incorrect.

Explanation:

Typology of stalking perpetrators:

- **Rejected stalkers:** pursue victims in order to reverse, correct, or avenge rejection (e.g. divorce, termination of relationship); previous sexual relationship; want to keep the victim in their lives
- **Intimacy seekers:** infatuated with their victims and believe their victims are infatuated with them; no previous sexual relationship
- **Incompetent suitors:** despite poor social or courting skills, have a fixation on (or in some cases a sense of entitlement to) an intimate relationship with those who have attracted their amorous interest
- **Resentful stalkers:** pursue a vendetta because of a sense of grievance against the victim; motivated by a desire to frighten and distress; not following a sexual relationship
- **Predatory stalkers:** initially spy on victims as a precursor to sexual assault; victim is often a stranger or acquaintance

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 742.

The correct answer is: Rejected stalkers

Question 12

Not answered

Marked out of 1

A newly appointed executive of a forensic secure hospital board is exploring ways of improving the security of the units. Which of the following is best described as a measure of relational security?

Select one:

- ☐ Provision of meaningful activities to patients
- ☐ Pre-employment checks on staff members
- ☐ Minimising the use of temporary members of staff
- ☐ Investigating and learning from incidents
- ☐ Removing hanging points

Your answer is incorrect.

Explanation:

Three concepts of security:

1. **Physical/environmental** – fence/locks and other means of securing against leaving the premises; also the use of soft cutlery, removing glasses, sharps and hanging points to reduce the risk of harm to self or others
2. **Relational** – training, good communication, meaningful activities, knowing patients well and developing good relationships with them; often a lack of relational security can decrease the actual safety net provided by the physical security measures
3. **Procedural** – searches, investigating and learning from incidents, staff pre-employment checks, other clinical governance arrangements, and risk management

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 750.

The correct answer is: Provision of meaningful activities to patients

Question 13

Not answered

Marked out of 1

The most common type of delusion among stalkers is a delusion of

Select one:

- ☐ Reference
- ☐ Persecution
- ☐ Love
- ☐ Jealousy
- ☐ Grandiosity

Your answer is incorrect.

Explanation:

Stalking is a behaviour, not a mental disorder. Having said that, research suggests that 50% of stalking offenders have some sort of mental disorder. Personality disorders (predominantly cluster B), substance use disorders, schizophrenia and other psychotic disorders, and depression are the most common. A significant minority of cases occur because of erotomanic delusions (delusions of love) in which the stalker believes the victim to be in love with them.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 110, 742-743.

The correct answer is: Love

Question 14

Not answered

Marked out of 1

Of those listed, what is the most common psychiatric diagnosis in perpetrators of arson?

Select one:

- ☐ Intellectual disability
- ☐ Autism spectrum disorder
- ☐ Personality disorder
- ☐ Anxiety disorder
- ☐ Schizophrenia

Your answer is incorrect.

Explanation:

Arson is a criminal act in which one willfully and maliciously sets fire to or aids in setting fire to a structure, dwelling, or property of another. Most arsonists are male, aged late adolescence-early adulthood, unmarried, poorly educated, living alone, and unemployed or unskilled labourers. Psychiatric disorders are over-represented in arsonists compared to the general population and other offender groups. Substance use disorders (especially alcohol) and personality disorders (particularly antisocial) are most common. Others include psychotic disorders and intellectual disability/low intellectual functioning.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 744.

Burton P et al. Firesetting, arson, pyromania, and the forensic mental health expert. Journal of the American Academy of Psychiatry and the Law Online. 2012 Sep 1;40(3):355-65.

The correct answer is: Personality disorder

Question 15

Not answered

Marked out of 1

A 23-year-old man is convicted of arson. Which of the following is true?

Select one:

- ☐ Less likely to have anxiety disorders than non-fire setters
- ☐ Less likely to have sexual motive than a female arsonist
- ☐ Psychoanalytic therapy is unlikely to be helpful
- ☐ Less likely to have mood disorders than non fire setters
- ☐ More likely to get psychiatric disposal than a female arsonist

Your answer is incorrect.

Treatments for fire setting are largely behavioural or focused on intervening in the family or intrapersonal stresses that may precipitate episodes of fire setting. Behavioural treatments like aversive therapy have helped fire setters (Koles and Jenson, 1985). Other treatment methods rely on positive reinforcement with threats of punishment and stimulus satiation. No established role has been shown for psychodynamic therapy.

The correct answer is: Psychoanalytic therapy is unlikely to be helpful



Question **16**

Not answered

Marked out of 1

Methods of assessing sex offenders may include:

Select one:

- ☐ Serum folate measurement.
- ☐ Penile plethysmography
- ☐ Full blood count
- ☐ Magnetic resonance imaging
- ☐ Truth serum analysis

Your answer is incorrect.

Penile plethysmography: Because sex offenders often show high levels of denial, assessment based on self-report alone is unreliable. Penile plethysmography provides a means of determining sexual arousal by measuring increases in penile volume or circumference in response to visual cues (slides) or auditory cues (stories) (Barker and Howell, 1992; Retrieved from <http://apt.rcpsych.org/content/aptrcpsych/10/1/73.full.pdf>).

The correct answer is: Penile plethysmography

Question 17

Not answered

Marked out of 1

Which of the following terms used in medical jurisprudence refers to the (voluntary) act of crime?

Select one:

- ☐ Habeas corpus
- ☐ Mens rea
- ☐ Actus reus
- ☐ Corpus delicti
- ☐ Compos mentis

Your answer is incorrect.

Explanation:

According to criminal law, committing a socially harmful act is not the sole criterion of whether a crime has been committed. Instead, the objectionable act must have two components: actus reus and mens rea. Actus reus refers to the (voluntary) act of crime. Mens rea refers to the (evil) intent of crime. An evil intent cannot exist when an offender's mental status is so deficient, so abnormal, or so diseased to have deprived the offender of the capacity for rational intent. Neither behaviour, however harmful, nor the intent to harm, alone, is grounds for criminal action.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 843.

The correct answer is: Actus reus

Question **18**

Not answered

Marked out of 1

The genetic heritability of antisocial behaviour overlaps with which of the following conditions?

Select one:

- ☐ Neuroticism
- ☐ ADHD
- ☐ Alcohol use disorder
- ☐ Movement disorders
- ☐ School refusal

Your answer is incorrect.

Explanation:

Antisocial behaviour and alcohol & drug abuse share a number of genetic risk factors.

Ref: Szalavitz M. Genetics: No more addictive personality. Nature. 2015 Jun;522(7557):S48-9.

The correct answer is: Alcohol use disorder



Question 19

Not answered

Marked out of 1

A 35-year-old man presented with features of medically unexplained physical symptoms. During a cognitive examination, when asked to designate a 10 pounds note he described it as printed paper. Which of the following is the most likely diagnosis?

Select one:

- ☐ Couvade syndrome
- ☐ Munchausen syndrome
- ☐ Ganser's syndrome
- ☐ Capgras syndrome
- ☐ Malingering

Your answer is incorrect.

In the original description of Ganser's syndrome, Ganser described a patient as follows: "when asked to read the paper he read out in a senseless fashion, Bismarck, King Albert, Rothschild. . . He designated a dollar bill, a mark and a fifty cent piece alike as a mark. A five-mark bill he described as printed paper, whose value he was not able to recognise." (Excerpt from Dwyer & Reid; Lancet 2004; 364: 471 -73)

The correct answer is: Ganser's syndrome

Question 20

Not answered

Marked out of 1

A patient admitted to a psychiatric intensive care unit has delusions of passivity. He believes that his "words are spoken by the devil's inner soul" and that his actions are controlled by the devil. When the devil takes over the patient's body, he punches his doctor who he believes is trying to poison him with ECT. This constellation of symptoms is called

Select one:

- ☐ Passivity loss syndrome
- ☐ Malingering
- ☐ Passive activity syndrome
- ☐ Threat-control override symptoms
- ☐ None of the listed options

Your answer is incorrect.

Explanation:

Threat-control override symptoms in which patients feel like they are being threatened or controlled (delusions of influence, delusions of threat, command hallucinations, threatening content of auditory hallucinations) have been found to be associated with violence. Having said that, most patients with these symptoms are never violent.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 746.

Haddock G et al. Psychotic symptoms, self-harm and violence in individuals with schizophrenia and substance misuse problems. Schizophrenia research. 2013 Dec 1;151(1-3):215-20.

The correct answer is: Threat-control override symptoms

Question **21**

Not answered

Marked out of 1

Which of the following is least likely to increase the risk of violence in those with schizophrenia?

Select one:

- ☐ History of victimisation
- ☐ History of conduct disorder
- ☐ Negative symptoms
- ☐ Depressive symptoms
- ☐ Persecutory delusions

Your answer is incorrect.

Explanation:

Negative symptoms, including blunting of affect and poverty of speech, have been associated with less violence in individuals with schizophrenia.

Ref: Buchanan A et al. Correlates of future violence in people being treated for schizophrenia. American Journal of Psychiatry. 2019 Sep 1;176(9):694-701.

The correct answer is: Negative symptoms



Question **22**

Not answered

Marked out of 1

A married man was accused of indecent exposure and you have examined his mental state. You have to write a report. What will reduce his chance of re-offending?

Select one:

- ☐ Psychosexual assessment
- ☐ Cyproterone acetate
- ☐ Court appearance
- ☐ Penile plethysmography
- ☐ Polygraph assessment

Your answer is incorrect.

Court appearance and sentencing would sometimes reduce the risk of recidivism although this obviously depends on the offender's personality. Multiagency working (e.g. co-ordinated risk management MAPPA- probation, housing, social services, psychiatry etc.) and Sex Offender Treatment Programmes (SOTP) are the most important ways of reducing the risk of recidivism. Psychiatric treatment is unlikely to be indicated in most sex offenders- including those convicted of indecent exposure. Polygraph assessment, psychosexual assessment, penile plethysmography are all helpful in risk assessment but not linked to reduction in recidivism, anti-androgens (this might reduce the risk of recidivism in serious sex offenders, but would obviously not be indicated for an offender whose offence is indecent exposure), SSRIs (these might be helpful for those who are depressed or obsessional, but only indicated in a small proportion).

The correct answer is: Court appearance

Question **23**

Not answered

Marked out of 1

The MacArthur Competence Assessment Tool is most commonly used to evaluate which of the following?

Select one:

- ☐ Fitness to plead
- ☐ Withdrawal of life support
- ☐ Parental decision making
- ☐ Prisoners' capacity to vote
- ☐ Gillick competence

Your answer is incorrect.

Explanation:

In the United States, Hoge et al. developed and validated the MacArthur Structured Assessment of the Competence of Criminal Defendants to assess a defendant's fitness to plead based on legal theory of competence. This 47-item measure was subsequently reduced to a 22-item clinical instrument, the MacArthur Competence Assessment Tool – Criminal Adjudication (MacCAT-CA). The MacCAT-CA was then modified to test abilities according to the criteria for fitness to stand trial in England and Wales. These changes include, for example, omitting reference to juries being involved in sentencing (a feature in some U.S. jurisdictions, but not in England and Wales) and adjusting offense designations. This process led to the development of a tool specific for England and Wales, the MacArthur Competence Assessment Tool – Fitness to Plead.

Ref: Akinkunmi AA. The MacArthur Competence Assessment Tool--Fitness to plead: A preliminary evaluation of a research instrument for assessing fitness to plead in England and Wales. Journal of the American Academy of Psychiatry and the Law. 2002.

The correct answer is: Fitness to plead

Question **24**

Not answered

Marked out of 1

Which of the following is most associated with reoffending in paedophiles?

Select one:

- ☐ Psychiatric illness
- ☐ Previous non sexual offending
- ☐ Depressed mood
- ☐ Lack of victim empathy
- ☐ Co morbid substance misuse

Your answer is incorrect.

Confirmed paedophiles have little insight into what constitutes a normal adult/child relationship; their lack of acceptance of the deleterious effects of their sexual behaviour on children is very resistant to change, reoffending is common (Seminars in practical forensic psychiatry).

The correct answer is: Lack of victim empathy



Question **25**

Not answered

Marked out of 1

The most commonly seen psychiatric side effect of cyproterone acetate is

Select one:

- ☐ Anxiety
- ☐ Depression
- ☐ Psychosis
- ☐ Mania
- ☐ Cognitive deterioration

Your answer is incorrect.

Explanation:

Anti-androgen therapy, such as cyproterone acetate, has been shown to place patients at a higher risk of neurocognitive and/or psychiatric complications, particularly depression. Depression may occur as a result of low androgen levels and/or activity, which is the major mechanism underlying anti-androgen therapy.

Ref: Alwhaibi A et al. Androgen deprivation therapy and depression in the prostate cancer patients: review of risk and pharmacological management. The Aging Male. 2022 Dec 31;25(1):101-24.

The correct answer is: Depression

Question **26**

Not answered

Marked out of 1

Which of the following personality traits is more commonly seen in elderly sexual offenders compared to non-offending older individuals?

Select one:

- ☐ Antisocial
- ☐ Dependent
- ☐ Paranoid
- ☐ Schizoid
- ☐ Schizotypal

Your answer is incorrect.

Elderly sex offenders and non-sex-offenders have similar prevalence rates of mental illness (Fazel, 2006). However, elderly sex offenders have increased rates of personality disorders (33% prevalence rates for any personality disorder), especially schizoid, obsessive-compulsive and avoidant traits. Psychol Med. 2002 Feb;32(2):219-26.

The correct answer is: Schizoid

Question **27**

Not answered

Marked out of 1

Pharmacotherapeutic options in the treatment of sex offenders include all of the following except

Select one:

- ☐ Gonadotrophin releasing hormone (GnRH) agonists
- ☐ Anti-androgens
- ☐ Selective serotonin re-uptake inhibitors (SSRIs)
- ☐ Combination of a GnRH agonist and SSRI
- ☐ Sildenafil citrate

Your answer is incorrect.

Explanation:

Since the 1990s, CBT has been the main modality of psychotherapeutic treatment for people convicted of sexual offences in the UK. Systematic use of medication only began to emerge in the early 2000s. The three classes of medications most often prescribed for problematic sexual behaviour include the SSRIs and two types of testosterone-lowering agents: the anti-androgens and the gonadotropin releasing hormone (GnRH) agonists. Treatment guidelines propose a hierarchical treatment protocol in which treatment is largely based on risk and progresses from psychotherapy (offered at all stages) to an ultimate stage where the combination of a GnRH agonist with an anti-androgen and/or SSRI is recommended. GnRH agonists are often considered preferable to anti-androgens whenever a testosterone-lowering medication is indicated as their side-effect profile appears to be relatively milder, and their effect more consistent.

Ref: Winder B et al. The pharmacologic treatment of problematic sexual interests, paraphilic disorders, and sexual preoccupation in adult men who have committed a sexual offence. International Review of Psychiatry. 2019 Feb 17;31(2):159-68.

The correct answer is: Sildenafil citrate

Question **28**

Not answered

Marked out of 1

Which of the following offences is the most common forensic presentation of conduct disorder?

Select one:

- ☐ sexual offence
- ☐ shoplifting
- ☐ arson
- ☐ robbery
- ☐ burglary

Your answer is incorrect.

Stealing without confronting the victim (e.g. shoplifting) is the most common offence that brings children with conduct disorder to the attention of a forensic setting.

The correct answer is: shoplifting



Question **29**

Not answered

Marked out of 1

The most common offence in people with learning disability is

Select one:

- ☐ Offences against property
- ☐ Fire setting
- ☐ Fight
- ☐ Stealing
- ☐ Sexual offence

Your answer is incorrect.

It is asserted that people with LD are more likely to commit sex or fire-raising offences with property offences being the most common (Hodgins).

The correct answer is: Offences against property



Question 30

Not answered

Marked out of 1

Which of the following is most likely to be associated with amnesia for offending?

Select one:

- ☐ Automotive theft
- ☐ Psychotic symptoms at the time of the offence
- ☐ Shoplifting
- ☐ Elevated mood in the perpetrator
- ☐ Extreme emotional arousal during the offence

Your answer is incorrect.

Explanation:

There may be three different explanations for memory loss in offenders. The first explanation contends that, during the time of the crime, some offenders suffer from a temporary (or permanent) brain dysfunction that prevents or undermines the storage of criminal events in memory (organic amnesia). The second explanation holds that some offenders are in an extreme emotional condition (e.g., rage) when committing a crime, particularly a violent crime. Later, the offender is unable to remember the crime because of a mismatch in emotional state between the encoding of crime-related events and the retrieval of such events (dissociative amnesia). The third explanation for crime-related amnesia is that a considerable number of offenders are pretending to be unable to remember crime-related details (feigned amnesia).

Ref: Jelcic M. Testing claims of crime-related amnesia. *Frontiers in Psychiatry*. 2018 Nov 21;9:617.

The correct answer is: Extreme emotional arousal during the offence

Question **31**

Not answered

Marked out of 1

A 31-year-old man accused of a homicide has a medical history of somnambulism. Which of the following is true about this condition?

Select one:

- ☐ The subject is actually awake when sleepwalking
- ☐ Sleepwalking is more common in the elderly than in children
- ☐ This diagnosis can be entertained as a plea for diminished responsibility
- ☐ An offence committed when sleepwalking is usually considered under sane automatism
- ☐ It occurs mostly during REM sleep

Your answer is incorrect.

Claims of diminished responsibility due to sleepwalking have been upheld in various jurisdictions in cases of attempted murder, grievous injuries and very rarely homicides.

The correct answer is: This diagnosis can be entertained as a plea for diminished responsibility



Question **32**

Not answered

Marked out of 1

Which one of the following is a structured risk assessment tool?

Select one:

- ☐ PCL-R
- ☐ STATIC
- ☐ VRAG
- ☐ SARA
- ☐ VRS

Your answer is incorrect.

Explanation:

The Spousal Assault Risk Assessment (SARA; Kropp 1995) is a 20-item structured risk assessment tool that can be used to help gauge the risk of future violence when there has been a history of domestic abuse, or the victim of a sexual offense has been a partner or a partner's child.

Ref: Russell K, Darjee R. Practical assessment and management of risk in sexual offenders. Advances in psychiatric treatment. 2013 Jan;19(1):56-66.

The correct answer is: SARA

Question **33**

Not answered

Marked out of 1

Factors that increase the risk of assault in association with stalking behaviour include all of the following except

Select one:

- ☐ Presence of narcissistic personality traits
- ☐ History of offending behaviour
- ☐ Substance misuse
- ☐ Female stalker
- ☐ Making threats of violence

Your answer is incorrect.

Factors likely to increase risk of assault in stalking (Palarea 1999, Mullen 2000) include substance use, male gender, making threats of violence, presence of personality disorder and history of non-compliance with treatment for mental health problems.

The correct answer is: Female stalker



Question **34**

Not answered

Marked out of 1

A 33-year-old man is charged with murder. His counsel pleads not guilty on the basis of sleepwalking. Sleepwalking may be accepted as an automatism provided that

Select one:

- ☐ The perpetrator has no previous history of sleepwalking
- ☐ The perpetrator was able to narrate the event on waking up
- ☐ The homicide took place within the first 2 hours of sleeping
- ☐ The perpetrator has partial recall for the act
- ☐ The perpetrator wakes up in the middle of the act with panic and screams

Your answer is incorrect.

Explanation:

If an individual commits an offence when their body is not under the control of their mind, (e.g. when asleep), they are not guilty of the offence. Legally, this is called an automatism. Insane automatisms are likely to recur and are due to an intrinsic cause (e.g. sleepwalking, brain tumours, epilepsy). Insane automatisms result in acquittal on the grounds of insanity. Sleepwalking involves complex, automatic behaviours (automatisms) such as wandering, attempting to dress or undress, carrying objects, eating, urinating in unusual places, and rarely driving a car or sexual behaviour. Behaviours usually occur 15-120 minutes following sleep onset. The activity never appears intentional or planned, and only rarely does aggressive behaviour occur. The person usually has no recollection of the episode. If awakened during the episode, they are usually confused and disoriented.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 460, 778.

The correct answer is: The homicide took place within the first 2 hours of sleeping

Question 35

Not answered

Marked out of 1

In forensic psychiatry, the Penrose hypothesis refers to which of the following?

Select one:

- ☐ The relationship between homicide rates and psychiatric bed numbers
- ☐ The relationship between the prison population and income inequality
- ☐ The relationship between the prison population and psychiatric bed numbers
- ☐ The relationship between income inequality and crime
- ☐ The relationship between the prison population and suicide rates

Your answer is incorrect.

Explanation:

Penrose proposed an inverse relationship between the relative number of psychiatric beds available to a population and its total number of prisoners, based on calculations from cross-sectional study of 18 countries.

Ref: Toynbee M. The Penrose hypothesis in the 21st century: revisiting the asylum. Evidence-based mental health. 2015 Aug 1;18(3):76.

The correct answer is: The relationship between the prison population and psychiatric bed numbers

Question 36

Not answered

Marked out of 1

Which of the following is considered to be a test for fitness to give evidence in court?

Select one:

- ☐ Ability to challenge a juror
- ☐ Understanding the concept of 'guilty mind'
- ☐ Ability to instruct counsel
- ☐ Understanding the concept of criminal responsibility
- ☐ Ability to understand questions posed by the court

Your answer is incorrect.

Explanation:

Though there is an absence of medical literature assessing the capacity to give evidence, legal competence in the UK requires that the person be able to understand questions and able to give answers that can be understood. Competence to give evidence is a decision for the judge, who may ask for the opinion of a psychiatrist or psychologist. The law presumes that ordinary individuals are competent and can be compelled to testify. Individuals may lack capacity to give evidence because of age or mental disorder.

Ref: Jones R, Elliott T. Capacity to give evidence in court: issues that may arise when a client with dementia is a victim of crime. Psychiatric Bulletin. 2005 Sep;29(9):324-6.

The correct answer is: Ability to understand questions posed by the court



© 2023 SPMM Course Limited, UK.

[Refund Policy](#) | [Terms & Conditions](#)

