

Started on	Sunday, 12 February 2023, 8:26 PM
State	Finished
Completed on	Sunday, 12 February 2023, 8:27 PM
Time taken	13 secs
Marks	0/60
Grade	0 out of 100

Question 1

Not answered

Marked out of 1

Which of the following best describes cognitive analytical therapy?

Select one:

- ☐ It is based on conflicts and role disturbances seen in immediate environment
- ☐ It is built on a hierarchy of goals
- ☐ Hypothesis testing is carried out by the patient as home work
- ☐ It is a structured therapy focussing on cognitive and emotional functioning along with reformulation
- ☐ It focuses on distorted cognitions

Your answer is incorrect.

CAT combines Cognitive Theory and Psychoanalytical principles in therapy and encourages clients to draw on their ability to develop the skills to change destructive patterns of behaviour. Negative ways of thinking are explored in structured and directive ways, involving diary-keeping, reformulation letters, progress charts, etc. Retrieved from http://www.bacp.co.uk/seeking_therapist/theoretical_approaches.php

The correct answer is: It is a structured therapy focussing on cognitive and emotional functioning along with reformulation

Question 2

Not answered

Marked out of 1

Which of the following is correct about Dialectical Behavioural Therapy?

Select one:

- ☐ It requires group participation
- ☐ It comprises of a hierarchy of goals
- ☐ It includes goodbye letters
- ☐ Low frustration tolerance is a contraindication
- ☐ It is based on psychodynamic formulations

Your answer is incorrect.

DBT uses many principles of cognitive behaviour therapy (CBT), along with a strong emphasis on the use of acceptance strategies. DBT is a dynamic and provocative treatment developed by Dr Marsha Linehan for patients with severe emotional dysregulation. One of the primary concepts of dialectical behavior therapy is that self-harming behaviors are learned, and provide evidence of maladaptive coping that is reinforced in an invalidating environment. Dialectical behavior therapy recommends a hierarchy of goals to effectively address the behaviors associated with dysregulation (Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/20088061>).

The correct answer is: It comprises of a hierarchy of goals

Question **3**

Not answered

Marked out of 1

In multisystemic therapy, the treatment is usually given for a period of

Select one:

- ☐ 2 years
- ☐ 3 months
- ☐ 4 weeks
- ☐ 6 months
- ☐ 12 months

Your answer is incorrect.

In multisystemic therapy, the clinicians take on only four to six cases at a time, since the work is intensive. Treatment is usually given for three months and then stopped.

The correct answer is: 3 months



Question **4**

Not answered

Marked out of 1

The target age range for functional family therapy in the treatment of conduct disorder is

Select one:

- ☐ 11-18 years
- ☐ 6-10 years
- ☐ 5-8 years
- ☐ 10-15 years
- ☐ 18-21 years

Your answer is incorrect.

One of the best-known interventions for serious antisocial behaviour is functional family therapy. It is designed to be practicable and relatively inexpensive. The target age range is 11-18 years.

The correct answer is: 11-18 years



Question 5

Not answered

Marked out of 1

Cognitive analytical therapy usually lasts for

Select one:

- ☐ 30-32 sessions
- ☐ 10-12 sessions
- ☐ 6-8 sessions
- ☐ 12-16 sessions
- ☐ 16-24 sessions

Your answer is incorrect.

A CAT therapy is weekly for 50-60 minutes sessions. A course of therapy can be from 16-24 sessions - this is negotiated with the therapist at the start of therapy. Between one and five follow-up sessions are offered after the end of regular therapy. Again this is negotiated with the therapist.

The correct answer is: 16-24 sessions



Question 6

Not answered

Marked out of 1

A psychotherapist makes a practical link between the patient's mental symptoms and disturbing life events that either trigger or follow from the onset of the symptoms. She also defines the problem as a medical illness and thus excuses the patient from self-blame. Which of the following patients is most likely to find this approach useful?

Select one:

- ☐ A 55 year old man convicted for an impulsive sexual offence
- ☐ A 14 year old boy with conduct disorder
- ☐ A girl with the diagnosis of bulimia
- ☐ A 66 year old woman with psychotic depression
- ☐ A 45 year old man with a head injury

Your answer is incorrect.

The description suits IPT - Inter-Personal Therapy. The focus for treatment includes events such as the death of someone important (complicated bereavement), struggle with a significant other (role dispute), or some other important life change (role transition); in the relatively infrequent absence of any of these, the default focus is on interpersonal deficits. IPT has good evidence base in support of treating bulimia (<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1414693/>).

The correct answer is: A girl with the diagnosis of bulimia

Question 7

Not answered

Marked out of 1

Which one of the following statements is incorrect with respect to the principles of psychodynamic therapy?

Select one:

- ☐ Interpretation of the transference is the only means to access the unconscious mind
- ☐ A therapist gains an insight of the client's issues by interpreting the unconscious motives
- ☐ Worsening of symptoms is not uncommon during the course of psychodynamic therapy
- ☐ Humans are instinctively irrational and destructive
- ☐ Object loss can be associated with manic states

Your answer is incorrect.

In psychoanalytic psychotherapy the interpretation of the unconscious could be accessed through dreams, free-association (it is the fundamental rule of psychoanalysis wherein the patient agrees to reveal everything that comes to mind during free association, no matter how embarrassing or socially unacceptable), slips of the tongue and abreaction. Object loss is associated with both depression and manic states. Both mania and depression are considered to be reactions to loss, may have no external basis in external reality and may be denied in pathological grief. Worsening of symptoms is not uncommon during the course of psychodynamic therapy. It is referred to as negative therapeutic reaction.

The correct answer is: Interpretation of the transference is the only means to access the unconscious mind

Question 8

Not answered

Marked out of 1

Interpersonal therapy has been particularly used in the treatment of

Select one:

- ☐ Ellis
- ☐ Arthur Janov
- ☐ Aaron Beck
- ☐ Klermann & Weismann
- ☐ Anna Freud

Your answer is incorrect.

IPT was developed in the 1970s by Klermann and Weissmann. It is a time-limited treatment that lasts for 12-16 hour long weekly sessions, disorder focused which deals with symptoms in the 'here and now'. The focuses of treatment are the current interpersonal relationships and their relationship to the development of illness. The focus is on role transitions like becoming a mother, job loss and role disputes like work difficulties and relationship difficulties. IPT is indicated for the treatment of mild to moderate depressive illness, bulimia nervosa and dysthymic disorder.

The correct answer is: Klermann & Weismann

Question 9

Not answered

Marked out of 1

At the second session of psychodynamic psychotherapy, a distressed young woman says she dreamed she was sexually abused by her father. What do you do?

Select one:

- ☐ Acknowledge her distress and ask if she was abused by her father
- ☐ Explore the feelings it arouses in you
- ☐ Ignore the comment and continue with the flow the patient's talk
- ☐ Leave it to the next session
- ☐ Acknowledge her distress and ask her to talk about the dream

Your answer is incorrect.

Acknowledging distress is an important aspect of therapeutic relationship across varying modalities of therapy.

The correct answer is: Acknowledge her distress and ask her to talk about the dream



Question 10

Not answered

Marked out of 1

Which one of the following groups is particularly helpful for carers with mental illness?

Select one:

- ☐ Problem focussed groups
- ☐ Psychodynamic groups
- ☐ Tavistock model analytic group
- ☐ Support groups
- ☐ Activity groups

Your answer is incorrect.

Carers find support groups particularly helpful.

The correct answer is: Support groups



Question 11

Not answered

Marked out of 1

Eye movement desensitisation and reprocessing is a treatment for

Select one:

- ☐ Social Phobia
- ☐ Acute stress reaction
- ☐ PTSD
- ☐ Dysthymia
- ☐ Dependent / avoidant personality disorder

Your answer is incorrect.

When a trauma occurs, it seems to get locked in the nervous system with the original picture, sounds, thoughts and feelings. This material can combine fact with fantasy and with images that stand for the actual emotions. The eye movements used in EMDR "unlock the nervous system" (desensitise) and allow the brain to process correctly the experience (reprocessing) and is useful for treating established cases of PTSD.

The correct answer is: PTSD



Question 12

Not answered

Marked out of 1

Which one among the following refers to the contents of the collective unconscious that describes a pancultural representation of human experience?

Select one:

- ☐ Animus
- ☐ Archetypes
- ☐ Persona
- ☐ Shadow
- ☐ Anima

Your answer is incorrect.

Examples of pancultural representation of human experience (i.e. the Jungian archetypes) include the mother, the child, the hero, the self, the shadow, anima and animus. 'Anima' refers to the unconscious feminine aspects of the male. 'Animus' refers to the unconscious masculine aspects of the female. 'Shadow' refers to the unacknowledged aspects of oneself that includes both creative impulses and destructive urges. 'Self' refers to the main archetype that gives the personality a sense of oneness. Persona refers to the outer concealed aspects of oneself.

The correct answer is: Archetypes



Question **13**

Not answered

Marked out of 1

Interpersonal therapy has been particularly useful in the treatment of which of the following conditions?

Select one:

- ☐ Moderate depressive illness
- ☐ Psychotic disorders
- ☐ Personality disorders
- ☐ Organic brain syndromes
- ☐ Anxiety disorders

Your answer is incorrect.

Interpersonal therapy is indicated in mild to moderate depressive illness and bulimia nervosa.

The correct answer is: Moderate depressive illness



Question **14**

Not answered

Marked out of 1

According to studies by Bateman and Fonagy, psychoanalytically oriented day hospital treatment has been shown to be superior to the standard psychiatric treatment of patients with which of the following conditions?

Select one:

- ☐ Borderline personality disorder
- ☐ Autistic spectrum disorders
- ☐ Eating disorders
- ☐ Chronic fatigue syndrome
- ☐ Psychosomatic disorders

Your answer is incorrect.

An improvement in depressive symptoms, a decrease in suicidal and self-mutilation acts, reduced inpatient days, and better social and interpersonal function was noted in borderline patients under partial hospitalisation program in an RCT.

The correct answer is: Borderline personality disorder



Question 15

Not answered

Marked out of 1

In which of the following therapies does positive transference get addressed early?

Select one:

- ☐ Brief dynamic therapy
- ☐ Cognitive analytical therapy
- ☐ Psychodrama
- ☐ Play therapy
- ☐ Trauma focussed CBT

Your answer is incorrect.

In brief dynamic therapy, a time limit is usually set in the beginning. The therapy lasts up to one year, usually 20-25 sessions. The techniques employed include goal setting, focus choosing (identification of currently active problems) and active involvement of the therapist in guiding the therapy. The other features include addressing positive transference early on and addressing countertransference issues, working through therapy termination.

The correct answer is: Brief dynamic therapy

Question 16

Not answered

Marked out of 1

Which of the following predicts poor response to psychodynamic psychotherapy?

Select one:

- ☐ Lower socioeconomic status
- ☐ History of multiple acting out incidents
- ☐ Ongoing multiple sexual relationships
- ☐ Average intelligence
- ☐ History of multiple previous relationships

Your answer is incorrect.

Lower socioeconomic status or average IQ need not be poor predictors of response by themselves. Multiple uncontrollable 'acting out' incidents may predict recurrences and hence poor response to psychoanalysis.

The correct answer is: History of multiple acting out incidents



Question 17

Not answered

Marked out of 1

In psychoanalytic terms, condensation refers to

Select one:

- ☐ Combining various unconscious impulses into a single image while dreaming
- ☐ Transfer of energy from an original object to a symbolic representation of the object
- ☐ Avoidance of unconscious impulses
- ☐ Enacting dreams as they occur
- ☐ Representing abstract concepts by using recognisable images

Your answer is incorrect.

Condensation is a term used in dream analysis. It refers to the process of several unconscious impulses being combined into a single image in the manifest dream content. e.g. One's father and a horrible teacher may be unified and occur as a single dreadful monster in a child's dream.

The correct answer is: Combining various unconscious impulses into a single image while dreaming

Question 18

Not answered

Marked out of 1

Which of the following terms is the key principle in the work of therapeutic community?

Select one:

- ☐ Basic assumption
- ☐ Fight-flight
- ☐ Reality Confrontation
- ☐ Dependence
- ☐ Pairing

Your answer is incorrect.

Reality Confrontation: Patients continually have other people presenting interpretations of their behaviour, so defence mechanisms such as denial, distortion and confusion become obvious. Over time, ongoing therapeutic interventions enable patients to reflect on how they are with other people, and to make changes. (Rapoport, 1960)

The correct answer is: Reality Confrontation

Question 19

Not answered

Marked out of 1

A middle-aged man with moderate depression is unwilling to take antidepressants but will have psychotherapy. Which factor would most make you choose brief psychodynamic therapy?

Select one:

- ☐ Continuous employment
- ☐ Co morbid anxiety
- ☐ History of one previous episode of depression
- ☐ Being over 50
- ☐ Wish to know more about self

Your answer is incorrect.

A wish to know more about oneself, the so-called psychological mindedness and a good degree of frustration tolerance are other positive indicators to choose brief psychodynamic therapy.

The correct answer is: Wish to know more about self



Question 20

Not answered

Marked out of 1

Mr. Z has a history of acrophobia (fear of heights). Although he is terrified of heights, his long-term ambition in life was to become a pilot and work for the air force. Which of the interventions given below is most likely to be successful for his condition?

Select one:

- ☐ Cognitive therapy
- ☐ SSRIs
- ☐ Problem solving therapy
- ☐ Systematic desensitisation
- ☐ Benzodiazepines

Your answer is incorrect.

Systematic Desensitisation (Wolpe): It consists of three steps: relaxation training, hierarchy construction, and desensitisation of the stimulus. The patient is exposed to a graded hierarchy of anxiety-provoking situations in a stepwise fashion. Other behaviour therapy techniques would include implosion (imaginal flooding) and 'flooding'. In flooding (in vivo) real life exposure takes place without any hierarchy. No relaxation exercises are used. Escaping from an anxiety-provoking experience, in fact, reinforces the anxiety through avoidance conditioning; in flooding this conditioning is targeted. The success of flooding depends on exposing patients for a reasonable duration until mastery, and calm composure are gained. Premature withdrawal will reinforce the avoidance. In implosion (in vitro) or imaginal flooding, the phobic situation is confronted through imagination, not in real life. Ref: Kaplan and Sadock's Synopsis of Psychiatry: Behavioural Sciences/Clinical Psychiatry, 10th Edition. Lippincott Williams and Wilkins 2007

The correct answer is: Systematic desensitisation

Question **21**

Not answered

Marked out of 1

Transactional analysis was first developed by

Select one:

- ☐ Eric Bern
- ☐ Anthony Ryle
- ☐ Carl Beck
- ☐ Marsha Linehan
- ☐ Carl Rogers

Your answer is incorrect.

Eric Bern first developed transactional analysis (TA). TA has elements of psychoanalytic, humanist and cognitive approaches. It is considered as a theory of personality and a systematic psychotherapy for personal growth and personal change.

The correct answer is: Eric Bern



Question **22**

Not answered

Marked out of 1

A depressed patient feels that only bad things happen to him. This is an example of

Select one:

- ☐ Elaboration
- ☐ Cognitive dissonance
- ☐ Catastrophisation
- ☐ Minimisation
- ☐ Selective abstraction

Your answer is incorrect.

Catastrophisation refers to a specific, relatively innocuous event being read in cataclysmic proportions. Negative recall bias (only bad things happen to me) is likely to be a selective abstraction.

The correct answer is: Selective abstraction



Question **23**

Not answered

Marked out of 1

The best treatment option for bulimia nervosa is

Select one:

- ☐ IPT
- ☐ Systemic family therapy
- ☐ CBT
- ☐ DBT
- ☐ Transdiagnostic enhanced CBT

Your answer is incorrect.

Fairburn et al. describe transdiagnostic CBT as an 'enhanced' version of CBT that features 2 versions of the treatment: a 'focused' form which focuses exclusively on the eating disorder and a 'broad' form which addresses some of the other issues commonly found in eating disordered patients (e.g., self-esteem, perfectionism, interpersonal problems, mood intolerance).

The correct answer is: Transdiagnostic enhanced CBT



Question **24**

Not answered

Marked out of 1

In major depressive disorder with comorbid personality disorder which of the following is true?

Select one:

- ☐ IPT is better than CBT
- ☐ CBT has same efficacy as IPT
- ☐ Neither IPT nor CBT is useful
- ☐ CBT is better than IPT
- ☐ Medication is better than CBT

Your answer is incorrect.

At least two studies have compared IPT with CBT head to head in patients with depression and comorbid personality disorders. Both concluded that IPT was less effective than CBT in such comorbid presentations. But in general patients without axis two disorders respond better to any form of psychotherapy for depression. (Rosenbluth, M et al. Depression and Personality p. 162).

The correct answer is: CBT is better than IPT



Question **25**

Not answered

Marked out of 1

Which of the following statements about CBT is incorrect;

Select one:

- ☐ CBT is helpful for treatment of alcohol dependence
- ☐ CBT and relaxation training are equally effective in generalized anxiety disorder
- ☐ With homework tasks CBT produces quicker, longer lasting progression
- ☐ CBT is better than waiting list in patients with HIV and depression
- ☐ CBT is helpful for treatment of delusions

Your answer is incorrect.

There is good evidence for the effectiveness of CBT in anxiety disorder, HIV and depression, alcohol dependence and treatment of delusions.

The correct answer is: CBT and relaxation training are equally effective in generalized anxiety disorder



Question **26**

Not answered

Marked out of 1

Which one of the following is a cognitive distortion?

Select one:

- ☐ Arbitrary inference
- ☐ Thought blocking
- ☐ Concrete thinking
- ☐ Circumstantiality
- ☐ Forced thinking

Your answer is incorrect.

Arbitrary inference refers to quickly drawing a conclusion without the requisite evidence

The correct answer is: Arbitrary inference



Question **27**

Not answered

Marked out of 1

Which of the following is not an accurate statement regarding humanistic psychotherapy?

Select one:

- ☐ Therapeutic genuineness is aimed for
- ☐ Using therapeutic alliance as a supportive element for therapy
- ☐ Accurate empathy is focussed
- ☐ Client centeredness
- ☐ Unconditional positive regard is given

Your answer is incorrect.

Humanistic psychotherapy originally was promoted as a third force in psychotherapy. There is a great deal of importance placed on the therapist-client relationship. Unlike other psychotherapies that view therapeutic relationship primarily as a means of delivering the treatment, in humanistic therapy, the relationship itself is the treatment.

The correct answer is: Using therapeutic alliance as a supportive element for therapy

Question **28**

Not answered

Marked out of 1

A couple sustained head on collision of their car but escaped with less than serious injuries. The husband does not want to discuss the accident but suffers from repeated nightmares and flashbacks. He has stopped going to work since this happened. His wife is pursuing insurance claims against the other driver. Which of the following could be a suitable treatment?

Select one:

- ☐ Psychoeducation
- ☐ Humanistic Therapy
- ☐ Rational Emotive Therapy
- ☐ Cognitive Analytical Therapy
- ☐ Eye Movement Desensitisation and Reprocessing Therapy

Your answer is incorrect.

When a trauma occurs it seems to get locked in the nervous system with the original picture, sounds, thoughts and feelings. This material can combine fact with fantasy and with 'images' that stand for the actual 'emotions'. The eye movements used in EMDR unlock the nervous system (desensitise) and allow the brain to process correctly the experience (reprocessing).

The correct answer is: Eye Movement Desensitisation and Reprocessing Therapy

Question 29

Not answered

Marked out of 1

A 19-year-old lady has a long-standing history of self-harm by taking overdoses and cutting herself in the past. She is impulsive, has low self-esteem and has constant fears of abandonment. She also has a history of sexual abuse by her stepfather in the past. Which mode of treatment is best recommended in this patient?

Select one:

- ☐ Rational emotive therapy
- ☐ Cognitive behavioural therapy
- ☐ Dialectical behavioural therapy
- ☐ Interpersonal therapy
- ☐ Cognitive analytical therapy

Your answer is incorrect.

An RCT showed that DBT was superior to community-based treatment as usual' in women with BPD associated with repeated self-harm (Linehan et al., 1991). Some benefit was retained at follow-up (Linehan et al., 1993, 1994). A further study showed similar benefits in women with BPD and substance misuse (Linehan et al., 1999) [Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1414693/>].

The correct answer is: Dialectical behavioural therapy

Question 30

Not answered

Marked out of 1

Availability of the therapist for telephone contact between appointments is one of the approaches used in

Select one:

- ☐ Interpersonal therapy
- ☐ Cognitive analytical therapy
- ☐ Cognitive behavioural therapy
- ☐ Dialectical behavioural therapy
- ☐ Rational emotive therapy

Your answer is incorrect.

Dialectical behavioural therapy (DBT) is an integrative therapy introduced in 1991 by Marsha Linehan and colleagues as a treatment for borderline personality disorder, drawing on ideas from behavioural, cognitive and psychodynamic therapies. DBT specifically aims at reduction of DSH and suicidal behaviour, improving the quality of life and personal relationships. DBT involves a variety of approaches: individual therapy, group skills training and availability of the therapist for telephone contact between appointments.

The correct answer is: Dialectical behavioural therapy



Question **31**

Not answered

Marked out of 1

Carl Rogers was associated with

Select one:

- ☐ Cognitive analytical therapy
- ☐ Dialectical behaviour therapy
- ☐ Patient centered therapy
- ☐ Cognitive behavioural therapy
- ☐ Rational emotive therapy

Your answer is incorrect.

The therapy proposed by Carl Rogers is known as Patient/Client-centered therapy. The main aim of the therapist is to help the client discover the distorted self-concepts that inhibit self-development. It is based on three cardinal principles- genuineness, accurate empathy and unconditional positive regard

The correct answer is: Patient centered therapy



Question **32**

Not answered

Marked out of 1

Reformulation letters are characteristic features of

Select one:

- ☐ Cognitive behavioural therapy
- ☐ Psychodynamic psychotherapy
- ☐ Rational emotive therapy
- ☐ Dialectical behavioural therapy
- ☐ Cognitive analytical therapy

Your answer is incorrect.

During CAT, by the third session, major maladaptive sequences and main reciprocal roles used by the patient are identified. . Between the third and fourth sessions the therapist writes a reformulation letter to the patient, which outlines the current situation and summarizes the repertoire of reciprocal roles and procedural sequences used by the patient. In the penultimate session, the therapist gives the patient another letter called goodbye letter that summarizes the therapy so far and outlines what remains to be achieved yet.

The correct answer is: Cognitive analytical therapy

Question **33**

Not answered

Marked out of 1

A child who is anxious about approaching dogs is repeatedly shown a dog when comfortably sitting on his mother's lap with an ice cream. Gradually the anxiety reduces. The principle employed here is

Select one:

- ☐ Participant modelling
- ☐ Stimulus preparedness
- ☐ Learned helplessness
- ☐ Reciprocal inhibition
- ☐ Cognitive dissonance

Your answer is incorrect.

When anxiety and a relaxed state are co-existent, then anxiety reduces - this is called reciprocal inhibition.

The correct answer is: Reciprocal inhibition



Question **34**

Not answered

Marked out of 1

The major principles on which a therapeutic community is based includes all of the following except

Select one:

- ☐ Democratisation
- ☐ Reality confrontation
- ☐ Permissiveness
- ☐ Aggregation
- ☐ Communalism

Your answer is incorrect.

The four major principles on which a therapeutic community is based are exemplified by the Henderson hospital model. According to this model, the major components are (mnemonic CPD-R) 1. Communalism (Staff are not separated from inmates by uniforms or behaviours, mutual helping and learning occurs) 2. Permissiveness (tolerating each other and realising unpredictable behaviour can happen within the community) 3. Democratisation (shared decision-making and joint running of the unit) and 4. Reality confrontation (self-deception or distortions from reality are dealt with honestly and openly by all members without formalities).

The correct answer is: Aggregation

Question **35**

Not answered

Marked out of 1

Tolerating each other and realising unpredictable behaviour can happen within the community?. This statement defines which one of the following principles of Therapeutic communities?

Select one:

- ☐ Communalism
- ☐ Democracy
- ☐ Non confrontation
- ☐ Reality confrontation
- ☐ Permissiveness

Your answer is incorrect.

Rapoport explained the key features of TCs that included communalism, democracy, permissiveness and reality confrontation. Permissiveness refers to tolerating each other and realising unpredictable behaviour can happen within the community.

The correct answer is: Permissiveness



Question **36**

Not answered

Marked out of 1

Which one among the following is not a key feature of Therapeutic Community?

Select one:

- ☐ Communalism
- ☐ Non-confrontation
- ☐ Permissiveness
- ☐ Reality confrontation
- ☐ Democracy

Your answer is incorrect.

Therapeutic communities are one of the important and successful treatment options for personality disorders. The Henderson and Cassell are well-known examples of therapeutic communities. In Henderson unit, the emphasis is on social analysis and in Cassell, psychoanalysis is primary. Tom Main (coined the term therapeutic community), Maxwell Jones (established the therapeutic community) and Rapoport explained the key features of TCs that includes communalism, democracy, permissiveness and reality confrontation.

The correct answer is: Non-confrontation

Question **37**

Not answered

Marked out of 1

A 40 yr old man with alcohol problems says that he has decided to quit drinking. He is ready to accept help and advice for this and acknowledges the harm caused by alcohol. Which of the following stages of change is he in?

Select one:

- ☐ maintenance
- ☐ action
- ☐ precontemplation
- ☐ preparation
- ☐ contemplation

Your answer is incorrect.

He is in preparatory stages of Prochaska's model.

The correct answer is: preparation



Question **38**

Not answered

Marked out of 1

Which one among the following terms refers to a patient's emotional attitude towards a therapist?

Select one:

- ☐ Interpretation
- ☐ Countertransference
- ☐ Transference
- ☐ Parapraxis
- ☐ Dream work

Your answer is incorrect.

The development, in the patient, of feelings and patterns of behaviour towards the therapist, which unconsciously partly recapitulates earlier relationships in their lives is known as transference. It usually takes a role earlier in the therapy and may even lead the patient to resist the process. Countertransference describes the equivalent reaction in the therapist towards the patient. Both were initially considered by Freud as hindrance to therapy. Parapraxis is a slip of the tongue commonly known as a Freudian slip. Occasionally it reveals unconscious meaning, particularly in affect-laden situations. Interpretation refers to a formulation of current conscious problems in terms of unconscious motives. It is also the expression of the therapist's understanding of the meaning of what is currently happening in therapy.

The correct answer is: Transference

Question 39

Not answered

Marked out of 1

A 25-year-old dysthymic lady feels that every around her are depressed but she is not unhappy. What is the possible defence mechanism?

Select one:

- ☐ Denial
- ☐ Repression
- ☐ Projection
- ☐ Projective identification
- ☐ Displacement

Your answer is incorrect.

One way of disowning thoughts and feelings is to project them onto other people. We convince ourselves that those unpleasant thoughts and feelings are coming from someone else and not from us.

The correct answer is: Projection



Question **40**

Not answered

Marked out of 1

Which of the following conditions in psychiatry was claimed to have insufficient evidence to justify supporting the use of existing therapies in practice?

Select one:

- ☐ Bipolar disorder
- ☐ Generalized anxiety disorder
- ☐ Depression
- ☐ Borderline personality disorder
- ☐ Schizophrenia

Your answer is incorrect.

Borderline personality disorder was claimed to have insufficient evidence to justify supporting the use of existing therapies in practice such as dialectical behaviour therapy, CBT, cognitive analytical therapy and mentalisation-based treatments.

The correct answer is: Borderline personality disorder

Question **41**

Not answered

Marked out of 1

All of the following are principles of motivational interviewing except

Select one:

- ☐ Developing cognitive dissonance
- ☐ avoiding argumentation
- ☐ Expressing sympathy
- ☐ Supporting self-efficacy
- ☐ rolling with resistance

Your answer is incorrect.

Expressing empathy, not sympathy, is a key process in motivational interviews.

The correct answer is: Expressing sympathy



Question **42**

Not answered

Marked out of 1

Klein's depressive position is related to the process of learning to cope with which of the following conflicts?

Select one:

- ☐ Depression
- ☐ Difficult relationships
- ☐ Hunger
- ☐ Sexual needs
- ☐ Ambivalence

Your answer is incorrect.

Melanie Klein described two positions - paranoid-schizoid and depressed position. The paranoid-schizoid position is associated with the use of splitting and projection as a defence mechanism. This position concerns an inability to perceive a whole object and splits all objects into their good and bad parts. But in the depressive position, the infant tolerates the ambiguity or ambivalence and can realise that an individual can have both good and bad qualities. In paranoid-schizoid position, the anxiety is about the survival of the self but in depressive position the anxiety is about the loss of the object.

The correct answer is: Ambivalence

Question **43**

Not answered

Marked out of 1

Which one of the following is not a technique used in DBT?

Select one:

- ☐ Distress tolerance training
- ☐ Interpersonal effectiveness training
- ☐ Role transition
- ☐ Core mindfulness training
- ☐ Emotion regulation skills training

Your answer is incorrect.

The four modes of treatment in DBT are (1) group skills training, (2) individual therapy, (3) phone consultations, and (4) consultation team. Role transitions are managed in interpersonal therapy.

The correct answer is: Role transition



Question **44**

Not answered

Marked out of 1

Which one among the following statements about dream work is true?

Select one:

- ☐ Dreams are always interpreted in psychoanalytic psychotherapy
- ☐ The dream work involves conversion of latent content into manifest content.
- ☐ Dream work converts the manifest content of the dream into its latent content
- ☐ Primary process thinking does not occur in dreaming
- ☐ The mental mechanisms involved in the psychoanalytic therapy of dreams include primary elaboration

Your answer is incorrect.

According to Freud, dreams are the royal road to the unconscious. In dreams, Freud distinguished between latent content (unconscious feelings that include the wishes, anxieties and fantasies) and manifest content (recalled dream). The dream work involves conversion of latent content into manifest content.

The correct answer is: The dream work involves conversion of latent content into manifest content.



Question **45**

Not answered

Marked out of 1

Cognitive analytical therapy was developed by which one of the following therapists?

Select one:

- ☐ Carl Rogers
- ☐ Anthony Ryle
- ☐ Elspeth Guthrie
- ☐ David Kingdon
- ☐ Robert Ellis

Your answer is incorrect.

Cognitive-analytic therapy (CAT) is a brief focal therapy informed by cognitive therapy, psycho- dynamic psychotherapy and certain developments in cognitive psychology. Anthony Ryle developed CAT specifically in response to the need for treatment approaches of short duration in the National Health Service (NHS).

The correct answer is: Anthony Ryle



Question 46

Not answered

Marked out of 1

Which one of the following is least used technique in CBT

Select one:

- ☐ Paradoxical Injunction
- ☐ Empirical reasoning
- ☐ Collaborative empiricism
- ☐ Socratic questioning
- ☐ Homework tasks

Your answer is incorrect.

The paradoxical injunction is a term used in family therapy. The therapist makes the patient intentionally engage in the unwanted behaviour (called the paradoxical injunction), e.g., avoid a phobic object or perform a compulsive ritual. This counterintuitive approach can provide new insight for some patients.

The correct answer is: Paradoxical Injunction

Question **47**

Not answered

Marked out of 1

An 18-year-old girl with relapse of bulimia is found to have poor engagement with the treating team. The most suitable psychological therapy would be

Select one:

- ☐ Flooding
- ☐ Dialectic behavioural therapy
- ☐ Exposure therapy
- ☐ Interpersonal therapy
- ☐ Implosion therapy

Your answer is incorrect.

Interpersonal therapy (IPT) has been identified as an effective treatment for bulimia nervosa that does not focus on bulimic symptoms. Rather, a detailed assessment culminating in an 'interpersonal inventory' identifies core associated interpersonal problem(s) that become the focus of treatment.

The correct answer is: Interpersonal therapy



Question **48**

Not answered

Marked out of 1

Choose the first step in treating a patient who is afraid of spiders and avoids them:

Select one:

- ☐ taking detailed history
- ☐ Flooding with spiders
- ☐ Graded exposure immediately
- ☐ explaining CBT
- ☐ Paroxetine treatment

Your answer is incorrect.

Taking a detailed history to establish the diagnosis is the most important step.

The correct answer is: taking detailed history



Question **49**

Not answered

Marked out of 1

Choose the most accurate statement about Interpersonal Therapy?

Select one:

- ☐ Focus is on social skills development
- ☐ Attempts to explain psychoanalytic ideas in cognitive terms
- ☐ Aim is to extinguish maladaptive behaviour and substitute more adaptive ones
- ☐ Empower patients to make use of their own strengths
- ☐ Therapy focuses on the 'here and now'

Your answer is incorrect.

Interpersonal therapy is a practical short-term, time-limited, 'here and now' therapy. Emotional problems are best understood by studying the interpersonal context (ongoing/current situation) in which they arise.

The correct answer is: Therapy focuses on the 'here and now'



Question **50**

Not answered

Marked out of 1

Which one of the following is not a technique used in psychodynamic psychotherapy?

Select one:

- ☐ Free association
- ☐ Exploration of transference/counter-transference
- ☐ Examination of parapraxes
- ☐ Examination of dreams
- ☐ Reinforcement

Your answer is incorrect.

Reinforcement (positive or negative) is a technique used in behavioural psychotherapy - not psychodynamic psychotherapy.

The correct answer is: Reinforcement



Question **51**

Not answered

Marked out of 1

Which of the following is a relative contraindication to brief time-limited psychotherapeutic approach?

Select one:

- ☐ Having simple and specific problems to be addressed
- ☐ Chronic maladaptive strategies
- ☐ Psychological mindedness
- ☐ Readiness for change
- ☐ Good ability to form trusting relationships

Your answer is incorrect.

Chronic maladaptive patterns tend to be deeply wired and need longer therapies.

The correct answer is: Chronic maladaptive strategies



Question **52**

Not answered

Marked out of 1

Which one of the following, if present, reduces the suitability of a patient for psychoanalytic psychotherapy?

Select one:

- ☐ Antisocial personality disorder
- ☐ High motivation
- ☐ Psychological mindedness
- ☐ Good ego strength
- ☐ Ability to form and sustain relationships

Your answer is incorrect.

Poor frustration tolerance and impulsivity seen in antisocial PD are among the factors indicating poor suitability for psychoanalytic modes of therapy.

The correct answer is: Antisocial personality disorder



Question **53**

Not answered

Marked out of 1

What predicts a poor response to psychodynamic psychotherapy?

Select one:

- ☐ Good impulse control and good frustration tolerance
- ☐ Patient can understand things in psychological terms
- ☐ Intense deliberate self harm
- ☐ Good perception of treatment alliance.
- ☐ Moderate depressive illness

Your answer is incorrect.

Repeated uncontrolled self-harm can interfere significantly with the psychotherapy. In general, such patients are not suitable for a psychodynamic approach.

The correct answer is: Intense deliberate self harm



Question **54**

Not answered

Marked out of 1

Which one of the following is a cognitive technique used in CBT?

Select one:

- ☐ Graded task assignments
- ☐ Relaxation training
- ☐ Socratic questioning
- ☐ Activity scheduling
- ☐ Hypothesis testing

Your answer is incorrect.

The overall purpose of Socratic questioning is to challenge accuracy and completeness of thinking in a way that acts to move people towards their ultimate goal.

The correct answer is: Socratic questioning



Question **55**

Not answered

Marked out of 1

Interpersonal therapy does not focus on one of the following areas.

Select one:

- ☐ Interpersonal deficits
- ☐ Role transitions
- ☐ Attachment pattern
- ☐ Grief
- ☐ Interpersonal role disputes

Your answer is incorrect.

Examples of areas covered are disputes with friends, family or co-workers, grief and loss and role transitions, such as retirement or divorce. Attachment issues are not dealt in IPT.

The correct answer is: Attachment pattern



Question **56**

Not answered

Marked out of 1

Unconditional positive regard is a feature of

Select one:

- ☐ Cognitive behavioural therapy
- ☐ Client centred psychotherapy
- ☐ Leader centred psychotherapy
- ☐ Group therapy
- ☐ Family therapy

Your answer is incorrect.

Client-centred therapy by Carl Rogers emphasises the importance of therapeutic attributes of genuineness, unconditional positive regard and accurate empathy.

The correct answer is: Client centred psychotherapy



Question **57**

Not answered

Marked out of 1

Which of the following therapies use 'mindfulness' as a therapeutic strategy?

Select one:

- ☐ Systems therapy
- ☐ Group therapy (Milan)
- ☐ Dialectical behaviour therapy
- ☐ Cognitive analytical therapy
- ☐ Rational emotive therapy

Your answer is incorrect.

Core mindfulness training i.e. learning to monitor internal mental states, is a characteristic feature of DBT.

The correct answer is: Dialectical behaviour therapy



Question **58**

Not answered

Marked out of 1

Which one of the following is a self-rating scale?

Select one:

- ☐ Montgomery Asberg depression rating scale
- ☐ Young mania scale
- ☐ Beck depression inventory
- ☐ Hamilton rating scale for depression
- ☐ Yale-brown obsessive compulsive scale

Your answer is incorrect.

Self-rating scale-Beck depression inventory, hospital anxiety and depression scale and Zung depression rating scale. Observer rating scales-Hamilton rating scale for depression, Montgomery Asberg depression rating scale, Young Mania Scale and Yale-brown obsessive compulsive scale.

The correct answer is: Beck depression inventory

Question 59

Not answered

Marked out of 1

Which of the following predicts poor treatment response in chronic fatigue syndrome when treated with CBT/graded exercise?

Select one:

- ☐ Continuing with occupation
- ☐ High sense of physical control
- ☐ Being passive with reduced activity
- ☐ Long duration of illness
- ☐ Less emphasis on physical symptoms

Your answer is incorrect.

Predictors of poor outcome when treating CFS include (Powell et al, 2004) 1. claiming a disability related benefit 2. low sense of control 3. strong focus on physical symptoms 4. being passive with reduced activity and 5. membership of self help group (Note: this does not mean that self help groups worsen CFS; membership may be an indicator of perceived severity)

The correct answer is: Being passive with reduced activity



Question **60**

Not answered

Marked out of 1

Defence mechanisms are a function of the

Select one:

- ☐ Id
- ☐ Superego
- ☐ Both Id and Ego
- ☐ Ego
- ☐ Both Ego and superego

Your answer is incorrect.

The first systematic account of defence mechanisms was written by Freud's daughter Anna Freud in the book 'Ego and the Mechanisms of Defence' in 1936.

The correct answer is: Ego



© 2023 SPMM Course Limited, UK.

[Refund Policy](#) | [Terms & Conditions](#)

