

---

**Started on** Sunday, 12 February 2023, 8:27 PM

**State** Finished

**Completed on** Sunday, 12 February 2023, 8:27 PM

**Time taken** 12 secs

**Marks** 0/69

**Grade** 0 out of 100

Question **1**

Not answered

Marked out of 1

Which of the following is a relative contraindication to brief time-limited psychotherapeutic approach?

Select one:

- ☐ Having simple and specific problems to be addressed
- ☐ Chronic maladaptive strategies
- ☐ Psychological mindedness
- ☐ Readiness for change
- ☐ Good ability to form trusting relationships

Your answer is incorrect.

Chronic maladaptive patterns tend to be deeply wired and need longer therapies.

The correct answer is: Chronic maladaptive strategies

## Question 2

Not answered

Marked out of 1

Which one of the following, if present, reduces the suitability of a patient for psychoanalytic psychotherapy?

Select one:

- ☐ Antisocial personality disorder
- ☐ High motivation
- ☐ Psychological mindedness
- ☐ Good ego strength
- ☐ Ability to form and sustain relationships

Your answer is incorrect.

Poor frustration tolerance and impulsivity seen in antisocial PD are among the factors indicating poor suitability for psychoanalytic modes of therapy.

The correct answer is: Antisocial personality disorder



## Question 3

Not answered

Marked out of 1

What predicts a poor response to psychodynamic psychotherapy?

Select one:

- ☐ Good impulse control and good frustration tolerance
- ☐ Patient can understand things in psychological terms
- ☐ Intense deliberate self harm
- ☐ Good perception of treatment alliance.
- ☐ Moderate depressive illness

Your answer is incorrect.

Repeated uncontrolled self-harm can interfere significantly with the psychotherapy. In general, such patients are not suitable for a psychodynamic approach.

The correct answer is: Intense deliberate self harm



## Question 4

Not answered

Marked out of 1

Which one of the following is a cognitive technique used in CBT?

Select one:

- ☐ Graded task assignments
- ☐ Relaxation training
- ☐ Socratic questioning
- ☐ Activity scheduling
- ☐ Hypothesis testing

Your answer is incorrect.

The overall purpose of Socratic questioning is to challenge accuracy and completeness of thinking in a way that acts to move people towards their ultimate goal.

The correct answer is: Socratic questioning



## Question 5

Not answered

Marked out of 1

Interpersonal therapy does not focus on one of the following areas.

Select one:

- ☐ Interpersonal deficits
- ☐ Role transitions
- ☐ Attachment pattern
- ☐ Grief
- ☐ Interpersonal role disputes

Your answer is incorrect.

Examples of areas covered are disputes with friends, family or co-workers, grief and loss and role transitions, such as retirement or divorce. Attachment issues are not dealt in IPT.

The correct answer is: Attachment pattern



## Question 6

Not answered

Marked out of 1

Unconditional positive regard is a feature of

Select one:

- ☐ Cognitive behavioural therapy
- ☐ Client centred psychotherapy
- ☐ Leader centred psychotherapy
- ☐ Group therapy
- ☐ Family therapy

Your answer is incorrect.

Client-centred therapy by Carl Rogers emphasises the importance of therapeutic attributes of genuineness, unconditional positive regard and accurate empathy.

The correct answer is: Client centred psychotherapy



## Question 7

Not answered

Marked out of 1

Which of the following therapies use 'mindfulness' as a therapeutic strategy?

Select one:

- ☐ Systems therapy
- ☐ Group therapy (Milan)
- ☐ Dialectical behaviour therapy
- ☐ Cognitive analytical therapy
- ☐ Rational emotive therapy

Your answer is incorrect.

Core mindfulness training i.e. learning to monitor internal mental states, is a characteristic feature of DBT.

The correct answer is: Dialectical behaviour therapy



## Question 8

Not answered

Marked out of 1

Which one of the following is a self-rating scale?

Select one:

- ☐ Montgomery Asberg depression rating scale
- ☐ Young mania scale
- ☐ Beck depression inventory
- ☐ Hamilton rating scale for depression
- ☐ Yale-brown obsessive compulsive scale

Your answer is incorrect.

Self-rating scale-Beck depression inventory, hospital anxiety and depression scale and Zung depression rating scale. Observer rating scales-Hamilton rating scale for depression, Montgomery Asberg depression rating scale, Young Mania Scale and Yale-brown obsessive compulsive scale.

The correct answer is: Beck depression inventory



## Question 9

Not answered

Marked out of 1

Which of the following predicts poor treatment response in chronic fatigue syndrome when treated with CBT/graded exercise?

Select one:

- ☐ Continuing with occupation
- ☐ High sense of physical control
- ☐ Being passive with reduced activity
- ☐ Long duration of illness
- ☐ Less emphasis on physical symptoms

Your answer is incorrect.

Predictors of poor outcome when treating CFS include (Powell et al, 2004) 1. claiming a disability related benefit 2. low sense of control 3. strong focus on physical symptoms 4. being passive with reduced activity and 5. membership of self help group (Note: this does not mean that self help groups worsen CFS; membership may be an indicator of perceived severity)

The correct answer is: Being passive with reduced activity

## Question 10

Not answered

Marked out of 1

Defence mechanisms are a function of the

Select one:

- ☐ Id
- ☐ Superego
- ☐ Both Id and Ego
- ☐ Ego
- ☐ Both Ego and superego

Your answer is incorrect.

The first systematic account of defence mechanisms was written by Freud's daughter Anna Freud in the book 'Ego and the Mechanisms of Defence' in 1936.

The correct answer is: Ego



## Question 11

Not answered

Marked out of 1

Which one of the following is not a goal of cognitive therapy?

Select one:

- ☐ Revise cognitive assumptions
- ☐ Test the validity of cognitions
- ☐ Identify learning styles
- ☐ Develop alternative behaviours
- ☐ Identify negative cognitions

Your answer is incorrect.

The goals of cognitive therapy are to identify negative cognitions, test the validity of those cognitions and revise cognitive assumptions.

The correct answer is: Identify learning styles



## Question 12

Not answered

Marked out of 1

Which one of the following is a behavioural technique used in CBT?

Select one:

- ☐ Socratic questioning
- ☐ Thoughts diary
- ☐ Graded assignments
- ☐ Psycho education
- ☐ Identifying automatic thoughts

Your answer is incorrect.

The therapist is very active in CBT and assists the patient to monitor cognitions, identify cognitive errors, understand maladaptive schema and explore with them strategies to challenge and change these and examine the resultant symptomatic effects. The cognitive techniques include psychoeducation, including reading assignments, identifying automatic thoughts, Socratic questioning, role play, thoughts diary, thought rehearsal, examining the evidence and working through the options. The behavioural techniques include activity scheduling, graded assignments, exposure, response prevention, distraction, relaxation training and assertiveness and social skills training.

The correct answer is: Graded assignments

## Question 13

Not answered

Marked out of 1

Leader assertiveness is one of the therapeutic factors described in

Select one:

- ☐ Group psychotherapy
- ☐ Behaviour therapy
- ☐ Family therapy
- ☐ Individual psychotherapy
- ☐ Interpersonal therapy

Your answer is incorrect.

The therapeutic factors described by Yalom in Group psychotherapy include altruism, catharsis, corrective recapitulation of the group, development of socialising techniques, existential awareness, group cohesiveness, instillation of hope, imparting of information, interpersonal learning, imitative behaviour and universality.

The correct answer is: Group psychotherapy



## Question 14

Not answered

Marked out of 1

Which of the following has a healing effect in a group setting?

Select one:

- ☐ initiation
- ☐ pairing
- ☐ debriefing
- ☐ cohesion
- ☐ triangulation

Your answer is incorrect.

Of all the described therapeutic factors (TFs), the mechanism of cohesion is the most central that facilitates the action of other TFs. Cohesion is the best definition of the therapeutic relationship in group (Retrieved from <http://www.agpa.org/home/practice-resources/practice-guidelines-for-group-psychotherapy>).

The correct answer is: cohesion

## Question 15

Not answered

Marked out of 1

Which one of the following is NOT a curative factor in group therapies?

Select one:

- ☐ Universality
- ☐ Pairing
- ☐ Instillation of hope
- ☐ Altruism
- ☐ Catharsis

Your answer is incorrect.

Yalom cited 11 'curative' factors responsible for change in groups. The curative factors include instillation of hope, universality, imparting information (feedback), altruism, corrective recapitulation, socialisation techniques, imitative behaviour, interpersonal learning, group cohesiveness, catharsis and existential factors. Of these, cohesiveness, and learning from feedback are valued positively though other factors may also be important.

The correct answer is: Pairing

## Question 16

Not answered

Marked out of 1

Which of the following is a component of the Dialectic Behavioural Therapy proposed by Linehan?

Select one:

- ☐ Dilemma management
- ☐ Fair treatment
- ☐ Paradoxical Injunction
- ☐ Mindfulness
- ☐ Interpersonal dependency

Your answer is incorrect.

The core skill areas in DBT include mindfulness, distress tolerance, emotion regulation, and interpersonal effectiveness.

The correct answer is: Mindfulness





## Question 17

Not answered

Marked out of 1

Which of the following is least likely to be correct concerning Camberwell Family Interview?

Select one:

- ☐ Audio-tapes are used
- ☐ Ratings based on interviewing parents singly had the most predictive value.
- ☐ It is a long interview
- ☐ It is used to measure EE levels
- ☐ The patient is not interviewed

Your answer is incorrect.

CFI is a long interview process where individual members of a family are interviewed (including the patient). If one relative is classified as high EE relative, the family is termed a high EE family. It was found that ratings based on interviewing parents individually had the most predictive value.

The correct answer is: The patient is not interviewed

## Question 18

Not answered

Marked out of 1

Rolling with resistance is a technique used as a major component in which of the following psychotherapies?

Select one:

- ☐ Jungian psychoanalysis
- ☐ Motivational interviewing
- ☐ Cognitive analytical therapy
- ☐ Object relation psychoanalysis
- ☐ Dialectical behavioural therapy

Your answer is incorrect.

Resistance is a predictive of poor treatment outcomes and lack of involvement in the therapeutic process. Rolling with resistance is a technique used in motivational interviewing.

The correct answer is: Motivational interviewing



## Question 19

Not answered

Marked out of 1

Which one among the following is not considered to be a Neo-Freudian?

Select one:

- ☐ Jean Piaget
- ☐ Erich Fromm
- ☐ Harry Stack Sullivan.
- ☐ Alfred Adler
- ☐ Karen Horney

Your answer is incorrect.

Neo-Freudians are thinkers who agree with the basis of Freud's psychoanalytic theory, but change or adapt it to incorporate their own beliefs, ideas and theories. Piaget is not considered as a Freudian theorist.

The correct answer is: Jean Piaget



## Question 20

Not answered

Marked out of 1

A depressed schoolteacher thinks 'one student in the class is sleeping, so the entire class is bored of my lecture'. The cognitive distortion in this scenario is

Select one:

- ☐ Personalisation
- ☐ Dichotomous thinking
- ☐ Magnification
- ☐ Selective abstraction
- ☐ Overgeneralisation

Your answer is incorrect.

This is one of the many cognitive distortions described by Beck.

The correct answer is: Overgeneralisation



Question **21**

Not answered

Marked out of 1

Interpersonal Psychotherapy is associated with the following except

Select one:

- ☐ Runs from 12 to 16 hourly sessions, usually occurring weekly
- ☐ Making a relationship inventory
- ☐ Focuses on snags and dilemmas
- ☐ Developed by Klerman and Weissman
- ☐ Establishing the sick role

Your answer is incorrect.

Cognitive analytical therapy focuses on snags and dilemmas. IPT focuses on grief and role transitions.

<http://pb.rcpsych.org/cgi/content/full/26/1/26>

The correct answer is: Focuses on snags and dilemmas



## Question 22

Not answered

Marked out of 1

which one among the following is not correctly paired?

Select one:

- ☐ Klerrmann and Weissmann-Interpersonal therapy
- ☐ Malan and Davanloo - Brief dynamic therapy
- ☐ Albert Ellis - Rational Emotive therapy
- ☐ Milan - Strategic family therapy
- ☐ Winnicott - Object Relations Theory

Your answer is incorrect.

Palazolli and colleagues from Milan are associated with systemic family therapy - not structural therapy. Strategic family therapy is associated with Haley. Haley called the therapy Strategic because "it is a therapy where the therapist initiates what happens during therapy, designs a specific approach for each person's presenting problem, and takes responsibility for directly influencing people" (Retrieved from [http://www.mri.org/strategic\\_family\\_therapy.html](http://www.mri.org/strategic_family_therapy.html)).

The correct answer is: Milan - Strategic family therapy

Question **23**

Not answered

Marked out of 1

Which of the following behavioural treatment is useful in trichotillomania?

Select one:

- ☐ Imaginal flooding
- ☐ Exposure and response prevention
- ☐ Massed negative practice
- ☐ Habit reversal
- ☐ Biofeedback

Your answer is incorrect.

Habit reversal training (HRT) is a behavioral treatment package useful in tics and other OCD spectrum disorders such as trichotillomania, nail biting, thumb sucking, and skin picking etc.

The correct answer is: Habit reversal



Question **24**

Not answered

Marked out of 1

Psychodynamic therapy can be used for a wide variety of disorders. But it is contraindicated for which of the following?

Select one:

- ☐ Personality disorders.
- ☐ Mild to moderate depressive illness
- ☐ Acute psychosis
- ☐ Neurotic disorders
- ☐ Behavioural disorders

Your answer is incorrect.

Psychodynamic psychotherapy is generally avoided in acute psychosis, very high risk of recurrent DSH and conditions where there is acute suicidal risk, delirium, dementia, very severe depressive illness and in drug and alcohol misuse. Indicators that a patient is likely to respond to psychotherapy include a good response to a trial interpretation, a tendency for self-reflection, presence of realistic expectations, a reasonable degree of personality integration and a motivation for change.

The correct answer is: Acute psychosis



## Question 25

Not answered

Marked out of 1

Which one of the following techniques is considered to be anti-therapeutic in group therapy?

Select one:

- ☐ Catharsis
- ☐ Interpretation
- ☐ Idealization of the therapist.
- ☐ Mirroring
- ☐ Amplification

Your answer is incorrect.

Scape goating, idealization of the therapist and coupling are considered to be anti-therapeutic in group therapy. Scapegoating is an explanation of discrimination based on the idea of frustration-aggression. The work of Bion, Ezriel and Foulkes were crucial in the development of group therapy.

The correct answer is: Idealization of the therapist.

## Question 26

Not answered

Marked out of 1

The basis of object relations theory is that the primary motivational drive of an individual is to seek

Select one:

- ☐ Satisfaction
- ☐ Money
- ☐ Self actualisation
- ☐ Pleasure
- ☐ Relationships

Your answer is incorrect.

Object relations theory describes with how a growing baby develops relationships with others. Gratification is thought to be obtained through relationships as well as through satisfaction of desires.

The correct answer is: Relationships



Question **27**

Not answered

Marked out of 1

A 25-year-old bride is getting ready for her wedding. She thinks that if she is late for her wedding, then her bridegroom will refuse to marry her, and she will be left on her own forever. Choose one cognitive distortion that is most likely

Select one:

- ☐ Rationalisation
- ☐ Projection
- ☐ Over-generalisation
- ☐ Catastrophising
- ☐ Splitting

Your answer is incorrect.

Catastrophisation is a cognitive distortion that refers to the tendency of always predicting the worst case scenario to happen. E.g. A woman with panic attacks can sense mild palpitation after some house work, and thinks she is going to have a massive heart attack and immediate death.

The correct answer is: Catastrophising



## Question 28

Not answered

Marked out of 1

Which one of the following is not included in Bion's basic assumptions?

Select one:

- ☐ Intellectualization
- ☐ Pairing
- ☐ Dependency
- ☐ Flight
- ☐ Fight

Your answer is incorrect.

According to Bion, whenever a group gets derailed from its task, it deteriorates into one of the three basic states: dependency, pairing, or fight-flight. A 4th basic assumption was introduced by Hopper called massification/aggregation where a rigid fusion of identities leads to loss of individuality, or extensive withdrawal leads to loss of mutual dependence.

The correct answer is: Intellectualization

## Question 29

Not answered

Marked out of 1

Which one of the following often seen in the process of supportive psychotherapy?

Select one:

- ☐ Good therapeutic alliance
- ☐ Good psychological mindedness
- ☐ Patient undergoing a crisis
- ☐ Patient seeking exploration of cause of symptoms
- ☐ Therapist having set boundaries

Your answer is incorrect.

Supportive therapy aims to offer practical and emotional support, provides an opportunity for ventilation of emotions and guided problem solving discussions. The primary aim of supportive psychotherapy is to support reality testing, provide ego support and attempt to reestablish usual level of functioning. It is usually employed in otherwise healthy patients with overwhelming ongoing crises and those with ego deficits.

The correct answer is: Patient undergoing a crisis



Question **30**

Not answered

Marked out of 1

What predicts a good response to dynamic psychotherapy?

Select one:

- ☐ Positive perception of the therapist throughout treatment
- ☐ Having had previous therapy
- ☐ Motivation developed after sustained effort from therapist
- ☐ Highly intelligent patient.
- ☐ Patient can understand issues in psychological terms

Your answer is incorrect.

A curiosity to understand and reformulate problems in terms of one's life experiences or psychological structure is an important predictor of response to psychotherapy.

The correct answer is: Patient can understand issues in psychological terms



Question **31**

Not answered

Marked out of 1

Genetic counselling is classified under which type of counselling?

Select one:

- ☐ Psychoeducation
- ☐ Risk counselling
- ☐ Information sharing
- ☐ Crisis management
- ☐ Problem based counselling

Your answer is incorrect.

Counselling may be thought of as a method of relieving distress undertaken by means of a dialogue between two people. The aim is to help the client or patient find their own solutions to problems, while being supported to do so and being guided by appropriate advice. The different types include psycho education, crisis management, problem-based counselling directed towards a specific primary problem, e.g. drug misuse and risk counselling used to guide an informed decision (e.g. pre-natal interventions, genetic counselling). The latter is differentiated from other forms of counselling by the fact that the counsellor is clearly the expert and has access to specialist information.

The correct answer is: Risk counselling



Question **32**

Not answered

Marked out of 1

Idealization and denigration are products of which of the following Kleinian processes?

Select one:

- ☐ Regression
- ☐ Repression
- ☐ Projective identification
- ☐ Projection
- ☐ Splitting

Your answer is incorrect.

Splitting refers to splitting of self into good and bad. Idealization and devaluation (denigration) are the results of splitting and are central in the Kleinian school of thought.

The correct answer is: Splitting





## Question 33

Not answered

Marked out of 1

Which one of the following group therapies has a high level of leader input?

Select one:

- ☐ Problem solving therapy
- ☐ Genetic counselling
- ☐ Psycho educational group
- ☐ Structured groups for drug or alcohol use
- ☐ Tavistock model analytic group

Your answer is incorrect.

Group therapies can be classified according to the objectives of the group and how the group is led or managed (leadership). Highly specific target oriented groups include structured groups for drug use or alcohol use, activity groups like occupational therapy, groups etc. These groups have high level of leader input. Psychodrama, music therapy, systems centred groups are some less specific therapies but are highly directed by the leader or therapist. Problem solving therapy and psycho educational groups are highly specific but have low level of therapist activity. Support groups, art therapy, interpersonal therapy and groups like Tavistock model analytic groups have low level of leader activity and have low specificity with respect to treatment goals

The correct answer is: Structured groups for drug or alcohol use

Question **34**

Not answered

Marked out of 1

The definite termination date is set at beginning of treatment in which of the following therapies?

Select one:

- ☐ Psychoanalytical psychotherapy
- ☐ Dialectical behavioural therapy
- ☐ Family therapy
- ☐ Brief dynamic therapy
- ☐ Cognitive analytical therapy

Your answer is incorrect.

Brief psychodynamic psychotherapy is a time-limited treatment that is based on psychoanalysis and psychodynamic theory. Brief psychodynamic therapy is an intervention where the concepts of symptom development and methods of therapy are based on those of psychoanalysis and are more focused on the here and now- the patient's current experience of the world, and techniques are employed to accelerate the process of therapy but where the time scale and number of sessions are reduced.

The correct answer is: Brief dynamic therapy



## Question 35

Not answered

Marked out of 1

Which of the following therapies uses techniques such as circular questioning and hypothesising?

Select one:

- ☐ Psychodynamic psychotherapy
- ☐ Systemic family therapy
- ☐ Interpersonal therapy
- ☐ Cognitive analytical therapy
- ☐ Cognitive behavioural therapy

Your answer is incorrect.

The Milan systemic approach (Selvini Palazzoli et al, 1978) has been modified over the years, from its original paradoxical prescriptions to a great emphasis on a particular style of interviewing – circular and reflexive questioning (Selvini Palazzoli et al, 1980). This technique focuses on questioning the various family members' beliefs and perceptions regarding relationships. Asking each to comment and reflect on the answers given by the various family members creates feedback that changes the fabric of family interactions (Retrieved from <http://apt.rcpsych.org/content/aptrcpsych/8/3/230.full.pdf>).

The correct answer is: Systemic family therapy

Question **36**

Not answered

Marked out of 1

Thought stopping is a behavioural technique used in

Select one:

- ☐ Borderline Personality Disorder
- ☐ Phobias
- ☐ OCD
- ☐ Somatisation disorder
- ☐ PTSD

Your answer is incorrect.

Additional behaviour and cognitive-behavioral techniques which may have some effectiveness for people who suffer from obsessive compulsive disorder include saturation and thought-stopping. Saturation is a process where the patient is directed to think repeatedly of the most intrusive obsessional thought. After an extended period of concentration on this one thought (e.g., 10-15 minutes at a time) over a number of days (3-5 days), the obsession can lose some of its strength. Thought-stopping is a technique where the patient learns how to halt obsessive thoughts through identification of the obsessional thoughts, and then averting them by carrying out a strong, incompatible response.

The correct answer is: OCD

Question **37**

Not answered

Marked out of 1

An effective technique for treating hypochondriasis is

Select one:

- ☐ Reassurance
- ☐ Psychodynamic psychotherapy
- ☐ Supportive psychotherapy
- ☐ Anxiety management
- ☐ Behavioural experiments

Your answer is incorrect.

In the treatment of hypochondriasis, it is important to break the cycle of reassurance and repeat presentation. Behavioural experiments focused on response prevention and exposure to illness cues is the most effective strategy in the management. This could be done as a part of CBT (The role of CBT is to identify and challenge misinterpretations, substitution of realistic interpretation, graded exposure to illness related situations and modification of core beliefs).

The correct answer is: Behavioural experiments

## Question 38

Not answered

Marked out of 1

Which of the following best describes the stance that is held by the therapist while administering Interpersonal Psychotherapy?

Select one:

- ☐ Befriending
- ☐ Antagonistic stance
- ☐ Patient's advocate
- ☐ Reflective stance
- ☐ Passive stance

Your answer is incorrect.

In IPT, the therapist is an explicit ally and advocate of the patient. The therapist is nonjudgmental, expresses warmth and positive regard for the patient and congratulates the patient as progress in the problem areas is made. The therapist works with the patient and for the patient and believes that the patient's problems can be solved. This does not imply that the therapist accepts all aspects of the patient, as that would preclude any stimulus for change. The therapist always tries to have the patient find the solution for the problems discussed in the session. However, the therapist is not afraid to make suggestions or provide direct advice when they seem useful.

The correct answer is: Patient's advocate

## Question 39

Not answered

Marked out of 1

Which one among the following is not one of the therapeutic techniques used in Interpersonal therapy?

Select one:

- ☐ Relationship problem solving
- ☐ Role-play
- ☐ Free association
- ☐ Facilitation of grief
- ☐ Catharsis

Your answer is incorrect.

The therapeutic techniques in IPT are specific to the problem under review and includes relationship problem-solving, facilitation of grief, catharsis and role-play. The role disputes can be resolved by renegotiation, acceptance of impasse or dissolution. Free association is a Freudian concept not relevant for IPT.

The correct answer is: Free association

## Question 40

Not answered

Marked out of 1

Systematic desensitization involves all of the following steps except

Select one:

- ☐ Desensitisation of the stimulus.
- ☐ Relaxation training
- ☐ Graded exposure
- ☐ Hierarchy construction
- ☐ Assertiveness training

Your answer is incorrect.

Systematic Desensitisation (Wolpe) is used on the behavioral principle of counter conditioning (i.e. gradual approach of feared situation in a psychophysiological state that inhibits anxiety leads to reduction of anxiety response.) Systematic desensitization consists of three steps: relaxation training, hierarchy construction, and desensitization of the stimulus. The patient is exposed to a graded hierarchy of anxiety-provoking situations in step wise fashion.

The correct answer is: Assertiveness training



Question **41**

Not answered

Marked out of 1

Which one of the following is considered as a mature defence mechanism?

Select one:

- ☐ Sublimation
- ☐ Regression
- ☐ Projection
- ☐ Repression
- ☐ Denial

Your answer is incorrect.

Primitive/immature defenses include splitting, projection, projective identification, idealization, denigration, introjection, passive aggression and turning against the self. Neurotic defences include repression, regression, denial, displacement, dissociation, isolation, reaction formation, intellectualisation, rationalization, externalisation and identification with the aggressor. Humour, anticipation, sublimation, altruism and suppression (SASHA) are all mature defence mechanisms.

The correct answer is: Sublimation

Question **42**

Not answered

Marked out of 1

Which of the following is a technique used in motivational interviewing?

Select one:

- ☐ Risk counselling
- ☐ Advice and education
- ☐ Reciprocal roles
- ☐ Rolling with resistance
- ☐ Role transition

Your answer is incorrect.

When using motivational interviewing, one does not fight client resistance, but instead 'rolls with it.' Resistance can take several forms, such as negating, blaming, etc.

The correct answer is: Rolling with resistance



## Question 43

Not answered

Marked out of 1

Which one among the following is NOT a part of the Beck's theory of depression?

Select one:

- ☐ Magnification
- ☐ Selective abstraction
- ☐ Arbitrary inference
- ☐ All or nothing thinking
- ☐ Turning against self

Your answer is incorrect.

Beck's cognitive model of depression includes two levels of cognitive errors that include automatic thoughts and core schemas. The cognitive errors include selective abstraction, arbitrary inference, minimisation, magnification, all or nothing thinking, personalisation, catastrophic thinking and overgeneralization. Turning against self is a psychodynamic concept of depression.

The correct answer is: Turning against self

Question **44**

Not answered

Marked out of 1

Who described the unconscious as composed of both the collective unconscious and personal unconscious?

Select one:

- ☐ Carl Gustav Jung
- ☐ Sigmund Freud
- ☐ Anna Freud
- ☐ Winnicott
- ☐ Alfred Adler

Your answer is incorrect.

Carl Gustav Jung described a holistic theory of psyche that could be applied across varied cultures. According to him, the unconscious mind includes the collective and personal unconscious. He postulated that the personal unconscious is unique to each but the collective unconscious is universal, common to mankind, contains the archetypes that include anima, animus, shadow or self and Hero.

The correct answer is: Carl Gustav Jung

## Question 45

Not answered

Marked out of 1

Nursing staff at a learning disability care home are concerned about the self-injurious behaviour of a 40-year-old man. A psychologist is called for conducting a functional behavioral analysis. On which of the following information would she concentrate?

Select one:

- ☐ Staff attitudes towards the patient
- ☐ The immediate consequences of the patient's self injury at the care home setting
- ☐ Past history of parental child abuse
- ☐ Activity schedule of the patient
- ☐ Psychiatric symptomatology in the patient

Your answer is incorrect.

In behavioural analysis, one looks at antecedent, behaviour and its consequences.

The correct answer is: The immediate consequences of the patient's self injury at the care home setting



## Question 46

Not answered

Marked out of 1

Strategic family therapy is offered to a dysfunctional family. What technique is mainly used in this therapy?

Select one:

- ☐ Relapse sign identification
- ☐ Counter conditioning
- ☐ Unbalancing
- ☐ Task setting
- ☐ Story telling

Your answer is incorrect.

Task setting (homework tasks) are an essential component of strategic family therapy. The use of tasks in family therapy is consistent with the model's emphasis on having family members 'do' rather than simply 'talk about'. The tasks are relatively easier in the early part of treatment to foster a sense of success. Tasks tend to be more complex later in the treatment. In all instances, the therapists create opportunities for family members to practice and demonstrate competence in the task, prior to assigning homework.

The correct answer is: Task setting

Question **47**

Not answered

Marked out of 1

The closeness between two family members may exclude a third member. Study of these triangular patterns of relationships in family takes place in

Select one:

- ☐ Structural therapy
- ☐ Couples therapy
- ☐ Strategic family therapy
- ☐ Paradoxical therapy
- ☐ Family systems therapy

Your answer is incorrect.

Bowen's model (family systems approach) emphasizes one's ability to retain individual self in the face of familial tension. The degree of enmeshment is analysed here. An emotional triangle is a three-party system where closeness of two members (in either positive or negative sense) tends to exclude a third. This hot triangle leads to symptom formation. Here therapist maintains minimal emotional contact with family members.

The correct answer is: Family systems therapy

## Question 48

Not answered

Marked out of 1

Which school of family therapy are associated with the use of an ONE-way mirror?

Select one:

- ☐ Family psychoeducation
- ☐ Strategic systemic therapy
- ☐ Bowen's model
- ☐ Milan's systemic approach
- ☐ Structural model

Your answer is incorrect.

The Milan systemic approach (Palazolli) gives great emphasis on circular and reflexive questioning. In a circular fashion, each family member is asked to comment and reflect on each others response. The Milan systemic school of family therapy is associated with the use of an ONE-way mirror.

The correct answer is: Milan's systemic approach





## Question 49

Not answered

Marked out of 1

You are called to see a newly admitted patient with personality disorder in the ward; you notice that the nursing staff are angry and feel unwanted by the patient. What is the most likely psychological explanation?

Select one:

- ☐ Sublimation
- ☐ Counter-transference
- ☐ Dissociation
- ☐ Suppression
- ☐ Projection

Your answer is incorrect.

Projective identification (not just projection) has taken place here. The patient has projected her own 'hatred' on the therapists (nurses), and they are identifying themselves with the projected material and are acting in a way showing the hatred (identification). This is a mechanism explaining counter-transference.

The correct answer is: Counter-transference

## Question 50

Not answered

Marked out of 1

The 'repressive barrier' to censor unacceptable wishes and desires is maintained by

Select one:

- ☐ Unconscious system
- ☐ Id
- ☐ Preconscious system
- ☐ Super-ego
- ☐ The conscious system

Your answer is incorrect.

The preconscious interfaces with both unconscious and conscious - contents of unconscious become conscious by 'squeezing' through the preconscious. It maintains the 'repressive barrier' to censor unacceptable wishes and desires (not the repressed contents).

The correct answer is: Preconscious system



Question **51**

Not answered

Marked out of 1

Which type of group psychotherapy is used for the treatment of phobias?

Select one:

- ☐ Analytically oriented group therapy
- ☐ Supportive group therapy
- ☐ Transactional group therapy
- ☐ Psychoanalysis of groups
- ☐ Behavioural group therapy

Your answer is incorrect.

Behavioural groups can be organised to provide group therapy in a resource efficient manner for phobias.

The correct answer is: Behavioural group therapy



## Question 52

Not answered

Marked out of 1

The focus of attention is childhood experience and exploration of the unconscious mind in which of the following forms of therapy?

Select one:

- ☐ Hypnotherapy
- ☐ Psychodynamic therapy
- ☐ Supportive psychotherapy
- ☐ Counselling
- ☐ CBT

Your answer is incorrect.

Psychodynamic therapy upholds psychic determinism i.e. unconscious events play a causal role in later experiences. It emphasizes developmental psychopathology as the source of adult life difficulties.

The correct answer is: Psychodynamic therapy



Question **53**

Not answered

Marked out of 1

The first step in management of someone presenting with fear of spiders is

Select one:

- ☐ Detailed history for a functional assessment
- ☐ CBT
- ☐ Systematic desensitisation
- ☐ Pharmacotherapy
- ☐ Exposure and response prevention

Your answer is incorrect.

Before applying any behavioural treatment approach, a detailed history is required.

The correct answer is: Detailed history for a functional assessment



## Question 54

Not answered

Marked out of 1

Which type of family therapy encourages clear and open boundaries between the parents and the children?

Select one:

- ☐ Milan therapy
- ☐ Structural family therapy
- ☐ Psychodrama
- ☐ Systemic family therapy
- ☐ Strategic family therapy

Your answer is incorrect.

There are several types of family therapy based on the psychotherapeutic method applied. Structural family therapy developed by Minuchin emphasises that the structure of the family is maintained by unspoken rules and when the roles are broken the structure is weakened leading to problems in the family. Minuchin originally worked with families of psychosomatic and eating disordered patients. The Milan systemic school of family therapy is associated with the use of a ONE-way mirror. It is also associated with the concepts of paradoxes (vicious circle in which the family is trapped), circular questioning, neutrality and hypothesising. The strategic family therapy emphasises more active interventions and specific solutions for clearly defined family problem.

The correct answer is: Structural family therapy

## Question 55

Not answered

Marked out of 1

During on of your psychotherapy sessions, a patient reveals that he has been abusing his children, but suddenly he changes the topic. What should you do?

Select one:

- ☐ Keep your boundaries and end the session as planned.
- ☐ Terminate the interview immediately and inform the social services.
- ☐ Tell him you will inform the concerned authority
- ☐ Clarify what he means by abuse
- ☐ Ignore and continue with the flow the patients talk

Your answer is incorrect.

People might hold varying meanings for commonly used expressions. It is necessary to seek clarification before proceeding further.

The correct answer is: Clarify what he means by abuse



## Question 56

Not answered

Marked out of 1

Positive prognostic indicators for psychotherapy include all of the following except

Select one:

- ☐ tendency to self-scrutiny
- ☐ good response to a trial interpretation
- ☐ some response to medication
- ☐ a diagnosis of paraphrenia
- ☐ psychological mindedness

Your answer is incorrect.

In general psychotic disorders have been traditionally considered to involve ego disintegration and hence not suitable for therapeutic working through.

The correct answer is: a diagnosis of paraphrenia





Question **57**

Not answered

Marked out of 1

In psychotherapy, the term psychological mindedness refers to which of the following?

Select one:

- ☐ Ability to form a therapeutic alliance
- ☐ Motivation for change
- ☐ The ability to understand problems in physical terms
- ☐ The ability to understand problems in psychological terms
- ☐ Adequate ego strength

Your answer is incorrect.

The term psychological mindedness refers to an ability to understand problems in psychological terms. Ego strength is a term coined by Freud, and it includes the ability to sustain feelings and fantasies without impulsively acting upon them.

The correct answer is: The ability to understand problems in psychological terms



## Question 58

Not answered

Marked out of 1

The cognitive outcome you would hope to achieve when treating someone with hypochondriasis is

Select one:

- ☐ The complete absence of thoughts of illness.
- ☐ Ability to avoid medical help seeking when symptoms are troublesome
- ☐ Less distress at thoughts of illness
- ☐ Better role functioning
- ☐ Better ability to tolerate frustration

Your answer is incorrect.

CBT for hypochondriasis aims to generate a noncatastrophic alternative psychological explanation for a patient's problems. An RCT concluded that a six-session CBT intervention modestly but significantly improved a range of hypochondriacal symptoms, beliefs, and attitudes at six and 12 months. These results may not be generalisable as only 30 percent of eligible patients agreed for CBT trial. It is unclear which components of CBT really work in hypochondriasis. Hypochondriasis often is refractory to treatment and improvement involved better coping methods rather than cure. Role functioning may improve as a functional outcome. Reduced distress associated with beliefs rather than eradication of beliefs is the primary outcome expected (Barsky et al 2004).

The correct answer is: Less distress at thoughts of illness

## Question 59

Not answered

Marked out of 1

Which one of the following is not a component of CBT for delusions in psychosis?

Select one:

- ☐ Decreasing the impact of positive symptoms
- ☐ Therapeutic alliance
- ☐ Improving medication adherence.
- ☐ Flooded reality testing
- ☐ Providing alternate explanations to unusual experiences

Your answer is incorrect.

Though CBT is gaining popularity in treating psychotic symptoms, often, clinicians are unsure about the symptoms targeted in CBT. According to Birchwood, the target is the emotional dysfunction that accompanies psychotic experience and not the psychotic symptoms per se. Turkington described the following elements in CBT for psychosis: Therapeutic alliance, Improving medication adherence, Providing alternate explanations to unusual experiences e.g. normalisation, Decreasing the impact of positive symptoms e.g. addressing the omnipotence of voices and Graded reality testing using peripheral questioning and inference chaining.

The correct answer is: Flooded reality testing

## Question 60

Not answered

Marked out of 1

Which of the following is the best description of Bion's principle of containing?

Select one:

- ☐ Therapist manages to exclude distortion of one patient's story from another patient's experiences.
- ☐ Therapist maintains objectivity and focuses on facts
- ☐ Therapist doesn't let his feelings out and restrains himself.
- ☐ Therapist advises the patient to contain the feelings without letting them out.
- ☐ Patient has to explore the feelings within and relate them to the surroundings.

Your answer is incorrect.

The cognitive capacity of the therapist to maintain objectivity and focus on selected facts during a discourse is called 'containing' (Bion). The affective disposition of the therapist, which helps in restraining oneself from retaliating to negative transferences, is called 'holding.'

The correct answer is: Therapist maintains objectivity and focuses on facts



Question **61**

Not answered

Marked out of 1

Which of the following is a correct description of transference?

Select one:

- ☐ Good eye contact throughout therapy session
- ☐ Patient's response to the therapist based on previous relationships
- ☐ Therapist's response to the patient
- ☐ Confidence in therapist patient relationship
- ☐ Empathy in relationship

Your answer is incorrect.

Transference is an unconscious process that involves transferring a part of one's past relationship in an at least partly inappropriate fashion to the present life situation.

The correct answer is: Patient's response to the therapist based on previous relationships



## Question 62

Not answered

Marked out of 1

A psychotherapist feels that she is talking a lot to a patient and offering a lot of advice during therapy. This is because something in the patient's life is causing anxiety in her. This phenomenon is known as

Select one:

- ☐ Sublimation
- ☐ Therapeutic alliance
- ☐ Interpretation
- ☐ Transference
- ☐ Counter-transference

Your answer is incorrect.

Transference is the phenomenon whereby we unconsciously transfer feelings and attitudes from a person or situation in the past on to a person or situation in the present (Retrieved from <http://apt.rcpsych.org/content/aptrcpsych/6/1/57.full.pdf>). Countertransference is the response that is elicited in the recipient (therapist) by the other's (patient's) unconscious transference communications.

The correct answer is: Counter-transference



## Question 63

Not answered

Marked out of 1

All of the following are described by Yalom as group curative factors, except

Select one:

- ☐ Imitation of adaptive behaviour
- ☐ Group cohesion
- ☐ Universality
- ☐ Altruism
- ☐ Identification

Your answer is incorrect.

Yalom cited 11 'curative' factors responsible for the change in groups. The curative factors include the instillation of hope, universality, imparting information (feedback), altruism, corrective recapitulation, socialisation techniques, imitative behaviour, interpersonal learning, group cohesiveness, catharsis and existential factors. Of these, cohesiveness, and learning from feedback are valued positively though other factors may also be important.

The correct answer is: Identification

Question **64**

Not answered

Marked out of 1

Which theory best suits the explanation for someone with an emotionally unstable personality disorder?

Select one:

- ☐ Prochaska's transtheoretical model.
- ☐ Bion's container theory
- ☐ Klein's depressive position
- ☐ Winnicott's false self
- ☐ Klein's paranoid schizoid position

Your answer is incorrect.

Klein's paranoid-schizoid position best explains the splitting, idealisation, denigration and persecutory anxiety seen in borderline personality disorder.

The correct answer is: Klein's paranoid schizoid position





## Question 65

Not answered

Marked out of 1

Your colleague asks you what has been shown to predict a good response to psychotherapy for depression. Your response will be

Select one:

- ☐ Lack of prior exposure to therapy
- ☐ The therapist's contribution to therapeutic alliance
- ☐ Young patients
- ☐ Intelligent patients
- ☐ The patient's perception of the working alliance with the therapist.

Your answer is incorrect.

The relationship between therapeutic alliance and treatment outcome was examined for depressed outpatients who received interpersonal psychotherapy, cognitive-behavior therapy, imipramine with clinical management, or placebo with clinical management in NIMH collaborative trial for depression treatment (Krupnick et al, 2006). Therapeutic alliance was found to have a significant effect on clinical outcome for both psychotherapies and for active and placebo pharmacotherapy. Ratings of patient contribution to the alliance were significantly related to treatment outcome; ratings of therapist contribution to the alliance and outcome were not significantly linked. These results indicate that the therapeutic alliance is a common factor with significant influence on outcome (The role of the therapeutic alliance in psychotherapy and ... (n.d.). Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8698947>). Positive perception of therapist throughout treatment (idealization) is not necessary and may even be counterproductive in some cases.

The correct answer is: The patient's perception of the working alliance with the therapist.

## Question 66

Not answered

Marked out of 1

Which of the following is a crucial step in the psychological treatment of phobias?

Select one:

- ☐ Addressing negative automatic thoughts
- ☐ Analyzing countertransference
- ☐ Psychoeducation & Desensitisation
- ☐ Response prevention following exposure
- ☐ Eliciting cognitive errors

Your answer is incorrect.

The important step and powerful step in the treatment of specific phobia is to begin confronting feared situations in real life. The desensitization process begins with making a hierarchy, a graded list of situations that cause anxiety. Systematic desensitization consists of three steps: relaxation training, hierarchy construction, and desensitization of the stimulus. Patient is exposed to a graded hierarchy of anxiety-provoking situations in stepwise fashion

The correct answer is: Psychoeducation & Desensitisation

Question **67**

Not answered

Marked out of 1

The approach of mentalisation is best utilised in which of the following therapies?

Select one:

- ☐ Psychodynamic therapy
- ☐ Nidotherapy
- ☐ Dialectical behavioural therapy
- ☐ Rational emotive therapy
- ☐ CBT

Your answer is incorrect.

MBT focuses on understanding your mental state, such as emotions, feelings, and thoughts and that of others so you can manage your emotions and improve your relationships. Mentalization-Based Therapy (MBT) provides a way to increase your capacity to make personal attachments and therefore reduce the emotional intensity you experience. Initial research has proved that mentalization is effective for BPD treatment as compared to the usual treatments in the field. It shows significant reduction in suicide attempts and suicidal ideation, hospitalization due to suicidal or feelings of/intent to self-harm, and trips to the emergency room.

The correct answer is: Dialectical behavioural therapy



Question **68**

Not answered

Marked out of 1

Which of the following types of psychotherapy is suitable in a child with learning difficulties and behavioural dyscontrol?

Select one:

- ☐ Behavioural therapy
- ☐ Mentalisation based therapy
- ☐ Cognitive therapy
- ☐ DBT
- ☐ Psychodynamic Psychotherapy

Your answer is incorrect.

Behavioural therapy is suitable in a child with learning difficulties and behavioural dyscontrol. It involves modification of behaviour using operant conditioning (rewarding good behaviour, removal of aversive stimuli, neutral response to attention seeking behaviour), secondary reinforcers, modelling and positive programming.

The correct answer is: Behavioural therapy



## Question 69

Not answered

Marked out of 1

Which of the following modes of therapies has the least empirical evidence in terms of efficacy for borderline personality disorder and not recommended by NICE?

Select one:

- ☐ Transference focussed psychotherapy
- ☐ Mentalisation based therapy
- ☐ Acceptance & commitment therapy
- ☐ DBT
- ☐ CBT

Your answer is incorrect.

Acceptance and Commitment Therapy or ACT has the least empirical evidence in terms of efficacy for borderline personality disorder. It is an empirically-based psychological intervention that uses acceptance and mindfulness strategies mixed in different ways with commitment and behavior-change strategies, to increase psychological flexibility. ACT has shown preliminary research evidence of effectiveness in randomized trials for a variety of problems including chronic pain, addictions, smoking cessation, depression, anxiety, workplace stress, diabetes management, weight management etc

The correct answer is: Acceptance & commitment therapy



[Refund Policy](#) | [Terms & Conditions](#)

