

Started on	Sunday, 12 February 2023, 8:52 PM
State	Finished
Completed on	Sunday, 12 February 2023, 8:53 PM
Time taken	25 secs
Marks	0/5
Grade	0 out of 100

Question 1

Not answered

Marked out of 1

The most important component in the development of a clinical guideline is

Select one:

- ☐ Systematic development
- ☐ Meta-analysis of the available data
- ☐ All stakeholders are consulted.
- ☐ Systematic literature search
- ☐ Economic factors are considered

Your answer is incorrect.

Practice guidelines are usually generated by a panel of stakeholders (e.g. NICE). It may not be always possible to include 'all' stakeholders for a particular issue considered. Guidelines based on same evidence base may vary in recommendations. This is because apart from being evidence based, guidances are also often value based, and local availability (cost) based. Though considering economic factors is important, the most important step in the actual process of developing guidelines is having a systematically reviewed base of evidence on which recommendations can be built. (This MCQ is from a sample provided by the RCPsych)

The correct answer is: Systematic literature search

Question 2

Not answered

Marked out of 1

Which of the following elements does not belong to the process of making treatment recommendations in the form of clinical guidelines?

Select one:

- ☐ Defining the decision to be made
- ☐ Thorough appraisal of available evidence based literature
- ☐ Incorporating values to decide course of action
- ☐ Preparing recommendations irrespective of local resources
- ☐ Specifying possible outcomes of a decision

Your answer is incorrect.

It is important to consider the availability of resources before drafting recommendations or guidelines. If certain resources are not available, one may wish to modify a recommended action even if it is based on the best evidence literature. (e.g. no depot clinics locally, no CPNs in an area, etc.).

The correct answer is: Preparing recommendations irrespective of local resources

Question 3

Not answered

Marked out of 1

A 77-year-old man falls out of NICE guidelines to prescribe antidementia medications. But he can get the same medications if he lived in France or Canada under their own national guidelines. How can this be explained in terms of the process of guideline production?

Select one:

- ☐ Antidementia drugs do not work that well in UK born patients
- ☐ Stakeholders such as Alzheimer's society were not consulted by NICE in UK
- ☐ Inadequate appraisal of literature took place while drafting NICE guidelines in UK
- ☐ Different set of research evidences were appraised by the different national bodies
- ☐ Guidelines are not only based on evidence but also resource availability, values and collective preferences.

Your answer is incorrect.

Steps to be followed in producing a treatment recommendation: Defining the decision (e.g. decision to switch antipsychotics when first episode of psychosis is not controlled even after 12 weeks of treatment) Specify the possible outcomes (control of symptoms, no effect of switch, increase in side effects, noncompliance, occurrence of depression etc). Using available evidence to specify the link between listed options and outcomes. (Olanzapine causes weight gain, lesser dropout rate for olanzapine and depots, etc.'') Incorporate values/value judgements to decide on optimal course of action (Female patient, poor lifestyle - weight gain is an important problem, etc) Consider local circumstances including resource availability and modify action if needed (no depot clinics locally, no CPNs etc). This might explain the differences in recommendation despite appraising same evidence base.

The correct answer is: Guidelines are not only based on evidence but also resource availability, values and collective preferences.

Question 4

Not answered

Marked out of 1

A new national guideline on management of Lewy Body dementia has been published. An NHS trust is keen to adapt these guidelines locally. Which of the following ways of implementation can ensure maximum effectiveness of the guidelines?

Select one:

- ☐ Setting up patient specific reminder during each new assessment
- ☐ Placing a bright yellow coloured placard in the local memory clinic
- ☐ Pharmacy sending patient specific feedback to clinicians citing the guidelines
- ☐ Setting up an ultimatum locally for using the guidelines
- ☐ Providing general feedback by auditing the effect before and after introduction of guidelines

Your answer is incorrect.

Guidelines are highly effective if they are; Developed by those who will use them Disseminated via a specific educational intervention e.g. focussed meeting, learning package with credits Implemented through patient-specific reminder during consultation e.g. filed in the notes etc.

The correct answer is: Setting up patient specific reminder during each new assessment



Question 5

Not answered

Marked out of 1

Which of the following best describes the Delphi method in clinical research

Select one:

- ☐ A cluster group technique in economic analysis
- ☐ A descriptive graph for survival analysis
- ☐ A consensus method for developing treatment guidelines
- ☐ A meta analysis technique for observational data
- ☐ A statistical technique for converting qualitative research to numerical data

Your answer is incorrect.

The Delphi survey is a group facilitation technique, which is an iterative multistage process, designed to transform opinion into group consensus. It is a flexible approach that is used commonly within the health and social sciences(Research guidelines for the Delphi survey technique. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11095242>)

The correct answer is: A consensus method for developing treatment guidelines



© 2023 SPMM Course Limited, UK.

[Refund Policy](#) | [Terms & Conditions](#)

