

Started on	Sunday, 12 February 2023, 8:56 PM
State	Finished
Completed on	Sunday, 12 February 2023, 8:56 PM
Time taken	11 secs
Marks	0.00/81.00
Grade	0.00 out of 100.00

Question 1

Not answered

Marked out of 1.00

The onset of very early onset schizophrenia is before the age of

Select one:

- ☐ 13
- ☐ 9
- ☐ 10
- ☐ 12
- ☐ 18

Your answer is incorrect.

In very early onset schizophrenia, the onset of psychotic symptoms is at or before the age of 13. Early onset schizophrenia is a term used to indicate the onset of psychotic symptoms before the age of 18.

The correct answer is: 13

Question **2**

Not answered

Marked out of 1.00

Tom, aged 12, has been diagnosed with NREM sleep disorder. The most likely diagnosis is;

Select one:

- ☐ Sleep walking
- ☐ Sleep Bruxism
- ☐ REM behavioural disorder
- ☐ Nightmares
- ☐ Sleep paralysis

Your answer is incorrect.

NREM Sleep disorders include Sleep walking, sleep terrors and confusion arousals. REM sleep parasomnias include REM behavioural disorder, Night mares and Sleep paralysis.

The correct answer is: Sleep walking



Question **3**

Not answered

Marked out of 1.00

Which of the following drug should be avoided in an ADHD child with liver dysfunction?

Select one:

- ☐ Atomoxetine
- ☐ Methylphenidate
- ☐ Clonidine
- ☐ Amitriptyline
- ☐ Dexamphetamine

Your answer is incorrect.

Atomoxetine should be avoided in children with preexisting liver disease. Parents should be warned of the possibilities of liver disease emerging and advised of the possible features they might notice. LFTs monitoring is needed.

The correct answer is: Atomoxetine



Question 4

Not answered

Marked out of 1.00

Which of the following is a feature of dual role transvestism?

Select one:

- ☐ Male wearing clothes of the other sex without desire for permanent change and no sexual arousal
- ☐ Male with persistent discomfort with his sex
- ☐ Male wearing clothes of the other sex to achieve sexual pleasure and associated with masturbation
- ☐ Male using shoes to obtain sexual pleasure
- ☐ Male with a persistent preoccupation with castration

Your answer is incorrect.

Various degrees of gender dysphoria exist. One mild form is recognized in ICD and DSM as dual role transvestism. The individual wears clothes of the opposite sex in order to experience temporary membership in the opposite sex. The individual experiences a sense of appropriateness by wearing clothes of the other gender. There is no sexual motivation for the cross-dressing. The individual has no desire for a permanent change to the opposite sex.

The correct answer is: Male wearing clothes of the other sex without desire for permanent change and no sexual arousal

Question 5

Not answered

Marked out of 1.00

Which of the following is most predictive of nocturnal enuresis in children?

Select one:

- ☐ Large family size
- ☐ Sleep disorder
- ☐ Hirschsprungs disease
- ☐ Eating of non nutritive substances
- ☐ Low birth weight babies

Your answer is incorrect.

Association of enuresis has been made with a high rate of stressful life events, UTI, constipation , and with children of low socioeconomic background, living in large families or overcrowded conditions.

The correct answer is: Large family size



Question 6

Not answered

Marked out of 1.00

Which of the following is not a feature seen in Oppositional Defiant Disorder?

Select one:

- ☐ Often been physically cruel to animals
- ☐ Often loses temper
- ☐ Often deliberately annoys people
- ☐ Often refuses to comply
- ☐ Often blames others

Your answer is incorrect.

Physical cruelty to animals is a feature seen in children with conduct disorder

The correct answer is: Often been physically cruel to animals



Question 7

Not answered

Marked out of 1.00

Which of the following psychotic disorders with onset in childhood has the worst prognosis?

Select one:

- ☐ Schizophrenic psychosis
- ☐ Depressive Psychosis
- ☐ Schizoaffective psychosis
- ☐ Drug-induced psychosis
- ☐ Bipolar Psychosis

Your answer is incorrect.

There is limited data on recovery rates from childhood onset schizophrenia but given that the emergence of the disorder disrupts psychosocial development and schooling. It is unsurprising the very early onset schizophrenia and early onset schizophrenia often has a relatively poor prognosis even with psychopharmacological treatment. 80-90% will have 2 or more episodes in a 5-year period. Presence of affective symptoms offers a better outcome.

The correct answer is: Schizophrenic psychosis

Question 8

Not answered

Marked out of 1.00

A teenager attends the CAMHS clinic for sleep problems and noted to have poor sleep hygiene. Which of the following advice should be given as a sleep hygiene measure?

Select one:

- ☐ Avoid large meals prior to bed
- ☐ Having a cold bath 1-2 hours before bedtime
- ☐ Try to do strenuous exercise in the 4 hours before bedtime
- ☐ Try to sleep even when you are not actually feeling tired or sleepy
- ☐ If you have a bad night sleep, change your daytime routine

Your answer is incorrect.

"A healthy, balanced diet will help you to sleep well, but timing is important. Some people find that a very empty stomach at bedtime is distracting, so it can be useful to have a light snack, but a heavy meal soon before bed can also interrupt sleep. Only try to sleep when you actually feel tired or sleepy, rather than spending too much time awake in bed. Even if you have a bad night sleep and are tired it is important that you try to keep your daytime activities the same as you had planned. That is, don't avoid activities because you feel tired. This can reinforce the insomnia. Having a hot bath 1-2 hours before bedtime can be useful, as it will raise your body temperature, causing you to feel sleepy as your body temperature drops again". (Excerpt from: www.cci.health.wa.gov.au Also see: Tan E, et al. BMC pediatrics. 2012 Dec 7;12(1):1.)

The correct answer is: Avoid large meals prior to bed

Question 9

Not answered

Marked out of 1.00

A young boy is referred to your CAMHS clinic. He has significant communication difficulties and more recently has been suffering from seizures. What is the most common feature of seizures in autism?

Select one:

- ☐ Generalised tonic-clonic type
- ☐ Predominance in males
- ☐ Onset before the age of 10
- ☐ Complex partial type
- ☐ Affect 75% of autistic individuals

Your answer is incorrect.

Roughly a third of individuals with autism will develop seizures. In the majority, seizures will begin after the age of 10, generalized tonic-clonic seizures predominate and prevalence is greater in females. (Bolton et al, 2011)

The correct answer is: Generalised tonic-clonic type



Question 10

Not answered

Marked out of 1.00

What specific dietary advice would you give to a mother whose child has been diagnosed with ADHD?

Select one:

- ☐ Add L-carnitine
- ☐ Omit complex carbohydrates
- ☐ Avoid refined sugars
- ☐ Omit proteins
- ☐ Stress the need for a balanced diet

Your answer is incorrect.

Studies on ADHD children showed that a well-balanced diet, including vegetables, complex carbohydrates, fruits, and plenty of protein, helped to bring hyperactive behaviour under control consistently.

The correct answer is: Stress the need for a balanced diet



Question **11**

Not answered

Marked out of 1.00

Which one among the following is an advanced sign of opioid withdrawal

Select one:

- ☐ Muscle spasm
- ☐ Lacrimation
- ☐ Piloerection
- ☐ Tachycardia
- ☐ Yawning

Your answer is incorrect.

The opioid withdrawal symptoms are given below; the early signs of withdrawal include anxiety & craving for opioids, fear of withdrawal. Agitation, insomnia, headache, lacrimation, rhinorrhoea, yawning, shivering and perspiration follow this. Dilated pupils and piloerection are late signs followed by tachycardia, increase in respiratory rate and abdominal cramps. The Advanced sign of opioid withdrawal is Muscle spasm.

The correct answer is: Muscle spasm

Question 12

Not answered

Marked out of 1.00

Which street drug, when taken, results in a 'fishy smell' in sweat and runny nose, with the user sometimes presenting with hallucinations?

Select one:

- ☐ Mephedrone
- ☐ Crack Cocaine
- ☐ Heroin
- ☐ Amphetamine
- ☐ MDMA

Your answer is incorrect.

Mephedrone, also known as meow-meow, is a white crystal or a powdery substance, sometimes sold as capsules or pills. It has a distinctive fishy odour though some products are described to have vanilla, bleach, or stale urine smell. This smell is often notable in the sweat of the user. Ref: <http://www.hmdb.ca/metabolites/HMDB41923>

The correct answer is: Mephedrone



Question **13**

Not answered

Marked out of 1.00

You are seeing a 47 year old man in the outpatients clinic. He is taking a drug called Varenicline. It is used in the treatment of which drug addiction

Select one:

- ☐ Nicotine
- ☐ Alcohol
- ☐ Cannabis
- ☐ Cocaine
- ☐ Amphetamines

Your answer is incorrect.

Varenicline tartrate is a partial agonist binding with high affinity to $\alpha 4 \beta 2$ nicotinic acetylcholine receptor. It significantly reduces nicotine withdrawal symptoms and makes smoking less rewarding so it may help prevent 'slips' developing into full relapse.

The correct answer is: Nicotine



Question 14

Not answered

Marked out of 1.00

Which of the following is true about Korsakoff's psychosis?

Select one:

- ☐ All patients need residential care
- ☐ All patients develop psychosis
- ☐ Thiamine is the treatment of choice
- ☐ About 10 % alcoholic patients recovering from Wernicke's encephalopathy develop Korsakoff's syndrome.
- ☐ Marked deficits noted in attention, procedural and working memory

Your answer is incorrect.

About 80 per cent of alcoholic patients recovering from Wernicke's encephalopathy develop Korsakoff's amnesic syndrome. It is characterized by marked deficits in anterograde and to some extent retrograde episodic memory, apathy, an intact sensorium, and relative preservation of other intellectual abilities such as attention, procedural memory and working memory. Diagnosis of Korsakoff's syndrome correlates best with the presence of lesions in the dorsomedial thalamus. But lesions in the mammillary bodies, the mammillo-thalamic tract and the anterior thalamus may be more important to memory dysfunction than lesions in the medial dorsal nucleus of the thalamus. Confabulation occurs due to a double lesion – frontal deficits and diencephalic deficits. In terms of treatment, continue oral thiamine replacement for up to 2 years.

The correct answer is: Thiamine is the treatment of choice

Question 15

Not answered

Marked out of 1.00

You are trying to promote safer drinking behaviours in a small community and decide to introduce a programme to schools in the area. School-based intervention for children and adolescents must include which of the following?

Select one:

- ☐ Include all age groups
- ☐ Exploration of the attitudes of children towards drinking
- ☐ Reinforce social drinking
- ☐ Include those with alcohol dependence
- ☐ Be led by parents

Your answer is incorrect.

Child and adolescent behaviour modification can occur at the national, school and individual level, depending on the type of intervention. School-based interventions would best; be focused on specific age groups (as the issue surrounding drinking will not apply to children of all ages), be more general (as any pupil with alcohol dependence will require an individual approach), explore existing attitudes of children towards drinking and be facilitated by a nominated member of staff or visiting expert speaker.

The correct answer is: Exploration of the attitudes of children towards drinking

Question 16

Not answered

Marked out of 1.00

Which one among the following test is a specific biomarker for heavy alcohol misuse?

Select one:

- ☐ Increased D- Dimer
- ☐ Increased carbohydrate deficient transferrin
- ☐ Reduced MCV
- ☐ Increased troponin levels
- ☐ Reduced GGT

Your answer is incorrect.

Carbohydrate Deficient Transferrin (CDT) is a biomarker for heavy alcohol consumption. The Carbohydrate Deficient Transferrin (CDT) test is a particularly specific in tracking heavy alcohol consumption over an extended period of time. The advantage of using CDT in testing for alcohol misuse is it is unaffected by mild alcohol related liver disease or fatty liver (due to obesity, diabetes etc) which can falsely increase γ -glutamyl transferase (GGT), the enzyme in blood most commonly used to indicate excess alcohol consumption. CDT is also unaffected by chronic disease or B12/folate deficiencies which can increase the mean corpuscular volume (MCV), another blood test used to detect alcohol misuse. Transferrin remains in the circulation for 7 to 14 days and CDT, therefore, can give an indication of alcohol consumption over this period of time, which gives it an advantage over blood or urine alcohol measurements, which only remain elevated for 24-48 hours after alcohol consumption. (Bergström, J.P. and Helander, A., 2008. Alcohol and alcoholism, 43(4), pp.436-441.)

The correct answer is: Increased carbohydrate deficient transferrin

Question 17

Not answered

Marked out of 1.00

You have received five calls from the Emergency Department to attend five separate cases. You are required to prioritise these cases. Which will you attend first ?

Select one:

- ☐ A 55-year-old male with schizophrenia and diabetes who is complaining of a sudden onset sharp left-sided chest pain
- ☐ A 28-year-old female with borderline personality disorder admitted for superficial lacerations to her forearm who is threatening to leave the ward
- ☐ A 45-year-old female with schizoaffective disorder who is attempting to hit the nurse
- ☐ An 88-year-old male with delirium who suffered a fall on the ward and is now complaining of pain in the hip
- ☐ A 37-year-old female with depression who is refusing to take her antidepressant medication

Your answer is incorrect.

This patient's presentation is typical of an acute myocardial infarction. Compared to the other patients in the list, he warrants the most immediate attention as his presentation is the most life-threatening.

The correct answer is: A 55-year-old male with schizophrenia and diabetes who is complaining of a sudden onset sharp left-sided chest pain



Question **18**

Not answered

Marked out of 1.00

A 23-year-old female student with anorexia nervosa has recently been admitted to your Eating Disorders unit. Which of the following would lead you to immediately referring this patient to a medical hospital?

Select one:

- ☐ Her blood pressure is 90/60 mmHg
- ☐ A potassium level of 2.6 mmol/L
- ☐ She has a BMI of 15
- ☐ Her heart rate is 58 beats per minute
- ☐ She refused to eat breakfast

Your answer is incorrect.

In patients with anorexia nervosa, hospital admission should only be considered in the presence of serious medical or psychiatric problems. This patient's blood test revealed a severe hypokalemia placing her at a greater risk of cardiac complications such as arrhythmias, which is the most common cause of death in patients with anorexia nervosa (mortality rate ~ 10%) (Ref: Oxford Handbook of Psychiatry).

The correct answer is: A potassium level of 2.6 mmol/L

Question 19

Not answered

Marked out of 1.00

You are reviewing a 52-year-old lady with paranoid schizophrenia who requires medical treatment for her symptoms. She has a history of breast cancer. Which of the following would be the LEAST appropriate medication to prescribe her?

Select one:

- ☐ Amisulpride
- ☐ Clozapine
- ☐ Olanzapine
- ☐ Aripiprazole
- ☐ Quetiapine

Your answer is incorrect.

Treatment with amisulpride can induce hyperprolactinaemia, a long-term consequence of which is a possible increase in the risk of breast cancer. The antipsychotics that are not usually associated with hyperprolactinaemia include Aripiprazole, Olanzapine, Quetiapine and Clozapine (for more details see SPMM Smart Revise Notes)

The correct answer is: Amisulpride



Question **20**

Not answered

Marked out of 1.00

Anorexia nervosa in adulthood is most commonly associated with which of the following?

Select one:

- ☐ Bulimia in childhood
- ☐ Paranoid schizophrenia
- ☐ Affective disorders
- ☐ High socioeconomic status
- ☐ Anxiety disorders

Your answer is incorrect.

Anorexia nervosa is most commonly associated with obsessive and other anxious personality traits.

The correct answer is: Anxiety disorders



Question **21**

Not answered

Marked out of 1.00

Sally has a long standing of paranoid schizophrenia. He was reviewed in the clinic for abnormal movements in her body and diagnosed with tardive dyskinesia. Which antipsychotic would be suitable to be used in this situation?

Select one:

- ☐ Haloperidol
- ☐ Clozapine
- ☐ Risperidone
- ☐ Amisulpride
- ☐ Zuclopenthixol

Your answer is incorrect.

The use of Clozapine is probably best supported in this regard, but Quetiapine, another weak striatal dopamine antagonist is also effective. (Ref: The Maudsley prescribing guidelines-11th edition; 107)

The correct answer is: Clozapine



Question **22**

Not answered

Marked out of 1.00

Which one among the following is a significant risk factor for the development of Neuroleptic malignant syndrome?

Select one:

- ☐ Dehydration
- ☐ Being on oral medication
- ☐ Female Patient
- ☐ History of personality disorder
- ☐ Low ambient temperature

Your answer is incorrect.

Risk factors for NMS: High ambient temperature, dehydration, patient agitation or catatonia, rapid antipsychotic initiation/dose escalation, withdrawal of anti-parkinsonian medication, use of high potency agents/depot IM preparations, history of organic brain disease such as dementia, history of affective disorder, previous history of NMS, Predisposing drugs like lithium and anticholinergic medications. (Ref: Oxford Handbook of Psychiatry-882)

The correct answer is: Dehydration

Question **23**

Not answered

Marked out of 1.00

Severe lithium toxicity presents with which of the following signs?

Select one:

- ☐ T-wave elevation
- ☐ Hypo active deep tendon reflexes
- ☐ Coloured vision
- ☐ Constipation
- ☐ Downbeat nystagmus

Your answer is incorrect.

Central Nervous System features of severe lithium toxicity: Blackout spells, epileptiform seizures, slurred speech, dizziness, vertigo, incontinence of urine or feces, somnolence, psychomotor retardation, restlessness, confusion, stupor, coma, acute dystonia, downbeat nystagmus. (Ref: McKnight RF, et al. Lithium toxicity profile: a systematic review and meta-analysis. The Lancet. 2012 Mar 2;379(9817):721-8.)

The correct answer is: Downbeat nystagmus

Question **24**

Not answered

Marked out of 1.00

Mary is a 42-year-old woman with bipolar disorder with frequent depressive episodes. What drug would you choose as the best prophylaxis against depression?

Select one:

- ☐ Carbamazepine
- ☐ Lamotrigine
- ☐ Sodium Valproate
- ☐ Topiramate
- ☐ Haloperidol

Your answer is incorrect.

Lamotrigine has good evidence in prophylaxis in bipolar affective disorder, especially when depressive episodes predominate.

The correct answer is: Lamotrigine



Question **25**

Not answered

Marked out of 1.00

A 26-year-old mother suffered a traumatic birth. For the last 3 months, she is experiencing intrusive thoughts of harming her 1-year-old baby and has flashbacks. She finds these thoughts anxiety provoking. She regards the birth as traumatic. What is her diagnosis?

Select one:

- ☐ Schizophrenia
- ☐ OCD
- ☐ Postnatal depression
- ☐ Generalized anxiety disorder
- ☐ PTSD

Your answer is incorrect.

Intrusive thoughts that cause anxiety are a feature of OCD, and there is a greater predisposition to this distressing experience following a traumatic birth.

The correct answer is: OCD



Question **26**

Not answered

Marked out of 1.00

What side effect would you monitor closely in a patient taking Donepezil?

Select one:

- ☐ Reduced Pulse rate
- ☐ Seizures
- ☐ Rash
- ☐ Tremors
- ☐ Elevated Blood pressure

Your answer is incorrect.

Acetyl cholinesterase inhibitors (AChEs) like Donepezil may have vagotonic effects on heart rate leading to bradycardia. AChE inhibitors can induce sinus bradycardia, sino-atrial block, and aggravate pre-existing sinus node disease and atrioventricular block (Savie et al, 1998). Donepezil overdose has been reported to result in profound sinus bradycardia that can be reversed with atropine (Shepherd et al, 1999; Hidetaka et al, 2003). Therefore it is important to monitor for signs like bradycardia.

The correct answer is: Reduced Pulse rate

Question **27**

Not answered

Marked out of 1.00

What type of memory is most affected by ECT?

Select one:

- ☐ Implicit memory
- ☐ Retrograde memory
- ☐ Semantic memory
- ☐ Episodic Memory
- ☐ Anterograde memory

Your answer is incorrect.

Retrograde memory i.e. the recall of events that occurred before the treatment, is the most commonly affected aspect of memory following Electro-Convulsive Therapy (ECT). It is an early side effect, occurs in 64% of patients receiving ECT and usually resolves completely.

The correct answer is: Retrograde memory



Question **28**

Not answered

Marked out of 1.00

Which of the following strategies has good evidence in prescribing in treatment resistant depression?

Select one:

- ☐ SSRI + SNRI
- ☐ SSRI + TCA
- ☐ SSRI + SSRI
- ☐ SNRI + NASSA
- ☐ SSRI+ MAOI

Your answer is incorrect.

"California rocket fuel" is a medical slang term for the combined administration of mirtazapine (a NaSSA) and venlafaxine (an SNRI). This combination is usually used in the treatment of treatment-resistant depression. It is usually well tolerated, has excellent literature support available and is widely used. However the patient should be informed of the theoretical risk of serotonin syndrome.

The correct answer is: SNRI + NASSA

Question **29**

Not answered

Marked out of 1.00

George is a 50-year-old man with a history of bipolar disorder. He recently had a renal transplant and subsequently became manic. He is already on an antipsychotic medication. Which mood stabiliser would you prescribe in this regard?

Select one:

- ☐ Lamotrigine
- ☐ Lithium
- ☐ Sodium valproate
- ☐ Gabapentin
- ☐ Topiramate

Your answer is incorrect.

With sodium valproate, approximately 2% excreted unchanged. Dose adjustments usually not required in renal impairment; however, free valproate levels may be increased. Therefore start at a low dose and increase slowly. ((Ref: The Maudsley prescribing guidelines-11th edition, 472)

The correct answer is: Sodium valproate

Question **30**

Not answered

Marked out of 1.00

Tom is a 37-year-old man with bipolar disorder and is on Lithium treatment. He is now presenting with mixed affective state. His recent Lithium level is 0.8. What would be your next step?

Select one:

- ☐ Start antidepressant medication
- ☐ Start sodium valproate
- ☐ Start antipsychotic medication
- ☐ Reduce the dose of lithium
- ☐ Increase the dose of lithium

Your answer is incorrect.

Mixed affective state should be treated as manic episode. The best strategy in this situation is to start him on an antipsychotic medication. Lithium levels are optimal and to increase could risk toxicity.

The correct answer is: Start antipsychotic medication



Question **31**

Not answered

Marked out of 1.00

Sally is a bipolar disorder patient who is currently on citalopram and lithium. She has relapsed into mania. What would be your next step?

Select one:

- ☐ Stop lithium completely
- ☐ Start her on another mood stabilizer
- ☐ Start on antipsychotic medication
- ☐ Stop the antidepressant medication
- ☐ Increase the dose of lithium

Your answer is incorrect.

Although rare, manic switch secondary to SSRIs (like citalopram) is well recognised and simple cessation may result in stabilisation.

The correct answer is: Stop the antidepressant medication



Question **32**

Not answered

Marked out of 1.00

Paul is a 4 year old boy with impaired social and communication skills. His mother is also concerned about his insomnia. The drug of choice would be;

Select one:

- ☐ Melatonin
- ☐ Lorazepam
- ☐ Diazepam
- ☐ Risperidone
- ☐ Citalopram

Your answer is incorrect.

This is a case of insomnia with autistic spectrum disorder. Children with autistic spectrum disorder have significant sleep problems. Melatonin has been shown in 17 studies to be beneficial in children with autistic spectrum disorder. Doses ranged from 1mg to 10 mg and they are very well tolerated. (Ref: The Maudsley prescribing guidelines; 11th edition: page 340)

The correct answer is: Melatonin

Question **33**

Not answered

Marked out of 1.00

Which of the following crimes is more common in people with Asperger's syndrome

Select one:

- ☐ Property destruction
- ☐ Motoring offences
- ☐ Fraud
- ☐ Sexual offences
- ☐ Drug offences

Your answer is incorrect.

From the given list, a person with Asperger's syndrome is more likely to conduct offences against property. Arson is also increased in prevalence among autistic spectrum disorders, but not as much as property damages.

The correct answer is: Property destruction



Question **34**

Not answered

Marked out of 1.00

Which scale is helpful to assess IQ of an 8-year-old child?

Select one:

- ☐ WISC
- ☐ Connors rating scale
- ☐ ADOS
- ☐ Strengths and difficulties questionnaire
- ☐ Behavioural assessment tool

Your answer is incorrect.

WISC stands for Wechsler Intelligence scale for children. WISC is an individually administered intelligence test for children between the ages of 6 and 16, which can be completed without reading or writing.

The correct answer is: WISC



Question **35**

Not answered

Marked out of 1.00

'Wringing hand' movements are associated with which of the following conditions?

Select one:

- ☐ Rett syndrome
- ☐ Angelman syndrome
- ☐ Edwards syndrome
- ☐ Prader Willi syndrome
- ☐ Smith Magenis syndrome

Your answer is incorrect.

Rett syndrome a type of pervasive developmental disorder that is seen exclusively in females. Repetitive stereotypic hand movements such as wringing and or repeatedly putting hand into the mouth are noted.

The correct answer is: Rett syndrome



Question **36**

Not answered

Marked out of 1.00

In which situation would it be LEAST appropriate to breach confidentiality?

Select one:

- ☐ A 26-year-old male, manic patient who is promiscuous and does not wear a condom despite harbouring a sexually transmitted infection
- ☐ A 19-year-old patient, recently diagnosed with schizophrenia who threatens to kill his brother
- ☐ A 76-year-old patient, recently diagnosed with Alzheimer's Disease who refuses to inform the DVLA despite your advice, and continues to drive
- ☐ A 37-year-old female patient who admitted to shoplifting
- ☐ A 47-year-old male patient who admitted to sexually abusing his five-year-old next door neighbor

Your answer is incorrect.

All the other scenarios put the patients and others at a significantly greater risk, requiring us to act with immediacy. Our duty of care as physicians encompasses a responsibility to protect the interests of both the patient and the public.

The correct answer is: A 37-year-old female patient who admitted to shoplifting



Question **37**

Not answered

Marked out of 1.00

'Sequential Diagrammatic reformulation' (SDR) is seen in what type of therapy?

Select one:

- ☐ Cognitive Behavioural therapy
- ☐ Dialectical behavioural therapy
- ☐ Psychodynamic psychotherapy
- ☐ Cognitive analytical therapy
- ☐ Interpersonal therapy

Your answer is incorrect.

Sequential Diagrammatic Reformulation (SDR) is a technique used in Cognitive analytical therapy. It involves collaboratively developed visual map of problematic patterns and reciprocal roles. The procedures are depicted and interlinked in flow-chart format. The aim of such diagrams is to amplify understanding of often-complex procedural sequences and their inter-relationships in a way that words alone cannot. SDR's are often elaborated by around the sixth or seventh session. (Ref: Denman, C., 2001. Cognitive-analytic therapy. *Advances in Psychiatric treatment*, 7(4), pp.243-252.)

The correct answer is: Cognitive analytical therapy

Question **38**

Not answered

Marked out of 1.00

Tom is a 12-year-old boy with severe depression. He was offered CBT that has failed. His parents are worried about side effects of medication and don't want him to have pharmacotherapy. The best alternative in this situation is

Select one:

- ☐ Cognitive analytical therapy
- ☐ Dialectical behavioural therapy
- ☐ Interpersonal therapy
- ☐ Parenting training
- ☐ Mindfulness training

Your answer is incorrect.

Interpersonal Therapy (IPT) and Transactional Analysis focus in particular on interpersonal relationships, and both may be effective in cases of depression. IPT has been adapted for use with children and adolescents. The areas of focus in IPT include conflicts with others, losses, changes in circumstances/difficulties adapting to new circumstances, and social isolation. Transactional Analysis looks at the way in which people take up roles of Parent, Adult or Child, exploring inner dialogue between different components of the self.

The correct answer is: Interpersonal therapy



Question 39

Not answered

Marked out of 1.00

Katie's mother had just died. However, she appears elated and happy to be keeping busy with funeral arrangements. The type of defense mechanism used in this situation is called

Select one:

- ☐ Suppression
- ☐ Omnipotence
- ☐ Identification with the aggressor
- ☐ Sublimation
- ☐ Manic Defense

Your answer is incorrect.

Ego defences exist in order to protect us from aspects of reality that might overwhelm us if we made too much contact with them. The defence mechanism employed in this situation is called Manic defence. The potential sadness and anguish about the mother's death is defended against by 'acting out the opposite': a sense of elation.

The correct answer is: Manic Defense



Question 40

Not answered

Marked out of 1.00

John was receiving psychodynamic psychotherapy. He started feeling depressed when his therapist mentioned he was making significant progress. This phenomenon is called

Select one:

- ☐ Dream interpretation
- ☐ Negative therapeutic reaction
- ☐ Transference
- ☐ Primary process thinking
- ☐ Counter transference

Your answer is incorrect.

Negative therapeutic reaction is a paradoxical phenomenon whereby a worsening of the patient's condition follows an acknowledgement of progress. Freud thought it was linked to unconscious guilt about the progress during psychotherapy.

The correct answer is: Negative therapeutic reaction



Question **41**

Not answered

Marked out of 1.00

The beneficial effects of a drug can be readily explained to the patients in terms of

Select one:

- ☐ Number needed to treat
- ☐ Side effects
- ☐ Cost
- ☐ Frequency
- ☐ Dose

Your answer is incorrect.

The number needed to treat (NNT) is the average number of patients who need to be treated to prevent one additional bad outcome (e.g. the number of patients that need to be treated for one to benefit compared with a control in a clinical trial). The ideal NNT is 1, where everyone improves with treatment and no one improves with control. The higher the NNT, the less effective is the treatment. (Ref: Bandolier 2009)

The correct answer is: Number needed to treat

Question 42

Not answered

Marked out of 1.00

The prevalence of a condition is 10%. A new diagnostic test ('test A') has a sensitivity of 95% and false positive rate of 5%. What is the post test probability?

Select one:

- ☐ 68% or 0.68
- ☐ 48% or 0.48
- ☐ 58% or 0.58
- ☐ 78% or 0.78
- ☐ 19% or 0.19

Your answer is incorrect.

According to Bayes' Theorem, if you know the pretest probability and likelihood ratio of a test, you can find the post test probability. There are two ways of computing post-test probability. If you have Bayesian nomogram, you can simply connect the dots indicating pretest probability and LR value, to get the third point indicating post-test probability. If you don't have a nomogram at hand, you have to then use the following formula: $\text{post-test odds} = \text{pretest odds} \times \text{LR}$ (Note that this formula uses odds, not probability rates). In this question, prevalence = pretest probability = 10/100. So pre test odds = 10/90 (as odds = probability / [1-probability]) $\text{LR+} = \text{true positivity rate} / \text{false positivity rate}$ (i.e. $\text{sensitivity} / [1 - \text{specificity}] = 95\% / 5\% = 19$ So, $\text{post-test odds} = \text{pretest odds} \times \text{LR} = (10/90) \times 19 = 190/90 = 19/9$. If you know the odds of an event, you can easily compute the probability. Use the formula $\text{probability} = \text{odds} / [1 + \text{odds}]$. So post test probability = $(19/9) / [1 + (19/9)] = 19/28 = 67.8\%$ nearly 68%

The correct answer is: 68% or 0.68

Question **43**

Not answered

Marked out of 1.00

Which of the following constitutes primary prevention of pre-natal depression?

Select one:

- ☐ Referral to the CMHT
- ☐ Pre-conception counseling
- ☐ Referral for Cognitive-Behavioural Therapy
- ☐ Admission after self-harm
- ☐ Treatment with antidepressants

Your answer is incorrect.

Primary prevention involves preventing the onset of illness with the overall aim of reducing the incidence of illness. Interventions are implemented before any evidence of disease. Pre-conception counseling is a form of pre-conceptual care that allows women and their partners to understand health issues which could affect them in pregnancy and encourages them to take measures to prepare actively for pregnancy, and be as healthy as possible. Based on their past psychiatric history, it identifies couples at increased risk of psychiatric complications during pregnancy and optimizes management of chronic mental illness. (Ref: Johnson, K., et al., 2006. Morbidity and Mortality Weekly Report, 55(4), pp.1-23.)

The correct answer is: Pre-conception counseling

Question 44

Not answered

Marked out of 4.00

Theme: Statistical Tests V

For each of the following vignettes choose the most appropriate statistical test/s.

1. Comparison of the risk of developing schizophrenia in two groups of 3000 people - one group comprises of people who have used cannabis and the other of people who have never used cannabis. ✖
2. Investigation of the association between several risk factors and the risk of developing schizophrenia. ✖
3. Comparing baseline and endpoint weight in the same individuals in a phase II study of an antipsychotic. The variable is normally distributed. (Choose TWO) ✖ ✖

Risk is a proportion; comparing proportions as in question 1 can be done using Chi-square test. In question 2, developing schizophrenia is a binary outcome (present or absent). Hence logistic regression is useful as a multivariate regression technique when several risk factors are evaluated. In question 3, two mean weight values are compared - but one measure is dependent on the other (paired). Hence either paired t test or repeated measures ANOVA can be used.

Correct Answer is : 1. Chi-squared. 2. Logistic regression. 3. ANOVA., Paired t-test.

Question 45

Not answered

Marked out of 4.00

Theme: Statistical Concepts and Definitions I

Give the appropriate measure from the list above that corresponds to the description below.

1. The spread of the raw data (Choose TWO)    
2. The range in which 19/20 identical experiments would be expected to find the same result.  
3. Synonymous with type I error rate.  

Explanations: Both standard deviation and variance are measures of the spread of raw data. The "margin of error" of a measurement obtained from a research sample represents the confidence interval. Using this, the researchers display their confidence that if other studies are done, then 95 per cent of the time - or 19 times out of 20 - the findings would fall in this range. The 95 per cent confidence level is used most often in research; it is a generally accepted standard. However, researchers can calculate CIs at any level of significance, such as 90 per cent or 99 per cent. The significance level simply indicates how precise they are willing to be. P value (generally 5%) is synonymous with type I error rate. It reflects the probability of making a false positive conclusion from the results.

Correct Answer is : 1. Standard Deviation, Variance 2. 95% confidence interval 3. P-value


Question 46

Not answered

Marked out of 7.00

RCTs of Antidepressants: Thirty-Year Meta-Analytic Review

The attached figure is obtained from Undurraga, J & Baldessarini RJ. Neuropsychopharmacology. 2012 Mar; 37(4): 851-864. Using the data presented in the figure choose the best answers from the above list.



For each of the following questions choose the correct option from the responses above

1. Choose the antidepressant drug with best efficacy when compared to placebo. (Choose ONE)



2. Choose the antidepressant drugs with highest weighting in the meta-analysis (Choose TWO)

3. Which drug with n>2 trials has the smallest effect size? (Choose ONE)

4. Choose an antidepressant drug with trial results favouring placebo (Choose ONE)

5. Choose the antidepressant drugs with highest number of eligible trials (Choose TWO)

×

Explanation: The forest plot provides a summary of relative risk of treatment response for individual antidepressants in comparison with placebo. For each drug, the number of available RCTs is indicated in parentheses. Several drugs have had only single eligible RCTs (Phenelzine, Clomipramine, Tranylcypromine, Trazodone and Nortriptyline); these drug trials are represented by squares with no horizontal whiskers (lines), as there are no confidence limits to be computed to denote variability across multiple trials. Of these, phenelzine, clomipramine, tranylcypromine and trazodone appear to be outliers. For each of 19 antidepressants, numerical values of efficacy are shown on the right (with confidence limits if there are more than 1 trials). The plot shows drugs listed by descending efficacy, with square size being proportional to the number of trials included (weighting). The vertical solid line represents line of no effect; vertical dotted line dissects the solid diamond that shows pooled RR for all agents tested. The pooled RR=1.42 indicates that antidepressants are 42% times more superior to placebos.

Correct Answer is : 1. Amitriptyline 2. Fluoxetine, Imipramine 3. Bupropion 4. None of them 5. Fluoxetine, Imipramine

Question 47

Not answered

Marked out of 4.00

Models and Tools for Quality Improvement

For each of the questions regarding quality improvement below, choose one or more appropriate answers as indicated from the list above.

1. An improvement system used to develop new processor products at superior performance levels. (Choose ONE)

2. An extension of PDSA model describing events and questions that must precede the initiation of a PDSA cycle (Choose TWO)

3. A visualisation tool to capture variations that occur within a system (Choose ONE)

Benefit Realisation Plan is a table used as a tool to ensure that the intended benefits originally planned in a QI project are actually delivered to the stakeholders in a timely fashion. The FOCUS approach to QI is shown in the figure below. FOCUS stands for Finding a process to improve, Organising a team, Clarifying current knowledge, Understanding causes of variation and Selecting the process improvement procedures. Model for improvement (MFI) is commonly employed now by many NHS organizations. MFI is an extension of PDSA and describes reflections that need to occur before embarking on a PDSA cycle. To measure outcomes over time, tools are available to generate time series graphs or frequency charts. Commonly used time series graphs include 'run charts' or Statistical Process Control charts (SPC). SPC enables the use of simple statistical concepts to understand processes that are being studied or implemented. In essence, SPC helps to assess the 'system' for predictability of outcome when a change is implemented; if unexpected variations are present, this provides an opportunity to study the external factors that may influence the desired outcome.

DMADV (Define-Measure-Analyze-Design-Validate) is a Six Sigma framework that focuses primarily on the development of a new service, product, or process as opposed to improving a previously existing one. It is especially useful when implementing new strategies and initiatives because of its basis in data, early identification of success, and thorough analysis. (Ref: <https://www.sixsigmadaily.com/what-is-dmadv/>)

Ans: 1. DMADV model

2. FOCUS approach, Model for Improvement (MFI)

3. Statistical Process Chart

Question **48**

Not answered

Marked out of 3.00

Types of prevention I

Identify the mode of prevention for each of the situations given below;

Woman with history of depression stopped venlafaxine after her dad had a heart attack

Choose...



ECG monitoring of patients on clozapine.

Choose...



Use of rivastigmine in dementia patients.

Choose...



Your answer is incorrect.

Explanation: 1) Indicated Primary Prevention. Given her familial risk of a cardiovascular event is increased, but has not yet happened to her, this is a form of Primary Intervention, particularly indicated for her. 2) Strictly speaking, ECG monitoring is not a preventative measure 3) Tertiary Prevention. This is indicated in cases of chronic illness, with the aim of slowing down progression.

The correct answer is: Woman with history of depression stopped venlafaxine after her dad had a heart attack → Indicated primary prevention., ECG monitoring of patients on clozapine. → Not a preventative measure, Use of rivastigmine in dementia patients. → Tertiary prevention



Question 49

Not answered

Marked out of 3.00

Types of prevention II

Identify the mode of prevention used in each of the following situations;

A 22-year-old female recently diagnosed with schizophrenia is started on an antipsychotic medication

Choose... ▾

The provision of substance abuse education as part of the school curriculum for all school children in England

Choose... ▾

A 45-year-old schizophrenia patient is transferred from an acute ward to a rehabilitation unit following treatment of a psychotic episode.

Choose... ▾

Your answer is incorrect.

Explanations: 1) Universal prevention strategies are designed to reach the entire population, without regard to individual risk factors and are intended to reach a very large audience. The program is provided to everyone in the population, such as a school or community. 2) Tertiary prevention focuses on reducing or minimizing the consequences of a disease once it has developed. The goal of tertiary prevention is to eliminate, or at least delay, the onset of complications and disability due to the disease. Most medical interventions fall into this category. 3) Secondary prevention attempts to identify a disease at its earliest stage so that prompt and appropriate management can be initiated. Successful secondary prevention reduces the impact of the disease

The correct answer is: A 22-year-old female recently diagnosed with schizophrenia is started on an antipsychotic medication → Secondary prevention, The provision of substance abuse education as part of the school curriculum for all school children in England → Universal prevention, A 45-year-old schizophrenia patient is transferred from an acute ward to a rehabilitation unit following treatment of a psychotic episode. → Tertiary prevention

Question 50

Not answered

Marked out of 5.00

Learning Disability Syndromes

Which is the most likely diagnosis for each of the given patients that have been referred to your LD clinic?

A 5-year-old boy with short stature and hypertelorism. His mother complains that he is hyperactive and very noisy.

Choose...

An 8-year-old boy who suffers from seizures and has recently developed mutism

Choose...

A short 12-year-old girl with a webbed neck, whose recent CT scan revealed a narrowed aorta

Choose...

A 14-year-old boy who has a history of social anxiety. In clinic he appears shy and avoids looking at your directly.

Choose...

A 9- year-old obese girl has a history of obsessional behaviour involving food that includes hyperphagia. She has a borderline IQ.

Choose...

Your answer is incorrect.

Explanation: 1) Cri-du-chat syndrome, also known as chromosome 5 depletion is a rare genetic disorder characterized by inappropriate laughter and a cat-like cry during infancy 2) Landau Kleffner Syndrome is an epilepsy syndrome of childhood. Its onset usually occurs before the age of six and is twice as likely to affect boys than girls. 3) Turner's Syndrome is a genetic disorder that only affects females causing distinctive characteristics such as short stature, webbed neck, widely spaced nipples and coarctation of the aorta. 4) Fragile X syndrome is an inherited cause of intellectual disability resulting in characteristics such as social anxiety, gaze avoidance and shyness. 5) Prader-Willi syndrome is a rare genetic disorder characterized by mild-to-moderate learning disability, food obsession, insatiable appetite, hyperphagia and obesity.

The correct answer is: A 5-year-old boy with short stature and hypertelorism. His mother complains that he is hyperactive and very noisy. → Cri-du-chat syndrome, An 8-year-old boy who suffers from seizures and has recently developed mutism → Landau Kleffner Syndrome, A short 12-year-old girl with a webbed neck, whose recent CT scan revealed a narrowed aorta → Turner's Syndrome, A 14-year-old boy who has a history of social anxiety. In clinic he appears shy and avoids looking at your directly. → Fragile X Syndrome, A 9- year-old obese girl has a history of obsessional behaviour involving food that includes hyperphagia. She has a borderline IQ. → Prader Willi Syndrome



Question 51

Not answered

Marked out of 5.00

Treatment of dementia

What would be the most appropriate treatment for?

A 74-year-old man with a diagnosis of Alzheimer's disease who presents with severe behavioural problems and is not responding to donepezil.

Choose...

A 78-year-old woman with a diagnosis of Alzheimer's disease who is taking donepezil and presents with a month's history of low mood, poor appetite, weight loss and insomnia. Her sister recently died from cancer.

Choose...

An 80-year-old man with a diagnosis of Parkinson's Disease who develops significant cognitive impairment

Choose...

A 92-year-old woman who has had a step-wise deterioration of cognition, which appears to coincide with historic strokes.

Choose...

A 73-year-old male on Sinemet therapy who experiences improvement in his Parkinsonian symptoms but within weeks complains of seeing goblins in the evening

Choose...

Your answer is incorrect.

Explanation: 1) Memantine, an NMDA-antagonist would be an appropriate choice for patients who are not responding to donepezil, an acetylcholinesterase inhibitor, and who present with challenging behaviour. It is important to check the renal function (including eGFR) of any patient prior to commencing memantine as it is eliminated by the kidneys and can cause renal impairment 2) This lady is presenting with symptoms of depression and would benefit from treatment with antidepressant medication. Mirtazapine is a popular choice for antidepressants in the elderly in view of its sedative effects (for those suffering from insomnia) and good side effect profile. 3) Rivastigmine, an acetylcholinesterase inhibitor, has been evaluated in the symptomatic treatment of patients with mild-to-moderate dementia associated with idiopathic Parkinson's disease. Although the evidence is preliminary, it is the best available evidence among the other options for Parkinson's related dementia. Rivastigmine appears to improve both cognition and activities of daily living in patients with PDD, resulting in a clinically meaningful benefit in a large number of cases. 4) This lady is suffering from vascular dementia. Anti-dementia drugs (acetylcholinesterase inhibitors) are not currently licensed for use in vascular dementia, although some studies suggest a possible benefit.

(Ref: SPMM). Aggressive treatment of risk factors remains the management of choice. 5) In patients with Parkinson's Disease who develop psychotic symptoms following treatment with dopamine agonists, use oral atypical antipsychotic medication, preferably ?quetiapine, (12.5 mg-75 mg but higher doses may be required) as they are associated with lower incidence of EPSEs. Furthermore, they do not require blood count surveillance. In refractory cases use clozapine in small doses as it is the most effective and only licensed antipsychotic in PD. Regular blood tests and close monitoring for neutropenia and agranulocytosis is essential in clozapine therapy.

The correct answer is: A 74-year-old man with a diagnosis of Alzheimer's disease who presents with severe behavioural problems and is not responding to donepezil. → Memantine, A 78-year-old woman with a diagnosis of Alzheimer's disease who is taking donepezil and presents with a month's history of low mood, poor appetite, weight loss and insomnia. Her sister recently died from cancer. → Mirtazapine, An 80-year-old man with a diagnosis of Parkinson's Disease who develops significant cognitive impairment → Rivastigmine, A 92-year-old woman who has had a step-wise deterioration of cognition, which appears to coincide with historic strokes. → Aggressive treatment of risk factors, A 73-year-old male on Sinemet therapy who experiences improvement in his Parkinsonian symptoms but within weeks complains of seeing goblins in the evening → Quetiapine



Question 52

Not answered

Marked out of 3.00

Psychotherapy - Types

What therapy could be offered in the following situations?

A 29 year old man with psychotic features who does not want antipsychotic medication

Choose...

A 36 year old man serving a 7 year sentence continually engages in repeated self harm

Choose...

A 23-year-old man murdered a friend in response to delusions. He has been remanded to hospital for treatment.

Choose...

Your answer is incorrect.

Explanation: 1) CBT is now used for the treatment of psychotic symptoms like delusions and hallucinations. There has been a particular emphasis on the factors maintaining the experience of psychotic symptoms like hallucinations and also on the distress & disturbance that can be associated with experiencing these symptoms. CBT strongly complements the recovery model for schizophrenia 2) DBT (alongside MBT- Mentalisation-Based Therapy) has a robust evidence base for effectiveness in people diagnosed with Personality Disorder (EUPD, Dissocial PD in particular) 3) Index Offense analysis involves the systematic gathering of reliable data on the characteristics of offenders and the offences which they have committed in order to develop understanding of the dynamics of offending and relevant intervention strategies. Index offence analysis should be a core task of any forensic clinician engaged in the assessment of offender/patients as it can provide a better understanding of crime scene actions and offence motivations. This can help guide treatment planning and improve risk assessments. (Ref: West AG, Greenall PV. Legal and criminological psychology. 2011 Feb 1;16(1):144-59.)

The correct answer is: A 29 year old man with psychotic features who does not want antipsychotic medication → Cognitive behavioural therapy for psychosis, A 36 year old man serving a 7 year sentence continually engages in repeated self harm → Dialectical behavioural therapy, A 23-year-old man murdered a friend in response to delusions. He has been remanded to hospital for treatment. → Index offence analysis.



© 2023 SPMM Course Limited, UK.

[Refund Policy](#) | [Terms & Conditions](#)

