

Started on	Monday, 29 November 2021, 1:17 PM
State	Finished
Completed on	Monday, 29 November 2021, 1:17 PM
Time taken	7 secs
Marks	0/10
Grade	0 out of 100

Question 1

Not answered

Marked out of 1

When sample size is same as population size, the type of survey carried out is often called

Select one:

- ☐ Epidemiological survey
- ☐ Cohort study
- ☐ Census survey
- ☐ Cross sectional survey
- ☐ Measurement survey

Your answer is incorrect.

Economic and practical limitations often make it difficult (even unethical) to collect measurements in an entire target population; hence we often resort to data from a small segment of a population for extrapolation to the target population. If the whole population is surveyed, it becomes a census survey.

The correct answer is: Census survey

Question 2

Not answered

Marked out of 1

A sample is selected in such a way that each group of the same size in the population has an equal chance of being selected; This type of sampling is known as

Select one:

- ☐ Stratified sampling
- ☐ Volunteer sampling
- ☐ Random sampling
- ☐ Convenience sampling
- ☐ Cluster sampling

Your answer is incorrect.

In a random sample, each individual in the parent population has an equal probability of being selected. One way of obtaining a random sample is to give each individual a number and then to use a computer-generated table of random numbers to decide which individuals should be included (SIMPLE RANDOM SAMPLE)

The correct answer is: Random sampling

Question 3

Not answered

Marked out of 1

A questionnaire concerning satisfaction with the Regional Mood Disorder Service was mailed to 50 patients selected from the patients register. The 50 patients in this survey are an example of a

Select one:

- ☐ Pilot group
- ☐ Parameter
- ☐ Sample
- ☐ Sampling frame
- ☐ Population

Your answer is incorrect.

A population represents the entire group of individuals in whom we are interested (all who had attended the mood disorder service). As it is costly and difficult to recruit all individuals in a population or a research project, we collect data on a sample of individuals who we believe are representative of this population. Hence, the study subjects must have similar characteristics to the individuals in the population.

The correct answer is: Sample

Question 4

Not answered

Marked out of 1

Netherlands mental health survey and incidence study (NEMESIS) used the following procedure to identify the sample for the survey. First, 90 municipalities were sampled randomly. Second, addresses of private households were selected randomly within those municipalities. Third, the person in the private household who had had the most recent birthday and was aged 18-64 years was asked to participate. Which of the following best describes the sampling method used?

Select one:

- ☐ Cluster sampling
- ☐ Multistage random sampling
- ☐ Simple random sampling
- ☐ Quota sampling
- ☐ Systematic sampling

Your answer is incorrect.

Multistage sampling is similar to cluster sampling except that there are two sampling events, instead of one. Initially, the larger primary units are randomly selected (e.g. municipalities in a country). Later individual units (households) within primary units are randomly selected for measurement. Note that in cluster sampling, all individuals in a selected primary unit are included in the study.

The correct answer is: Multistage random sampling

Question 5

Not answered

Marked out of 1

A case-control study investigating various risk factors for a disease in hospitalised patients chooses hospitalised patients with other diseases as controls. Which of the following is a major disadvantage of this process?

Select one:

- ☐ Higher cost of recruitment
- ☐ Baseline differences in socioeconomic and geographical residence
- ☐ Underestimation of true association
- ☐ Low availability of such controls
- ☐ Longer time duration required to recruit controls

Your answer is incorrect.

This approach has the advantage of selecting controls who come to the hospital through the same referral mechanisms as the cases, from the same region and are likely to be socioeconomically similar to the cases. The controls are readily available, and the control data is available at comparatively cheaper cost. But as the controls are diseased with a different diagnosis, they may not represent the entire population from which the cases arise. If the cause of the disease being studied is also associated with predictors of other diseases for which subjects get hospitalized, then the design may suffer from "overmatching," i.e. unintentionally the chosen controls will be more similar to the cases leading to a failure to detect or underestimate the strength of risk factors.

The correct answer is: Underestimation of true association

Question 6

Not answered

Marked out of 1

Which of the following is false concerning stratified random sampling?

Select one:

- ☐ Proportional stratified random sampling yields a representative sample
- ☐ It helps in addressing confounding effect
- ☐ Proportions of groups in the sample must always match their population proportions
- ☐ It involves a random selection process within identified subgroups
- ☐ All subjects within a subgroup have an equal chance of being selected

Your answer is incorrect.

Stratified sampling enables representation of not only the overall population, but also key subgroups of the population, especially small minority groups. We can use the same sampling fraction (proportion) for each stratum to produce a proportionate stratified random sampling. If a subgroup of interest is extremely small, we may also use different sampling fractions (f) within the different strata to randomly over-sample the small group. This is known as disproportionate stratified random sampling. Within each stratum, all individuals have an equal chance of being selected for the study.

The correct answer is: Proportions of groups in the sample must always match their population proportions

Question 7

Not answered

Marked out of 1

Which of the following method best represents the population from which a sample has been taken

Select one:

- ☐ A small sample picked on 'first come first serve basis' from a large number subgroups
- ☐ A large sample from small number of clusters
- ☐ A large sample based on simple random sampling
- ☐ A small sample based on simple random sampling
- ☐ A large sample based on the convenience sampling technique

Your answer is incorrect.

Large samples are generally more representative; convenient samples and quota samples may be less representative than simple random samples.

The correct answer is: A large sample based on simple random sampling

Question 8

Not answered

Marked out of 1

The type of sampling in which each member of the population selected for the sample is returned to the population before the next member is selected is called

Select one:

- ☐ Simple random sampling
- ☐ Systematic sampling
- ☐ Sampling without replacement
- ☐ Sampling with replacement
- ☐ Two stage sampling

Your answer is incorrect.

Suppose we generate a random list using a computer program to select a simple random sample. After we pick a number (unit) from the list, we can remove the number, or we can put it back into the list. If we put the number back in the list it may be selected more than once; if we put it aside, it can be selected only one time. "When a population element can be selected more than one time, we are sampling with replacement. When a population element can be selected only one time, we are sampling without replacement"(Retrieved from <http://stattrek.com/sampling/populations-and-samples.aspx>).

The correct answer is: Sampling with replacement

Question 9

Not answered

Marked out of 1

A cross-sectional epidemiological survey is carried out to determine the presence of quasi-psychotic symptoms in healthy working age adults. In this survey the target population is

Select one:

- ☐ All individuals on a list from which the sample was chosen
- ☐ All individuals that participated in the survey
- ☐ All individuals to whom the results should apply
- ☐ All individuals remaining in the list after the sample was chosen
- ☐ The total population in area surveyed

Your answer is incorrect.

The group to which we wish to generalize the results of a study is called the target population. The listing of the accessible population from which we draw the sample is called the sampling frame.

The correct answer is: All individuals to whom the results should apply

Question 10

Not answered

Marked out of 1

There were no new cases of CJD in the United Kingdom from January 1, 2007 through January 1, 2008. This measure refers to

Select one:

- ☐ Period prevalence
- ☐ Lifetime prevalence
- ☐ Incidence
- ☐ Lifetime expectancy
- ☐ Point prevalence

Your answer is incorrect.

Incidence is the number of new cases of a given disease at a specified time divided by the population at risk for that disease at that time.

The correct answer is: Incidence



© 2021 SPMM Course Limited, UK.

[Refund Policy](#) | [Terms & Conditions](#)

