Started on	Wednesday, 8 March 2023, 1:14 PM
State	Finished
Completed on	Wednesday, 8 March 2023, 1:18 PM
Time taken	4 mins 17 secs
Grade	**0.00** out of 150.00 (**0**%)

Question 1

Not answered

Marked out of 4.00

Defence mechanisms in operation

Identify the defence mechanism used in each of the following situations:

A boy who is being bullied at school takes on the identity of Superman.

A woman is not able to recall the abuse she sustained as a child.

An idle woman who is not performing well at her job is fired. She states this is because her colleague is bad-mouthing her rather than admitting her poor performance.

A man who was diagnosed to have cancer says, "This cannot be true... I follow a healthy lifestyle."

Sublimation
Reaction formation
Dissociation
Suppression
Intellectualization
Regression
Rationalisation
Repression
Denial

Choose...

Your answer is incorrect.

Explanation:

Reaction formation is the management of unacceptable impulses by permitting expression of the impulse in antithetical form.

Repression is characterised by preventing unacceptable aspects of internal reality from coming to conscious attention. (For example, a victim of childhood abuse may not have conscious awareness of the abuse as an adult.) The associated emotional reaction may remain in the conscious mind but divorced from its accompanying memory.

Rationalisation involves finding excuses that will justify unacceptable behaviours when self-esteem is threatened.

Denial is defined as the expressed refusal to recognise a threatening reality. It is of relevance especially to those with serious physical illnesses, where the patient denies being told of the presence of illness in themselves or their loved ones. It may persist despite constant reiteration of the facts.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1067.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 892.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. pp. 129-130.

The correct answer is: A boy who is being bullied at school takes on the identity of Superman. → Reaction formation, A woman is not able to recall the abuse she sustained as a child. → Repression, An idle woman who is not performing well at her job is fired. She states this is because her colleague is bad-mouthing her rather than admitting her poor performance. → Rationalization, A man who was diagnosed to have cancer says, "This cannot be true... I follow a healthy lifestyle." → Denial

Question 2

Not answered

Marked out of 3.00

Diagnosing neurotic disorders

Choose the correct diagnosis for each of the following clinical situations;

The medical teams have reviewed a woman several times over four years for symptoms of abdominal pain and shortness of breath. No organic cause has ever been found.

A man presents with low mood and anhedonia. He also has fatigue, poor sleep and headache.

A man presents with an abscess in his left groin. He requires constant reassurance from the nursing staff. A member of the night staff observes him injecting from a soiled syringe into the area adjacent to the abscess.

Normal adaptive reaction
Conversion disorder
Malingering
Bodily distress disorder
Factitious disorder
Post-traumatic stress disorder
Adjustment disorder

Your answer is incorrect.

Explanation:

Case 1: It is a case of bodily distress disorder, which is the experience of bodily symptoms that are distressing to the individual with excessive attention directed toward the symptoms.

Case 2: This man has core features of depression such as low mood, anhedonia and low energy levels.

Case 3: It is a case of factitious disorder in which a person deliberately produces, feigns, or exaggerates symptoms.

The correct answer is: The medical teams have reviewed a woman several times over four years for symptoms of abdominal pain and shortness of breath. No organic cause has ever been found. → Bodily distress disorder, A man presents with low mood and anhedonia. He also has fatigue, poor sleep and headache. → Moderate depressive episode, A man presents with an abscess in his left groin. He requires constant reassurance from the nursing staff. A member of the night staff observes him injecting from a soiled syringe into the area adjacent to the abscess. → Factitious disorder

Question 3

Not answered

Marked out of 3.00

Diagnosing culture-bound syndromes

For each of the descriptions given below chose one most appropriate culture-bound syndrome as a diagnosis from the list

A 32-year-old Greek woman presents with extreme sorrow, nausea, insomnia and fatigue Ch

A 44-year-old Mexican woman presents feeling despondent with involuntary tics and diarrhoea since her husband met with a sudden death in her presence Ch

A 22-year-old Nepalese man presents with the fear that he is losing semen in the urine. He describes symptoms suggestive of Urinary Tract infection along with tiredness. Ch

Amok
Dhat syndrome
Windigo
Tajin
Susto
Latah
Brain fag
Pa-leng
Koro
nerfiza

Your answer is incorrect.

Explanation:

Nerfiza (also called nervios or nevra) is a culture-related syndrome widespread throughout Latin America, and also found in Egypt and Greece, characterised by chronic anxiety and sorrowfulness, interpreted as a general state of vulnerability to psychosocial stressors brought about by personal or family difficulties, with a broad range of signs and symptoms, commonly including anorexia, dyssomnias, headaches, irritability, stomach upsets, tearfulness, loss of concentration, tremor, paraesthesia, and dizziness or vertigo.

Susto is a culture-related syndrome found in Peru and other parts of Latin America and among other Spanish-speaking communities. Following a frightening experience, the soul is believed to have departed the body, resulting in symptoms such as psychomotor agitation, anorexia, insomnia or hypersomnia, nightmares, and depressed mood, together with somatic complaints such as muscle and joint pain, headaches, and diarrhoea.

Dhat syndrome has been studied in South Asia and is rooted in Ayurvedic traditions about bodily production of semen as representing an end-product of energy-demanding metabolism. Dhat is recognised by a whitish discharge in the urine assumed to be semen. This 'loss of semen' is associated with somatic symptoms (weakness, exhaustion), severe anxiety, hypochondriasis, and sexual dysfunction.

Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 990.

The correct answer is: A 32-year-old Greek woman presents with extreme sorrow, nausea, insomnia and fatigue → Nerfiza, A 44-year-old Mexican woman presents feeling despondent with involuntary tics and diarrhoea since her husband met with a sudden death in her presence → Susto, A 22-year-old Nepalese man presents with the fear that he is losing semen in the urine. He describes symptoms suggestive of Urinary Tract infection along with tiredness. → Dhat syndrome

Question 4

Not answered

Marked out of 3.00

Supportive techniques

For each of the following examples, identify the type of supportive techniques aimed at eliciting information

Positive reinforcement
Reassurance
Disapproval
Statement of respect
Postponement
Validation
Parenting
Acknowledgement of affect
Advice

I note that these side effects are rather inconvenient for you. But you have shown some notable improvement, and if you continue with the treatment it is likely that you will achieve sustained response.

I can see that you look anxious and upset when talking about those voices

You have been through a lot. I am impressed at how you have hung in there and you must be a strong person

Your answer is incorrect.

Explanation: Reassurance: Used to instil positive hope and avoid or reduce despair. Must not be falsely reassuring. Acknowledgement of affect: Making a remark about patient's affect can facilitate disclosure.Statement of respect: Affirmative statements (must be genuine and appropriate) indicating respect and dignity along with positive reinforcement. Ref: Adapted from Kay J & Tasman A. Essentials of Psychiatry, 2nd edition, 2006. John Wiley & Sons, Ltd.

The correct answer is: I note that these side effects are rather inconvenient for you. But you have shown some notable improvement, and if you continue with the treatment it is likely that you will achieve sustained response. → Reassurance, I can see that you look anxious and upset when talking about those voices → Acknowledgment of affect, You have been through a lot. I am impressed at how you have hung in there and you must be a strong person → Statement of respect

Question 5

Not answered

Marked out of 5.00

Mok Assessment CXEMI038

Various forms of OCD and related disorders

A 35-year-old gentleman has been diagnosed with Obsessive Compulsive Disorder (OCD). He enquires about the various forms of this illness.

1. Which one symptom-based subtype confers a higher genetic risk to first-degree relatives?

2. Which two types predict a poor response to treatment with SSRI antidepressants?

3. Which two types predict a poor response to CBT?

Trichotilomania
Counting rituals
Aggressive/checking rituals
Excessive checking of facial defects
Sexual/religious obsessional thoughts
Hoarding rituals
Symmetry/ordering rituals
Contamination/wash rituals

Explanation: Symmetry/ordering rituals tend to be chronic and confers a higher genetic risk on first-degree relatives.Sexual/religious obsessional thoughts and hoarding rituals predict a poor response to SSRIs and CBT. Compulsive hoarding may be a neurobiologically distinct form of obsessive-compulsive disorder. Hoarding is notoriously difficult to treat by either psychological or pharmacological means.

The correct answer is:

Which one symptom domain confers a higher genetic risk to first-degree relatives? - Symmetry/ordering rituals,

Which TWO symptom domains predict a poor response to treatment with SSRI antidepressants? - Sexual/religious obsessional thoughts, Hoarding rituals,

Which TWO symptom domains predict a poor response to CBT? - Sexual/religious obsessional thoughts, Hoarding rituals

Question 6

Not answered

Marked out of 6.00

Ambitendency
Gegenhalten
Mitgehen
Cataplexy
Catalepsy
Waxy Flexibility
Automatic obidence
Stereotypy
Mannerism

Mok Assessment CXEMI039

Catatonic signs

From the above list chose the most appropriate terms describing each of the following clinical observations;

1. A patient moves his arm on slightest pressure regardless of any instruction given

2. Despite being asked to comply with, a patient resists with same pressure an attempt to move his arm. ✖

3. When asked to put out his hand, a patient extends, withdraws and then extends his hand repeatedly for 23 times before allowing it to rest on the examiner's hand. ✖

4. On hearing a sudden noise, a patient feels that his knees are buckling. He falls and hurts himself. ✖

5. A patient allows his arms to be manipulated freely without much resistance and maintains the posture in which the examiner leaves him for few minutes. (TWO) ✖ ✖

Explanation:

Mitmachen is a motor symptom of catatonia where the patient's limbs can be moved without resistance to any position. The limbs return to their resting state once the examiner lets go, in contrast with catalepsy where the limbs remain in their set positions for prolonged periods. Mitgehen is an extreme form of mitmachen where the patient's limbs can be moved to any position by very slight or fingertip pressure.

Gegenhalten is when an individual with catatonia opposes all passive movements with the same degree of force as that which is being applied by the examiner.

Ambitendency is a motor symptom of catatonia in which there is an alternating mixture of automatic obedience and negativism.

Cataplexy is a symptom of narcolepsy in which there is sudden loss of muscle tone, leading to collapse. This usually occurs following emotional stress.

Waxy flexibility (also known as catalepsy or flexibilitas cerea) is a rare motor symptom of catatonia that describes a situation in which the patient's limbs can be passively moved to any posture, which will then be held for a prolonged period of time.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 102, 104, 114.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 94.

The correct answer is:
A patient moves his arm on slightest pressure regardless of any instruction given - Mitgehen

Despite being asked to comply with, a patient resists with same pressure an attempt to move his arm - Gegenhalten

When asked to put out his hand, a patient extends, withdraws and then extends his hand repeatedly for 23 times before allowing it to rest on the examiner's hand - Ambitendency

On hearing a sudden noise, a patient feels that his knees are buckling. He falls and hurts himself - Cataplexy

A patient allows his arms to be manipulated freely without much resistance and maintains the posture in which the examiner leaves him for few minutes (TWO) - Catalepsy, Waxy flexibility

Question 7

Not answered

Marked out of 5.00

Drug Detection in Urine

How long could each of the following drugs be detected in urine?

Cannabis (chronic user)	Choose...
Cocaine (metabolites)	Choose...
Amphetamine	Choose...
Heroin	Choose...
Methadone (maintenance dosing)	Choose...

Up to 48 hours
Up to 30 days
2-4 days
1-3 days
60 days
Two weeks
90 days
Up to 7 days
Up to 12hrs

Your answer is incorrect.

Explanation:

Occasional use of cannabis could be detected in the urine up to 3 days. Heavy (daily) use could be detected in the urine up to 30 days. Cocaine could be found in urine for 6-8 hours; metabolites found up to 2-4 days. The half-life of methadone is longer than heroin and could be detected up to 7 days in the setting of maintenance dosing. Heroin could be detected for 1-3 days. Amphetamines could be detected up to 48 hours.

Ref: Taylor DM, Barnes TRE, Young AH. The Maudsley Prescribing Guidelines in Psychiatry, 14th Edition. 2021. p. 531-532.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 631.

The correct answer is: Cannabis (chronic user) → Up to 30 days, Cocaine (metabolites) → 2-4 days, Amphetamine → Up to 48 hours, Heroin → 1-3 days, Methadone (maintenance dosing) → Up to 7 days

Question 8

Not answered

Marked out of 3.00

Swapping antidepressants

For each of the following change of antidepressants, choose ONE method of switching from the options. Each option may be used once, more than once or not at all.

Venlafaxine to mirtazapine in a patient who has developed severe gastrointestinal intolerance.

Fluoxetine to tranylcypromine due to the inefficacy of the former.

Paroxetine to fluoxetine due to anticholinergic side effects of the former.

Withdraw the first and start the second immediately after stopping it.
Cross taper the two drugs
Offer combined therapy
Withdraw, wait for five weeks and then start the second drug
Keep full dose of the first drug and start the second drug at half dose
half full dose of the first drug and start the second drug at half dose
Withdraw, wait for 2 weeks and then start the second drug
Half the dose of the first drug and start the second drug at full dose
Withdrawal both the agents
Keep full dose of the first drug and start the second drug simultaneously

Your answer is incorrect.

Explanation:

When changing from one antidepressant to another, cross-tapering is preferred in which the ineffective or poorly tolerated drug is slowly reduced while the new drug is slowly introduced. This is generally recommended when switching from venlafaxine to mirtazapine. However, in this case the side effects are severe and intolerable, so a direct switch can be made.

Serotonin syndrome occurs more frequently with a combination of serotonergic drugs or in overdose. Most severe cases of serotonin syndrome involve an MAOI plus an SSRI. When switching from fluoxetine to phenelzine, stop fluoxetine and then wait for 5-6 weeks before starting the MAOI.

In some cases, cross-tapering may not be necessary. An example is when switching from one SSRI to another; their effects are so similar that administration of the second drug is likely to ameliorate withdrawal effects of the first. A direct switch is possible when switching from paroxetine to fluoxetine.

Ref: Taylor DM, Barnes TRE, Young AH. The Maudsley Prescribing Guidelines in Psychiatry, 14th Edition. 2021. p. 336-341.

The correct answer is: Venlafaxine to mirtazapine in a patient who has developed severe gastrointestinal intolerance. → Withdraw the first and start the second immediately after stopping it, Fluoxetine to tranylcypromine due to the inefficacy of the former. → Withdraw, wait for

five weeks and then start the second drug, Paroxetine to fluoxetine due to anticholinergic side effects of the former. → Withdraw the first and start the second immediately after stopping it

Question 9

Not answered

Marked out of 5.00

Inheritance of psychosis

A 26-year-old woman has been admitted to hospital for bizarre delusions, auditory hallucinations and disorganized speech. She is also flat in her affect, unkempt and is self-neglecting. There is no history of mania or depression. For each question below, choose the correct answer from the given list:

What is the chance of her 12-year-old son inheriting the disorder?	50%
	26-30%
	12-15%
She has a sister who has an adopted 6-year-old son with Down's syndrome. What is the chance of him developing the same psychotic disorder?	16-20%
	46%
What is the chance of her identical twin sister to inherit the disorder?	36-40%
	33%
If her child's father is also suffering from the same psychotic disorder, what is the risk of inheritance in the child?	0.5%
	21-25%
What is the risk of her non-identical twin brother to inherit the disorder?	40%
	6-10%

Your answer is incorrect.

Explanation:

Siblings and offspring of patients with schizophrenia have about a 10-fold increase in their lifetime risk of developing the disorder compared to the general population, whereas in second- degree relatives (e.g. nephews, nieces, grandchildren) the risk is about 3-4 times the population risk. People with Down's syndrome have lower than the average population risk for schizophrenia (choose the lowest number).

Family member(s) affected	Risk of schizophrenia (approximate) (%)
Identical twin	46
One sibling/fraternal twin	12-15
Both parents	40
One parent	12-15
One grandparent	6
No relatives affected	0.5-1

Adapted from Oxford Handbook of Psychiatry, 2019. p. 191.

SPMM COURSE

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 191.

The correct answer is: What is the chance of her 12-year-old son inheriting the disorder? → 12-15%, She has a sister who has an adopted 6-year-old son with Down's syndrome. What is the chance of him developing the same psychotic disorder? → 0.5%, What is the chance of her identical twin sister to inherit the disorder? → 46%, If her child's father is also suffering from the same psychotic disorder, what is the risk of inheritance in the child? → 40%, What is the risk of her non-identical twin brother to inherit the disorder? → 12-15%

Question 10

Not answered

Marked out of 3.00

Chromosomal profiles

Identify the karyotypic / genetic profile for each of the following conditions:

Mr. X is a 21-year-old tall, thin man with a moderate level of intellectual disability especially with reading. He has diabetes and presents with behavioural outbursts. Cho

Miss. Y is a 19-year-old woman with a moderate intellectual disability and has a developed a schizophrenia-like psychosis. She has cardiac problems and cleft palate. Cho

Mr. Z is an 11-year-old boy with growth retardation who is very sociable with markedly better verbal than spatial intelligence. He has cardiac problems and sensory-motor impairments. Cho

Deletion of part of chromosome 22
Trisomy 18
Deletion of part of chromosome 15
Deletion of part of chromosome 21
Deletion of part of chromosome 7
Triplication of part of chromosome 22
47XXY
46 YY
Deletion of part of chromosome 5

Your answer is incorrect.

Explanation:

Case 1 describes a man with Klinefelter syndrome. Klinefelter syndrome (KS) (47, XXY) is one of the most common chromosomal disorders and is a relatively frequent cause of male infertility and hypogonadism. Men with 47XXY makeup have an increased risk of diabetes and metabolic syndrome related to adiponectin dysregulation.

Case 2 is descriptive of Velocardio Facial Syndrome (VCFS). VCFS was first described by Kirkpatrick and DiGeorge in 1968 as a constellation of immunologic deficiencies (hence the name DiGeorge syndrome). It was renamed Shprintzen Syndrome following a thorough description of the common presenting symptoms by Shprintzen, et al., in 1978 who included palate anomalies (Velo), congenital cardiovascular defects (Cardio), and mild facial dysmorphism (Facial) as characteristic features of VCFS . Nearly 25% develop psychosis. It is associated with a microdeletion in the long arm of chromosome 22 at band 22q11.2.

Williams syndrome (case 3) is a rare neurodevelopmental disorder classically characterized by the triad of unusual facies, infantile hypercalcemia, and supravalvular aortic stenosis. Deletion of elastin or protein kinase gene at chromosome 7q11.23 is associated with this syndrome.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 808, 809, 817.

The correct answer is: Mr. X is a 21-year-old tall, thin man with a moderate level of intellectual disability especially with reading. He has diabetes and presents with behavioural outbursts. → 47 XXY, Miss. Y is a 19-year-old woman with a moderate intellectual disability and has a developed a schizophrenia-like psychosis. She has cardiac problems and cleft palate. → Deletion of part of chromosome 22, Mr. Z is an 11-year-old boy with growth retardation who is very sociable with markedly better verbal than spatial intelligence. He has cardiac problems and sensory-motor impairments. → Deletion of part of chromosome 7

Question 11

Not answered

Marked out of 5.00

Neurotransmitters & disorders

Identify the disorders associated with each of the neurotransmitters listed below

Substance P	Choose...
Neurotensin	Choose...
Neuropeptide Y	Choose...
Corticotropin-releasing factor	Choose...
Endocannabinoids	Choose...

Alzheimer's disease
Circadian function disorders
Schizophrenia
Migraine
Depression
OCD
Huntington's disease
Parkinson's disease
Feeding

Your answer is incorrect.

Substance P is the primary neurotransmitter in most primary afferent sensory neurons (Pain) and is also seen in the striatonigral pathway. Levels of substance P have been reported to be markedly reduced in patients with Huntington's chorea.

Neurotensin coexists with dopamine in some axon terminals. Neurotensin-related peptides are being investigated in the treatment of psychotic symptoms.

Neuropeptide Y may play an important role in feeding disorders through its effects on energy balance. It is secreted by the hypothalamus and in addition to increasing food intake, it increases the proportion of energy stored as fat and blocks nociceptive signals to the brain.

Corticotropin releasing factor (CRF) is the neurotransmitter associated with anxiety and depression, and CRF receptor antagonists are currently under clinical evaluation as anxiolytics and antidepressants.

The endocannabinoids are associated with schizophrenia and psychotic disorders. Anandamide (weak ligand) and 2-arachnidonylglycerol (strong) are formed from arachidonic acid and ethanolamine - belong to endocannabinoid family. The two types of cannabinoid receptors, central (CB1) and peripheral (CB2), bind tetrahydrocannabinol (THC), the active ingredient of marijuana. CB1 receptors are found in the brain and are potential targets for the treatment of both pain and psychosis. CB2 receptors are associated with immune regulation.

The correct answer is: Substance P → Huntington's disease, Neurotensin → Schizophrenia, Neuropeptide Y → Feeding, Corticotropin-releasing factor → Depression, Endocannabinoids → Schizophrenia

Question 12

Not answered

Marked out of 1.00

Tay-Sachs disease is caused by deficiency of

Select one:

- Neuropeptide Y
- Glucoronidase
- Monoamine oxidase
- Hexosaminidase A
- Hypoxanthine phosphoribosyl transferase

Your answer is incorrect.

Explanation:

Tay-Sachs disease belongs to the group of autosomal-recessive lysosomal storage metabolic disorders. It is caused by hexosaminidase A (Hex A) enzyme deficiency. In most cases, the disease manifests itself during infancy, which characterises the most severe disorders of the nervous system. The typical features are muscle weakness, ataxia, speech, and mental disorders. Currently disease treatment is based on symptom relief.

Ref: Solovyeva VV et al. New approaches to Tay-Sachs disease therapy. Frontiers in physiology. 2018:1663.

The correct answer is: Hexosaminidase A

Question 13

Not answered

Marked out of 1.00

Which of the following is a characteristic feature of autosomal recessive transmission?

Select one:

- Pure maternal transmission
- Everyone with mutant gene is diseased
- High male to male transmission
- Phenotype not expressed in females
- High risk with consanguinity

Your answer is incorrect.

As carriers are possible in AR transmission, 'hidden relatives' may give birth to unwell children.

The correct answer is: High risk with consanguinity

Question 14

Not answered

Marked out of 1.00

The anticipation in paternal transmission is most likely to show up in which of the following conditions?

Select one:

- Noonan syndrome
- Hurler syndrome
- Lesch-nyhan syndrome
- Huntington disease
- Tuberous sclerosis

Your answer is incorrect.

Explanation:

Huntington disease is characterised by the phenomenon of anticipation, where the age of onset tends to decrease in successive generations. This decrease is due to the unstable nature of the CAG repeats that tend to increase in size, particularly when passed through the male germline. Although germline instability can explain the phenomenon of anticipation, somatic instability has been proposed as a mechanism underlying the tissue specificity of the disease.

Ref: Jimenez-Sanchez M et al. Huntington's disease: mechanisms of pathogenesis and therapeutic strategies. Cold Spring Harbor perspectives in medicine. 2017 Jul 1;7(7):a024240.

The correct answer is: Huntington disease

Question 15

Not answered

Marked out of 1.00

If working memory deficit is inherited as a multifactorial trait then its recurrence risk will be

Select one:

- same in all races
- not dependent on number of affected family members
- influenced by the severity of deficit
- same in all families
- same in both sexes

Your answer is incorrect.

Involvement of more family members and higher severity of expression of a multifactorial trait may mean higher polygenic contribution to the inheritance. Ethnic and gender differences are well noted in multifactorial inheritances.

The correct answer is: influenced by the severity of deficit

Question 16

Not answered

Marked out of 1.00

Which of the following is most likely with regard to Fragile X syndrome?

Select one:

- Locus q25 is deleted
- Microcephaly is often seen
- Clinical features are apparent only during adolescence
- CTG repeats are notable
- Intelligence is correlated to the load of chromosomal damage

Your answer is incorrect.

Explanation:

Fragile X syndrome (FXS) is the most common single gene disorder, which causes autism and intellectual disability. The fragile X mental retardation 1 (FMR1) gene is silenced when CGG Triplet repeats exceed 200, which is the full mutation that causes FXS. Carriers of FXS have a CGG repeat of between 55 and 200, which is defined as a premutation and transcription of the gene is overactive with high levels of the FMR1 mRNA. Most carriers of the premutation have normal levels of fragile X mental retardation protein (FMRP) and a normal intelligence, but in the upper range of the premutation (120-200) the FMRP level may be lower than normal. The clinical problems associated with the premutation are caused by the RNA toxicity associated with increased FMR1mRNA levels, although for some mildly lowered FMRP can cause problems associated with FXS.

Ref: Tassanakijpanich N et al. Fragile X premutation and associated health conditions: A review. Clinical Genetics. 2021 Jun;99(6):751-60.

The correct answer is: Intelligence is correlated to the load of chromosomal damage

Question 17

Not answered

Marked out of 1.00

Which of the following refers to the stage between two successive cell divisions wherein the cell replicates its genetic material?

Select one:

- Metaphase
- Interphase
- Prophase
- Anaphase
- Telophase

Your answer is incorrect.

Interphase is the 'holding' stage

The correct answer is: Interphase

Question 18

Not answered

Marked out of 1.00

When experiencing disgust, which of the following parts of brain show consistent changes in functional imaging?

Select one:

- Anterior insular region
- Supplementary motor area
- Dorso medial thalamus
- Posterior hippocampus
- Anterior cingulate cortex

Your answer is incorrect.

Explanation:

The anterior insula is crucial in transforming unpleasant sensory input into visceromotor reactions and the accompanying feeling of disgust.

Cantone M et al. Fear and disgust: case report of two uncommon emotional disturbances evoked by visual disperceptions after a right temporal-insular stroke. BMC neurology. 2019 Dec;19(1):1-4.

The correct answer is: Anterior insular region

Question 19

Not answered

Marked out of 1.00

Which of the following is seen in anterior cerebral artery occlusion?

Select one:

- Contralateral lower limb weakness
- Sensory aphasia
- Unilateral sensory loss
- Ipsilateral facial palsy
- Bulbar palsy

Your answer is incorrect.

Occlusion of the anterior cerebral artery may result in weakness and sensory loss of the contralateral foot and leg; gait apraxia and urinary incontinence (with bilateral damage in the acute phase). This sometimes resembles the triad of Normal Pressure Hydrocephalus where identical anatomical structures are implicated albeit presenting with an insidious or subacute onset. In left-sided lesions, some cases present with transcortical motor aphasia.

The correct answer is: Contralateral lower limb weakness

Question 20

Not answered

Marked out of 1.00

A 56-year-old gentleman suffers from ataxia and nystagmus. His CT scan shows ischemia in cerebellum. Which part of cerebellum is most likely to be affected?

Select one:

- Vermis
- Anterior lobe
- Flocculo-nodular lobe
- Posterior lobe
- Lateral lobe

Your answer is incorrect.

Explanation:

The flocculonodular lobe is the evolutionarily oldest part of the cerebellum, consisting of a pair of irregular structures (flocculi) joined in the middle by a node or nodulus and separated by a deep fissure from the back of the cerebellum, connected to the vestibular system and concerned chiefly with posture, balance, and ballistic movements.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: Flocculo-nodular lobe

Question 21

Not answered

Marked out of 1.00

Which area of the brain is implicated with recognition of faces?

Select one:

- Amygdala
- Fusiform gyrus
- Nucleus accumbens
- Prefrontal cortex
- Left inferior frontal area

Your answer is incorrect.

Explanation:

The fusiform gyrus is a spindle-shaped gyrus on the medial (inside) surface of each temporal lobe, at the junction of the occipital lobe, occupied by Area V4 of the visual cortex, largely responsible for colour perception and also implicated in face recognition. Face perception is an ability that involves many areas of the brain. Brain imaging studies have typically shown a great deal of activity in the fusiform gyrus, an area also known to cause prosopagnosia when damaged bilaterally.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: Fusiform gyrus

Question 22

Not answered

Marked out of 1.00

Which of the following enzymes is responsible for the degradation of noradrenaline?

Select one:

- Acetylcholinesterase
- Choline Acetyl Transferase
- Catechol-O-Methyl transferase
- Tryptophan hydroxylase
- Tyrosine hydroxylase

Your answer is incorrect.

Explanation:

Norepinephrine (NE) action is terminated by two principal destructive or catabolic enzymes that turn NE into inactive metabolites. The first is monoamine oxidase A or B, which is located in mitochondria in the presynaptic neuron and elsewhere. The second is catechol-O-methyltransferase (COMT), which is thought to be located largely outside of the presynaptic nerve terminal. The action of NE can be terminated not only by enzymes that destroy NE, but also by a transport pump for NE that removes it from acting in the synapse without destroying it.

Ref: Stahl SM. Stahl's Essential Psychopharmacology: Neuroscientific Basis and Practical Applications, 5th Edition. 2021. pp. 253-254.

The correct answer is: Catechol-O-Methyl transferase

Question 23

Not answered

Marked out of 1.00

Which of the following substance is the most active ingredient in the Ethiopian plant Catha edulis, whose leaves on chewing provide amphetamine-like stimulation?

Select one:

- Coumarin
- Cathine
- Ketamine
- Ephedrine
- Cathinone

Your answer is incorrect.

Explanation:

Catha edulis, or 'Khat', is a plant indigenous to Eastern Africa and Southern Arabia. It releases a variety of psychoactive chemicals, including the stimulants cathinone and cathine. Khat is chewed like tobacco and held between the cheek and gums after chewing. The stimulant effect of the Khat plant was originally attributed to cathine, but this was disputed by reports showing the plant extracts from fresh leaves contained another substance more active than cathine - called cathinone. Cathinone is not very stable and breaks down to produce cathine and norephedrine. This explains why fresh leaves are more stimulating, and the habit of chewing Khat is more or less confined to the regions where it is cultivated.

In the mid-2000s, synthetic cathinones began to appear, marketed as 'legal high' chemicals. They were packaged under labels including 'screen cleaner', 'jewelry cleaner', and most notoriously 'bath salts'. These compounds are substituted cathinones that are chemically modified to avoid regulatory control. Generally, substituted cathinones increase synaptic catecholamine levels by inhibiting the dopamine, serotonin, and norepinephrine reuptake transporters in a fashion similar to cocaine. Most also facilitate catecholamine release, similar to methamphetamine.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 303.

The correct answer is: Cathinone

Question 24

Not answered

Marked out of 1.00

Hyperpolarization of the neuron is influenced by which of the following?

Select one:

- chloride and sodium
- calcium and chloride
- sodium and calcium
- potassium and sodium
- chloride and potassium

Your answer is incorrect.

In CNS neurons, inhibitory transmitters open chloride channels. In addition, second messengers that mediate inhibition work by operating on potassium channels.

The correct answer is: chloride and potassium

Question 25

Not answered

Marked out of 1.00

The receptors which have both pre and postsynaptic presence are

Select one:

- 5-HT2C receptors
- D2 receptors
- D1 receptors
- D5 receptors
- 5-HT1A receptors

Your answer is incorrect.

Explanation:

Dopamine receptors are divided into two groups. The first group is the D1-like receptors, including both D1 and D5 receptors. D1-like receptors are excitatory, and positively linked to adenylate cyclase. The second group is the D2-like receptors, including D2, D3, and D4 receptors. D2-like receptors are inhibitory and negatively linked to adenylate cyclase. Thus, the neurotransmitter DA can be either excitatory or inhibitory, depending on which DA receptor it binds. All five DA receptors can be located postsynaptically, but D2 and D3 receptors can also both be located presynaptically, where, due to their inhibitory actions, they act as autoreceptors to inhibit further DA release.

Ref: Stahl SM. Stahl's Essential Psychopharmacology: Neuroscientific Basis and Practical Applications, 5th Edition. 2021. pp. 81-82.

The correct answer is: D2 receptors

Question 26

Not answered

Marked out of 1.00

Which of the following is the most powerful endogenous opioid peptide?

Select one:

- ß-Endorphin
- Morphine
- Dynorphin
- Enkephalin (?-endorphin)
- Neoendorphin

Your answer is incorrect.

Explanation:

Dynorphin A has been described as ‘extraordinarily potent’ (‘dyn from the Greek, dynamis = power and ‘orphin’ for the endogenous morphine peptide). Dynorphin peptides are released in the brain during stress exposure. After they are released, they activate kappa opioid receptors distributed throughout the brain and spinal cord, where they trigger cellular responses resulting in different stress responses: analgesia, dysphoria-like behaviours, anxiety-like responses, and increased addiction behaviours in experimental animals.

Ref: Chavkin C. Dynorphin–still an extraordinarily potent opioid peptide. Molecular pharmacology. 2013 Apr 1;83(4):729-36.

The correct answer is: Dynorphin

Question 27

Not answered

Marked out of 1.00

All of the following intrauterine infections cause intellectual disabilities except

Select one:

- CMV
- Toxoplasmosis
- Treponema pallidum
- Rubella
- Epstein Barr Virus

Your answer is incorrect.

Explanation:

Maternal infections during pregnancy, especially viral infections, have been known to cause fetal damage and intellectual disability. Syphilis (caused by Treponema pallidum) in pregnant women was once the leading cause of various neuropathic changes in their offspring, including intellectual disability. Rubella (German measles) has now replaced syphilis as the primary cause of congenital malformations and intellectual disability caused by maternal infection. Children with intellectual disability from cytomegalovirus frequently have cerebral calcification, microcephaly, or hydrocephalus. Toxoplasmosis causes mild to severe intellectual disability and, in severe cases, hydrocephalus, seizures, microcephaly, and chorioretinitis. EBV (Infectious Mononucleosis) has not been implicated in intellectual disability syndromes.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 104.

The correct answer is: Epstein Barr Virus

Question 28

Not answered

Marked out of 1.00

Which of the following is least likely to occur in schizophrenia?

Select one:

- Increase in neuron density
- Cytoarchitectural anomalies
- Reduced cell size affecting neurons
- Gliosis
- Reduced cell number affecting neurones

Your answer is incorrect.

Explanation:

Macroscopic brain changes that occur in schizophrenia include enlarged ventricle size and reductions in grey matter volume, whole-brain volume, and white matter anisotropy. The reduced neuropil (intertwined axons and dendrites of neurons) hypothesis has been suggested as a microanatomical explanation to account for these macroscopic changes. Studies have found both increased and decreased neuronal density in the cerebral cortex. The literature provides support for increased neuron density in schizophrenia, with variation dependent on age. Although the brain is abnormal, gliosis is absent.

Ref: Bakhshi K, Chance SA. The neuropathology of schizophrenia: a selective review of past studies and emerging themes in brain structure and cytoarchitecture. Neuroscience. 2015 Sep 10;303:82-102.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 189.

The correct answer is: Gliosis

Question 29

Not answered

Marked out of 1.00

An intravenous heroin user presents with significant slowing of cognitive ability. He has fatuous affect and impaired concentration that has developed insidiously. Which of the following is a likely diagnosis?

Select one:

- Opioid intoxication
- Bacterial encephalitis
- Heroin withdrawal
- HIV-associated neurocognitive disorder
- Depression

Your answer is incorrect.

Explanation:

HIV-associated neurocognitive disorder (HAND) is an umbrella term for the spectrum of neurocognitive impairment induced by HIV (direct effects of HIV). It ranges from asymptomatic impairment to severe impairment with pronounced deficits in self-efficacy and interference with day-to-day activities. Early indicators of HAND include impairments in psychomotor function, in simple and divided attention and concentration, in mental agility, and in mental flexibility. HAND also relates to personality changes (e.g. apathy, irritability, and inertia) and changes in behaviour (e.g. social withdrawal and impaired activities of daily living). Neurological signs of HAND include poor concentration, loss of balance, and tremor.

Ref: Knights MJ et al. HIV infection and its psychiatric manifestations: a clinical overview. BJPsych Advances. 2017 Jul;23(4):265-77

Taylor DM, Gaughran F, Pillinger T. The Maudsley Practice Guidelines for Physical Health Conditions in Psychiatry. 2021.

The correct answer is: HIV-associated neurocognitive disorder

Question 30

Not answered

Marked out of 1.00

The term used to describe EEG more than 13Hz is

Select one:

- Mu waves
- Beta waves
- Gamma waves
- Theta waves
- Alpha waves

Your answer is incorrect.

Explanation: Beta waves have a frequency of more than 13 Hz.

EEG Rhythm	Frequency range	Amplitude	Main scalp area	Subject condition
Delta	0.1-3.5 Hz	50-350 uV	Variable	Drowsiness and sleep Hyperventilation Infancy and childhood
Theta	4-7.5 Hz	10-150 uV	Variable	Drowsiness and sleep Hyperventilation Infancy and childhood Small amount in awake adults
Alpha	8-13 Hz	20-100 uV	Posterior (occipital-parietal)	Relaxed wakefulness with eyes closed
Beta	Above 13 Hz	10-30 uV	Frontal or diffuse	Increase during cognitive efforts as well as during drowsiness and sleep
Mu	7-12 Hz	10-50 uV	Central	Relaxed wakefulness with both eyes open and closed Blocked by movements of the contralateral body parts
Lambda	Sharp transients of 200-300 msec duration	Below 50 uV	Occipital	Visual exploration

Adapted from Standard electroencephalography in clinical psychiatry: A practical handbook. 2011.

SPMM COURSE

Ref: Boutros NN, Galderisi S, Pogarell O, Riggio S. Standard electroencephalography in clinical psychiatry: A practical handbook. 2011.

The correct answer is: Beta waves

Question 31

Not answered

Marked out of 1.00

The disorder associated with ambiguous genitalia is

Select one:

- Cri du chat syndrome
- Cushing syndrome
- Conn syndrome
- Congenital adrenal hyperplasia
- Cotard syndrome

Your answer is incorrect.

Explanation:

The congenital adrenal hyperplasias comprise a family of autosomal recessive disorders that disrupt adrenal steroidogenesis. Infant females with classical congenital adrenal hyperplasia generally present in the neonatal period with ambiguous genitalia.

Ref: Witchel SF. Congenital adrenal hyperplasia. Journal of pediatric and adolescent gynecology. 2017 Oct 1;30(5):520-34.

The correct answer is: Congenital adrenal hyperplasia

Question 32

Not answered

Marked out of 1.00

A 58 year old man presents with postural hypotension, anorexia, nausea and weakness. He appears to be extremely apathetic, with significant anhedonia compounded by fatigue and depressed mood.Which of the following hormonal imbalance is most likely?

Select one:

- Vasopressin
- Insulin
- Thyroxine
- Corticosteroids
- Adrenaline

Your answer is incorrect.

Explanation:

Addison syndrome may result from adrenal destruction by autoimmune disease, metastatic cancer, tuberculosis, or human immunodeficiency virus infection or secondary to suppression of ACTH secretion by chronic corticosteroid therapy. Addison disease (primary adrenal insufficiency) is characterised by chronic glucocorticoid and/or mineralocorticoid deficiency due to failure of the adrenal cortex. It should be suspected in all acutely or chronically ill patients with general fatigue or severe weakness and unexplained dehydration, hypotension, weight loss, fever, abdominal pain, and hyperpigmentation. Psychiatric symptoms include apathy, anhedonia, fatigue, and depressed mood. It may be misdiagnosed as major depressive disorder. Anorexia is common in both depression and Addison disease, but the presence of nausea, vomiting and hyperpigmentation may suggest Addison disease.

Ref: Barthel A et al. An update on Addison's disease. Experimental and Clinical Endocrinology & Diabetes. 2019 Feb;127(02/03):165-75.

The correct answer is: Corticosteroids

Question 33

Not answered

Marked out of 1.00

Choose the antidepressant which, if given to a pregnant mother, is most likely to cause withdrawal symptoms in the neonate?

Select one:

- Citalopram
- Paroxetine
- Fluoxetine
- Clomipramine
- Sertraline

Your answer is incorrect.

Explanation:

When antidepressants are taken during pregnancy, the neonate may experience discontinuation symptoms, which are usually mild, such as agitation and irritability, or rarely respiratory distress and convulsions (with SSRIs). The risk is assumed to be particularly high with short half-life drugs such as paroxetine and venlafaxine. Continuing to breastfeed and then 'weaning' by switching to mixed (breast/bottle) feeding may help reduce the severity of reactions.

Ref: Taylor DM, Barnes TRE, Young AH. The Maudsley Prescribing Guidelines in Psychiatry, 14th Edition. 2021. p. 689.

The correct answer is: Paroxetine

Question 34

Not answered

Marked out of 1.00

The drug that is most likely to result in a hypertensive crisis from the given list is

Select one:

- Trifluoperazine
- Tryptophan
- Venlafaxine
- Duloxetine
- Phenelzine

Your answer is incorrect.

Explanation:

The most worrisome side effect of MAOIs (such as phenelzine) is the tyramine-induced hypertensive crisis ('cheese reaction'). The amino acid tyramine is normally transformed via GI metabolism. However, MAOIs inactivate GI metabolism of dietary tyramine, thus allowing intact tyramine to enter the circulation. A hypertensive crisis may subsequently occur because of a powerful pressor effect of the amino acid.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 645.

The correct answer is: Phenelzine

Question 35

Not answered

Marked out of 1.00

Lucy on antidepressant medication for more than 8 years. She has developed pins and needles and a burning type of pain on her feet and legs. The most likely offending medication is

Select one:

- Imipramine
- Fluoxetine
- Trazodone
- Phenelzine
- Fluvoxamine

Your answer is incorrect.

Explanation:

Among the most common drugs that can cause toxic neuropathies are phenelzine and phenytoin. Phenelzine has a higher likelihood of causing peripheral neuropathy than the other antidepressants listed here.

Ref: Grisold W, Carozzi VA. Toxicity in Peripheral Nerves: An Overview. Toxics. 2021 Sep;9(9):218.

The correct answer is: Phenelzine

Question 36

Not answered

Marked out of 1.00

A 50-year-old lady on treatment with Tranylcypromine consumed cheese and developed hypertensive reaction. What is the most suitable choice for treatment of this condition?

Select one:

- Captopril
- Flumazenil
- Propranalol
- Phentolamine
- Hemodialysis

Your answer is incorrect.

Explanation:

An MAOI-induced hypertensive crisis should be treated with alpha-adrenergic antagonists (for example phentolamine or chlorpromazine). These drugs lower blood pressure within 5 minutes. IV furosemide can be used to reduce fluid load, and a beta-adrenergic receptor antagonist can control tachycardia.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 646.

The correct answer is: Phentolamine

Question 37

Not answered

Marked out of 1.00

Mrs. X is 30 years old a patient with bipolar affective disorder with secondary amenorrhea. The investigation to be done is

Select one:

- Serum prolactin levels
- Pregnancy test
- FSH and LH levels
- Thyroid levels
- Serum testosterone levels

Your answer is incorrect.

Pregnancy is an obvious cause of amenorrhea and is the most common reason for secondary amenorrhea.

The correct answer is: Pregnancy test

Question 38

Not answered

Marked out of 1.00

The best treatment among the following for lithium induced tremors is

Select one:

- Procyclidine
- Levo-dopa
- Propranalol
- Selegiline
- Amantadine

Your answer is incorrect.

Explanation:

Early adverse effects of lithium include gastrointestinal upset and fine tremor. Propranolol can be useful in lithium-induced tremor. Course tremor is associated with lithium toxicity and may be seen together with muscle weakness/twitching, drowsiness, confusion, and ataxia.

Ref: Taylor DM, Barnes TRE, Young AH. The Maudsley Prescribing Guidelines in Psychiatry, 14th Edition. 2021. p. 250.

The correct answer is: Propranalol

Question 39

Not answered

Marked out of 1.00

Which of the following is not a significant risk factor for the development of antidepressant induced SIADH?

Select one:

- Increasing age
- Polypharmacy
- Medical co-morbidity
- Female sex
- Psychotic symptoms

Your answer is incorrect.

Explanation:

Most antidepressants have been associated with hyponatraemia. It has been suggested that serotonergic drugs are relatively more likely than noradrenergic drugs to cause hypnonatraemia, although this is disputed. Risk factors include: older age (most important), female sex, major surgery, history of hyponatraemia or a low baseline sodium concentration, co-therapy with other drugs known to be associated hyponatraemia (e.g. diuretics, NSAIDs, antipsychotics, carbamazepine, cancer chemotherapy, calcium antagonists, ACE inhibitors, and laxatives), reduced renal function, medical co-morbidity (e.g. hypothyroidism, diabetes, COPD, hypertension, head injury, congestive cardiac failure, cerebrovascular accident, various cancers), and low body weight.

Ref: Taylor DM, Barnes TRE, Young AH. The Maudsley Prescribing Guidelines in Psychiatry, 14th Edition. 2021. pp. 393-394.

The correct answer is: Psychotic symptoms

Question 40

Not answered

Marked out of 1.00

Which of the following has been associated with the use of SSRIs in mid-late pregnancy?

Select one:

- Cleft palate
- Hypospadias
- Pulmonary hypertension
- Pyloric stenosis
- Neural tube defects

Your answer is incorrect.

Explanation:

A specific problem that has been associated with serotonergic antidepressant use in mid to late pregnancy is persistent pulmonary hypertension of the newborn (PPHN). Although initial reports suggested an approximate six-fold increase, recent literature has shown the absolute risk to be small and more modest. Although the risk of PPHN in babies exposed in utero to SSRIs is increased, the absolute risk is low.

Ref: McAllister-Williams RH et al. British Association for Psychopharmacology consensus guidance on the use of psychotropic medication preconception, in pregnancy and postpartum 2017. Journal of Psychopharmacology. 2017 May;31(5):519-52.

The correct answer is: Pulmonary hypertension

Question 41

Not answered

Marked out of 1.00

What effect can carbamazepine have on cardiac function?

Select one:

- Exacerbation of pre-existing cardiac disease
- Hypersensitivity myocarditis
- Reduction of cardiac conduction
- Sinus tachycardia
- All of the listed options

Your answer is incorrect.

Explanation:

Carbamazepine decreases cardiac conduction and can thus exacerbate pre-existing cardiac disease. Carbamazepine use rarely elicits an immune hypersensitivity response consisting of fever, rash, eosinophilia, and possibly fatal myocarditis. Carbamazepine can cause sinus tachycardia.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 661.

Taylor DM, Gaughran F, Pillinger T. The Maudsley Practice Guidelines for Physical Health Conditions in Psychiatry. 2021. p. 5.

The correct answer is: All of the listed options

Question 42

Not answered

Marked out of 1.00

Deaths in neuroleptic malignant syndrome (NMS) are usually due to

Select one:

- Renal failure
- Respiratory failure
- Disseminated intravascular coagulation
- Arrhythmias
- All of the listed options

Your answer is incorrect.

Deaths from neuroleptic malignant syndrome are usually due to respiratory failure, cardiovascular collapse, myoglobinuric renal failure, sepsis, arrythmias, thromboembolism, or disseminated intravascular coagulation (DIC).

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 1018.

The correct answer is: All of the listed options

Question 43

Not answered

Marked out of 1.00

Which one among the following is more likely to induce akathisia?

Select one:

- Risperidone
- Quetiapine
- Amisulpride
- Olanzapine
- Clozapine

Your answer is incorrect.

Explanation:

The relative liability of individual antipsychotic medications for akathisia is uncertain, but there is consensus that the incidence is lowest for olanzapine, quetiapine, and clozapine. The incidence of akathisia is relatively high with haloperidol, risperidone, aripiprazole, and ziprasidone.

Ref: Taylor DM, Barnes TRE, Young AH. The Maudsley Prescribing Guidelines in Psychiatry, 14th Edition. 2021. pp. 110, 114.

The correct answer is: Risperidone

Question 44

Not answered

Marked out of 1.00

Mr.X is a 33 year old gentleman with a long standing diagnosis of paranoid schizophrenia. He was started on atypical anti psychotic medication 6 months ago and when reviewed in the clinic, he complains of side effects such as reduced sexual function and has also developed enlarged breasts. Which of the following anti psychotic drugs is most likely to be associated with these side effects?


Select one:

- Clozapine
- Aripiprazole
- Quetiapine
- Risperidone
- Olanzapine

Your answer is incorrect.

Explanation:

When D2 antagonists are administered, D2 receptors in the tuberoinfundibular dopamine pathway are blocked. This causes plasma prolactin concentrations to rise, a condition called hyperprolactinaemia. This can be associated with galactorrhea (i.e. breast secretions), sexual dysfunction, gynaecomastia (enlargement of the breasts) in men, and amenorrhea (i.e. irregular or lack of menstrual periods) in women.



Ref: Stahl SM. Stahl's Essential Psychopharmacology: Neuroscientific Basis and Practical Applications, 5th Edition. 2021. pp. 164-165.

Taylor DM, Barnes TRE, Young AH. The Maudsley Prescribing Guidelines in Psychiatry, 14th Edition. 2021. p. 168.

The correct answer is: Risperidone

Question 45

Not answered

Marked out of 1.00

The site of action of thyroid hormones is at the level of;

Select one:

- Nucleus
- Cell membrane
- Plasma membrane
- Cytoplasm
- None of the listed options

Your answer is incorrect.

Thyroid hormones diffuse through cell membranes to bind with receptors that are members of a large super-family of nuclear receptors that act as hormonally-activated transcription factors. The hormone-receptor complex interacts with the DNA of to activate gene transcription processes downstream.

The correct answer is: Nucleus

Question 46

Not answered

Marked out of 1.00

Zolpidem acts as an agonist at

Select one:

- Omega 1 receptor
- Omega 3 receptor
- Alpha receptor
- Beta receptor
- Omega 2 receptor

Your answer is incorrect.

Explanation:

Z drugs bind to benzodiazepine-sensitive GABA-A receptors. A different nomenclature (omega system) denotes the specific sites of GABA-A receptors where these 'Z' drugs act as omega-1 receptors. Conventional benzodiazepines interact at both ω1 and ω2 sites, the latter likely contributing to side effects.

Zolpidem and zaleplon bind selectively to alpha-1 subunits of these sites. By contrast, benzodiazepines, zopiclone and eszopiclone bind to four alpha subunits (alpha-1, alpha-2, alpha-3, and alpha-5). The functional significance of alpha-1 selectivity is not yet proven, but may contribute to lower risk of tolerance and dependence. The alpha-1 subtype is known to be critical for producing sedation and thus is targeted by every effective GABA-A PAM (positive allosteric modulator) hypnotic, both benzodiazepines and Z-drugs.

Ref: Stahl SM. Stahl's Essential Psychopharmacology: Neuroscientific Basis and Practical Applications, 5th Edition. 2021. pp. 422-423.

Madari S, Golebiowski R, Mansukhani MP, Kolla BP. Pharmacological management of insomnia. Neurotherapeutics. 2021 Jan;18(1):44-52.

The correct answer is: Omega 1 receptor

Question 47

Not answered

Marked out of 1.00

Which of the following medications is not associated with an increased risk of depression?

Select one:

- Varenicline
- Alpha interferons
- Ezetimibe
- Orlistat
- Isoretinoin

Your answer is incorrect.

Explanation:

Orlistat interferes with the absorption of dietary fats, causing reduced caloric intake. It works by inhibiting gastric and pancreatic lipases, the enzymes that break down triglycerides in the intestine. The most common subjective side effects of orlistat are GI related and include steatorrhea, flatulence, fecal incontinence, and frequent or urgent bowel movements. Hepatic and renal injuries are potentially severe side effects or orlistat use. It has not been associated with an increased risk of depression.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. pp. 708-709.

The correct answer is: Orlistat

Question 48

Not answered

Marked out of 1.00

Which of the following selective serotonin reuptake inhibitors has notable anticholinergic activity?

Select one:

- Escitalopram
- Paroxetine
- Fluoxetine
- Citalopram
- Fluvoxamine

Your answer is incorrect.

Explanation:

Paroxetine has the highest anticholinergic effects of the SSRIs. Paroxetine's mild anticholinergic actions can enhance the rapid onset of anxiolytic and hypnotic efficacy, but also cause mild anticholinergic side effects.

Ref: Taylor DM, Barnes TRE, Young AH. The Maudsley Prescribing Guidelines in Psychiatry, 14th Edition. 2021. pp. 421-422.

Stahl SM. Stahl's Essential Psychopharmacology: Prescriber's Guide, 7th Edition. 2021. p. 599.

The correct answer is: Paroxetine

Question 49

Not answered

Marked out of 1.00

Risperidone increases the levels of

Select one:

- Prolactin
- Growth hormone
- LH
- FSH
- Neuropeptides

Your answer is incorrect.

Explanation:

When D2 antagonists, such as risperidone, are administered, D2 receptors in the tuberoinfundibular dopamine pathway are blocked. This causes plasma prolactin concentrations to rise, a condition called hyperprolactinaemia. This can be associated with galactorrhea (i.e. breast secretions), sexual dysfunction, gynaecomastia (enlargement of the breasts) in men, and amenorrhea (i.e. irregular or lack of menstrual periods) in women.

Ref: Stahl SM. Stahl's Essential Psychopharmacology: Neuroscientific Basis and Practical Applications, 5th Edition. 2021. pp. 164-165.

The correct answer is: Prolactin

Question 50

Not answered

Marked out of 1.00

Which combination could be prescribed together, in which the second drug is used to minimize the adverse effect of the first one?

Select one:

- Clozapine and tranylcypromine
- Reboxetine and Doxazosin
- Citalopram and carbamazepine
- Chlorpromazine and diazepam
- Bupropion and Fluoxetine

Your answer is incorrect.

Explanation:

Reboxetine is a selective norepinephrine reuptake inhibitor. Side effects may appear 'anticholinergic', but reboxetine does not directly block muscarinic receptors. Constipation, dry mouth, and urinary retention are noradrenergic, due in part to peripheral alpha 1 receptor stimulation causing decreased acetylcholine release. Thus, antidotes for these side effects can be alpha 1 antagonists such as tamsulosin or doxazosin, especially in men over 50 with borderline urine flow.

Ref: Stahl SM. Stahl's Essential Psychopharmacology: Prescriber's Guide, 7th Edition. 2021. p. 684.

The correct answer is: Reboxetine and Doxazosin

Question 51

Not answered

Marked out of 1.00

What happens to GABA-A receptors in benzodiazepine withdrawal state?

Select one:

- Reduced glutamate transmission
- Reduced activity at GABA receptors
- Increased dopamine levels
- Sensitisation of GABA receptor system
- Reduced inhibitory tone in the CNS

Your answer is incorrect.

Chronic benzodiazepine administration may lead to uncoupling of the linkage between GABAA and benzodiazepine receptor sites and subsequent preferential degradation of certain GABAA receptor units. Such a change results in a long-term homeostatic adaptation to the chronic presence of benzodiazepines leading to a reduction in GABA receptor density. Sudden withdrawal of the benzodiazepine once tolerance has developed would expose the recipient to underactive GABA system without the counterbalancing presence of benzodiazepine drugs.

The correct answer is: Reduced activity at GABA receptors

Question 52

Not answered

Marked out of 1.00

A middle-aged gentleman suffers from alcohol withdrawal symptoms and has history of liver impairment. A suitable drug for detoxification would be;

Select one:

- Diazepam
- Oxazepam
- Carbamazepine
- Temazepam
- Chlordiazepoxide

Your answer is incorrect.

Explanation:

Shorter-acting benzodiazepines (e.g. oxazepam) may be preferred in those for whom over-sedation must be avoided, in people with liver disease who may not be able to metabolise long-acting agents efficiently, and in people with COPD. However, there is a greater risk of breakthrough seizures and greater potential for misuse.

Ref: National Institute for Health and Clinical Excellence. The NICE guideline on diagnosis, assessment and management of harmful drinking and alcohol dependence. Updated edition 2020.

The correct answer is: Oxazepam

Question 53

Not answered

Marked out of 1.00

Which one among the following is more related to causing discontinuation syndrome?

Select one:

- Citalopram
- Fluvoxamine
- Fluoxetine
- Sertraline
- Paroxetine

Your answer is incorrect.

Explanation:

Discontinuation effects can be more common or more severe with paroxetine than with other SSRIs. Paroxetine's discontinuation effects may be related to the fact that it inhibits its own metabolism. Thus, when paroxetine is withdrawn, the rate of its decline can be faster as it stops inhibiting its metabolism. Readaptation of cholinergic receptors after prolonged blockade may also contribute to the discontinuation effects of paroxetine.

Ref: Stahl SM. Stahl's Essential Psychopharmacology: Prescriber's Guide, 7th Edition. 2021. p. 596.

The correct answer is: Paroxetine

Question 54

Not answered

Marked out of 1.00

Which of the following drugs follows zero-order kinetics?

Select one:

- Carbamazepine
- Topiramate
- Lamotrigine
- Phenytoin
- Sodium valproate

Your answer is incorrect.

Explanation:

Phenytoin follows zero-order kinetics.

Ref: Taylor DM, Barnes TRE, Young AH. The Maudsley Prescribing Guidelines in Psychiatry, 14th Edition. 2021. p. 836.

The correct answer is: Phenytoin

Question 55

Not answered

Marked out of 1.00

Which of the following is correct with regard to the pharmacokinetics of depot antipsychotics?

Select one:

- Test doses are not useful to avoid severe adverse effects
- Higher doses are significantly more effective than lower doses
- Adverse effects are usually short lived than oral medications due to the low frequency of administration
- Plasma level of depot antipsychotics could be as high as 4 times the initial levels, 2 to 3 months after starting treatment
- Shortening the dosing intervals can improve efficacy in non-responders

Your answer is incorrect.

Explanation:

For first-generation antipsychotic (FGA) depots, a test dose should be given. Because of its long half-life, any adverse effects from the administration of a depot are likely to be long-lived. For FGAs, a test dose consisting of a small dose of active drug in a small volume of oil serves a dual purpose: it is a test of the patient's sensitivity to EPS and of any sensitivity to the base oil. For second-generation antipsychotic (SGA) depots, test doses may not be required (less propensity to cause EPS and aqueous base not known to be allergenic) although they could be considered appropriate where a patient is suspected of being non-adherent to oral antipsychotic medication and the depot will be the first exposure to guaranteed antipsychotic medication delivery. For both types of depots, prior treatment with the equivalent oral formulation is preferred to assess efficacy and tolerability. At the start of therapy, plasma drug levels increase over several weeks to months without increasing the given dose. (This is due to accumulation: steady state is only achieved after at least 6-8 weeks.) There are few data showing clear dose-response effects for depots. There is some information indicating that low doses (within the licensed range) may be at least as effective as higher ones. Depots should be administered at the longest possible licensed interval. There is no evidence to suggest that shortening the dose interval improves efficacy.

Ref: Taylor DM, Barnes TRE, Young AH. The Maudsley Prescribing Guidelines in Psychiatry, 14th Edition. 2021. pp. 67-68.

The correct answer is: Plasma level of depot antipsychotics could be as high as 4 times the initial levels, 2 to 3 months after starting treatment

Question 56

Not answered

Marked out of 1.00

Which of the following is the most important property of a depot antipsychotic medication?

Select one:

- High water solubility
- The elimination is almost entirely renal
- Presence of a hydroxyl group
- Presence of a carboxyl ester
- A central dicyclobenzene ring

Your answer is incorrect.

Depot antipsychotics are esters formed between the alcohol group of the antipsychotic compound and a long chain of fatty acids (enantic, decanoic, palmitic, and undecylenic acid), and this ester is dissolved in oil (sesame oil or Viscoleo). Antipsychotic compounds are generally fat-soluble, but the esterification increases this fat solubility 1000 times (the longer the carbon chain of the fatty acid esters, the more the fat solubility). The ester is released from the oil phase by slow diffusion into the body water phase. The higher the fat solubility, the slower the release. (Kane et al., European Neuropsychopharmacology 1998, 8: 55-56)

The correct answer is: Presence of a carboxyl ester

Question 57

Not answered

Marked out of 1.00

Mrs. P suffers from bipolar disorder and is on lithium carbonate treatment. She also suffers from systemic hypertension. The ideal drug of choice for treating her hypertension would be

Select one:

- Candesartan
- Indapamide
- Ramipril
- Bendroflumethazide
- Propranalol

Your answer is incorrect.

Explanation:

ACE inhibitors (e.g. captopril, ramipril) can reduce thirst (which can lead to mild dehydration) and increase renal sodium loss (leading to increased sodium re-absorption by the kidney), resulting in an increase in lithium plasma levels. The magnitude of this effect is variable, from no increase to a four-fold increase. Care is also required with angiotensin II receptor antagonists such as candesartan. Diuretics can reduce the renal clearance of lithium, the magnitude of this effect being greater with thiazide than loop diuretics. Lithium levels usually rise within 10 days of a thiazide diuretic (e.g. bendroflumethiazide, chlortalidone, inadapamide, xipamide) being prescribed. Beta-blockers like propranalol are relatively safe to be used with lithium.

Ref: Taylor DM, Barnes TRE, Young AH. The Maudsley Prescribing Guidelines in Psychiatry, 14th Edition. 2021. pp. 252-253.

The correct answer is: Propranalol

Question 58

Not answered

Marked out of 1.00

Pharmacokinetic properties of SSRIs include all except

Select one:

- Rapid absorption from the gut
- Presence of active metabolite with longer half life for certain compounds
- Requirement for a washout period prior to administration of an MAOI
- Non linear elimination kinetics in high doses
- Long elimination half life

Your answer is incorrect.

Explanation:

SSRIs are well, but not rapidly, absorbed after oral administration. They have their peak effects in the range of 3-8 hours. Absorption of sertraline may be slightly enhanced by food. All SSRIs have a long half-life, which allows once daily administration. Fluoxetine has the most prolonged half-life: 4-6 days; its active metabolite has a half-life of 7-9 days. The half-life of sertraline is 26 hours, and its less active metabolite has a half-life of 3-5 days. Fluoxetine displays non-linear kinetics probably owing to the saturation of liver enzymes. Higher doses can result in disproportionately higher plasma levels.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 625.

The correct answer is: Rapid absorption from the gut

Question 59

Not answered

Marked out of 1.00

Choose the correct option regarding effects of smoking?

Select one:

- Doesn't activate P-450 1A2
- Doesn't activate P-450 2D6
- Increase the levels of haloperidol
- Increase levels of Fluvaxamine
- Reduce clozapine levels

Your answer is incorrect.

Explanation:

Tobacco smoke contains polycyclic aromatic hydrocarbons that induce (increase the activity of) certain hepatic enzymes (CYP1A2 in particular). The extent of enzyme induction is determined by the number and type of cigarettes smoked and the degree of smoke inhalation. For some drugs used in psychiatry, smoking significantly reduces drug plasma levels and higher doses are required than in non-smokers. Smoking reduces plasma clozapine levels by up to 50%.

Ref: Taylor DM, Barnes TRE, Young AH. The Maudsley Prescribing Guidelines in Psychiatry, 14th Edition. 2021. p. 856.

The correct answer is: Reduce clozapine levels

Question 60

Not answered

Marked out of 1.00

A child attains gender identity in which stage of Freud's psychosexual development?

Select one:

- Phallic phase
- Oral phase
- Latent phase
- Anal phase
- Genital phase

Your answer is incorrect.

Explanation:

The phallic stage begins sometime during year 3 and continues until approximately the end of year 5. The phallic stage lays the foundation for gender identity and serves to integrate residues from the previous stages into a predominantly genital-sexual orientation.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1062-1063.

The correct answer is: Phallic phase

Question 61

Not answered

Marked out of 1.00

At your CAMHS inpatient unit, you observe Paolo, a four-year-old boy playing with his dinosaur toy. He treats it as though it is a living object. According to Piaget's model, which stage of cognitive development is he in?

Select one:

- Pre-operationial
- Sensorimotor
- Post-operational
- Concrete Operational
- Formal operational

Your answer is incorrect.

Explanation.

This child is displaying animism. In Piaget's pre-operational stage, children use animistic thinking, which is the tendency to endow physical events and objects with life-like psychological attributes, such as feelings and intentions.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1008.

Stage	Age	Piaget's Cognitive Developmental Characteristics
Sensorimotor	0-2 years	Divided into six phases, characterised by: 1. Inborn motor and sensory reflexes 2. Primary circular reaction - coordinates activities of own body and five senses 3. Secondary circular reaction – seeks out new stimuli 4. Use of familiar means to obtain ends; plays peek-a-boo 5. Tertiary circular reaction and discovery through active experimentation 6. Insight and object permanence
Preoperational	2-7 years	Deferred imitation, symbolic play, graphic imagery (drawing), mental imagery, and language Egocentrism, phenomenalistic causality, and animistic thinking
Concrete operations	7-11 years	Conservation of quantity, weight, volume, length, and time based on reversibility by inversion or reciprocity; operational thought; class inclusion and seriation Syllogistic reasoning – logical conclusion formed from two premises
Formal operations	11 years-end of adolescence	Combinatorial system, whereby variables are isolated and all possible combinations are examined; hypotheticodeductive reasoning Complex use of language and abstract thinking

Ref: Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition, 2022



The correct answer is: Pre-operationial

Question 62

Not answered

Marked out of 1.00

Pauline is a first-time mother to Stephanie, her 18-month-old daughter. She is worried that Stephanie isn't talking as well as the other children at the nursery. If an 18-month-old child has normal development, how many words should he or she know at this stage?

Select one:

- 20-50
- 0-10
- 50-85
- 11-15
- 85-125

Your answer is incorrect.

Explanation:

By age 18 months, a child is generally using 20 single words. They will understand up to 150 words.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 889.

The correct answer is: 20-50

Question 63

Not answered

Marked out of 1.00

In your CAMHS clinic, you observe Mollie, a five-year-old girl, as she plays with her doll. You notice that she is talking to her doll as though it is real. What is the name of this phenomenon?

Select one:

- Animism
- Artificialism
- Centration
- Symbolic play
- Egocentrism

Your answer is incorrect.

Explanation:

Animism is the belief that plants, inanimate objects and natural phenomena possess a living soul and have human-like feelings and intentions. In Piaget's pre-operational stage, children use such animistic thinking to endow physical events and objects with life-like psychological attributes, such as feelings and intentions.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1008.

The correct answer is: Animism

Question 64

Not answered

Marked out of 1.00

According to Kohlberg, which one among the following represent Conventional Stage?

Select one:

- Authority
- Social Contract
- Universal Ethics
- Obedience
- Punishment

Your answer is incorrect.

Explanation:

Lawrence Kohlberg integrated Jean Piaget's concepts and described multiple stages of moral development within three significant levels of morality. The first level is pre-conventional morality, in which punishment and obedience are the determining factors. The second level is morality of conventional role-conformity, in which children try to conform to gain approval and maintain good relationships with others and are also guided by rigid codes of 'law and order' (authority orientation). The third and highest level is the morality of self-accepted moral principles (post-conventional), in which children voluntarily comply with rules based on a concept of universal ethical principles and make exceptions to rules in certain circumstances.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 902.

Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: Authority

Question 65

Not answered

Marked out of 1.00

Which of the following is seen in a normal child development?

Select one:

- Babbling with consonant sounds at 7 months
- Social smile at 6 months
- Toilet training at 5 years
- Putting two words together at 4 years
- Beginning to crawl at 15 months

Your answer is incorrect.

Babbling is the production of articulate sounds that are not yet recognizable words. It begins shortly after birth. At around 7 months, babbling contains alternations of vowels and consonants e.g. baba, mama (Ref. NHS)

The correct answer is: Babbling with consonant sounds at 7 months

Question 66

Not answered

Marked out of 1.00

In a strange situation experiment, Bobby, a 2-year-old boy, is only comforted when his mother returns to the room and is not comforted by the stranger. What type of attachment is seen here?

Select one:

- Secure attachment
- Anxious Avoidant
- Anxious Resistant
- Disorganized
- Insecure

Your answer is incorrect.

Explanation:

Mary Ainsworth devised the Strange Situation procedure with 12- to 18-month infants, which categorised infant-parent relationships. The procedure involves separations and reunions with caregivers and observes response. In secure attachment, the child is distressed by separation but can quickly be soothed on reunion with the caregiver.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 658-659.

The correct answer is: Secure attachment

Question 67

Not answered

Marked out of 1.00

Which of the following is NOT a risk factor for delayed speech development?

Select one:

- Small family size
- Absence of a father
- Low levels of parental education
- Rapid successive pregnancies
- Smoking in pregnancy

Your answer is incorrect.

The important family risk factors associated with delayed language development in children include large family size, smoking in pregnancy, rapid successive pregnancies, absence of father, low levels of parental education, divorce, marital disharmony & violence.

The correct answer is: Small family size

Question 68

Not answered

Marked out of 1.00

Which of the following has the strongest correlation with developmental language delay?

Select one:

- Maternal smoking during pregnancy
- Maternal alcohol use during pregnancy
- Low socio-economic status of mother
- High fibre diet
- Immigration

Your answer is incorrect.

Smoking during pregnancy and large family size are the strongest risk factors for developmental language delay

The correct answer is: Maternal smoking during pregnancy

Question 69

Not answered

Marked out of 1.00

The preferential attachment of a baby to one adult female figure is called;

Select one:

- Monotropy
- Secure attachment pattern
- Universal attachment
- Proximity seeking
- Separation distress

Your answer is incorrect.

Explanation:

Attachment theory originated in the work of John Bowlby. Being monotropic, infants tend to attach to one person, but they can form attachments to several people. A basic sense of security and safety is derived from a continuous and close relationship with a caregiver, usually the mother.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1011.

The correct answer is: Monotropy

Question 70

Not answered

Marked out of 1.00

Which one of the following is an objective test of personality?

Select one:

- Rorschach inkblot test
- Minnesota multiphasic personality inventory
- Sentence completion test
- Draw a person test
- Thematic apperception test

Your answer is incorrect.

Explanation:

The Minnesota Multiphasic Personality Inventory (MMPI) is a self-report questionnaire consisting of 567 questions covering eight areas of psychopathology and two areas of personality type. The scales are: Hypochondriasis, Depression, Hysteria, Psychopathic deviate, Masculinity-femininity, Paranoia, Psychasthenia, Schizophrenia, Hypomania, and Social introversion. There are also four additional validity (test-taking attitude) scales. It is NOT a projective test.

Projective tests of personality include the Rorschach test, Thematic Apperception Test (TAT), Sentence Completion Test, Holtzman Inkblot Technique (TIT), Figure drawing (typically human forms but can involve houses or other forms) and Make-a-Picture Story (MAPS). Projective tests use ambiguous stimuli, and the patient is expected to generate narratives based on them; rating scales may or may not be used in conjunction to rate such measures.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 96.

Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 59.

The correct answer is: Minnesota multiphasic personality inventory

Question 71

Not answered

Marked out of 1.00

The best neuropsychological test of semantic memory is

Select one:

- Naming objects and describing their meanings
- Address recall
- Rivermead behavioural memory test
- Vocabulary test
- Weschler's Memory Scale

Your answer is incorrect.

Explanation:

Semantic memory is a type of long-term memory for factual information about the world, excluding personal episodes in one's life. Typical examples are knowledge of the dates of the world wars and the chemical formula for water. The term is from the Greek 'semantikos', so called because it is necessary for the use of language.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: Naming objects and describing their meanings

Question 72

Not answered

Marked out of 1.00

Which of the following scales utilizes the theory of Big Five personality constructs?

Select one:

- Eysenck's Personality Inventory
- IPDE
- NEO Inventory
- Thematic Apperception Test
- Minnesota Multiphasic Personality Inventory

Your answer is incorrect.

Explanation:

NEO stands for Neuroticism, Extraversion, and Openness to experience. The inventory is a questionnaire designed to measure the Big Five personality factors, namely those three plus Agreeableness and Conscientiousness.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: NEO Inventory

Question 73

Not answered

Marked out of 1.00

In Skinner box experiments, rats receive food pellets in response to pressing a lever. What type of learning is this?

Select one:

- Backward conditioning
- Operant conditioning
- Forced conditioning
- Classical conditioning
- Forward conditioning

Your answer is incorrect.

Explanation:

The basic principles of operant conditioning are that if a response to a stimulus produces positive consequences for the individual, it will tend to be repeated, while if it is followed by negative consequences, it will not. In Thorndike and Skinner's original experiments, rats were placed in a box containing a level that, when pressed, delivered a pellet of food. Eventually, the rat would press the lever and be rewarded. The rat would then press the lever with increasing frequency.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 907.

The correct answer is: Operant conditioning

Question 74

Not answered

Marked out of 1.00

Star charts are effective in managing certain behavioural problems. They work on the basis of

Select one:

- Modelling
- Social learning
- Operant conditioning
- Classical conditioning
- Covert conditioning

Your answer is incorrect.

Explanation:

Behaviour modification is based on operant conditioning. Forms of behavioural modification include the use of secondary reinforcement, such as 'token economy', in which socially desirable/acceptable behaviours are rewarded with tokens that can be exchanged for other material items or privileges, or 'star charts' where children's good behaviour is rewarded when a certain level is achieved.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 908.

The correct answer is: Operant conditioning

Question 75

Not answered

Marked out of 1.00

A child was told to finish homework tasks before he can play outside. This is based on

Select one:

- Reciprocal inhibition
- Classical conditioning
- Cognitive mapping
- Premack principle
- Operant conditioning

Your answer is incorrect.

Explanation:

The Premack principle asserts that access to a preferred action will reinforce any action that is less preferred. Any behaviour that is preferred to a second behaviour will theoretically reinforce the second behaviour.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1019.

The correct answer is: Premack principle

Question 76

Not answered

Marked out of 1.00

With respect to memorising a list of words, which one among the following statements is true?

Select one:

- In immediate recall the primacy effect is stronger than the recency effect
- The retroactive inhibition refers to prior learning interfering with subsequent learning
- Primacy effect is related to initial items receiving most consolidation
- Elaborative rehearsal helps to store information in STM.
- Retrieval of material from semantic long term memory is an implicit process which requires clear knowledge of the time of acquisition of the semantic knowledge

Your answer is incorrect.

Explanation:

In serial learning, the primacy effect is a tendency for the items near the beginning of the series to be better recalled than those near the middle. An accurate recollection of an item is more likely if it is one of the first (primacy effect) or last items (recency effect) to be learnt. Primacy is because initial items receive most consolidation and recency is because immediate information is still in STM. The Proactive inhibition of 'interference theory' refers to prior learning interfering with subsequent learning. Retroactive inhibition is based on the fact that over time, new memories interfere with existing ones and this interference effect is increased if there is a considerable similarity between the new memory and existing ones. The other hypothesis for forgetting is decay theory due to the trace of the memory fading with time.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: Primacy effect is related to initial items receiving most consolidation

Question 77

Not answered

Marked out of 1.00

Which of the following is a type of non-declarative memory?

Select one:

- Semantic memory
- Autobiographical memory
- Procedural memory
- Episodic memory
- All of the listed options

Your answer is incorrect.

Explanation:

Non-declarative memory is memory for non-declarative knowledge, involving memory systems that do not draw on the individual's general knowledge. Non-declarative knowledge does not involve awareness and understanding of factual information about the world. It includes procedural knowledge (information about how to carry out sequences of operations – knowing **how** in contrast to knowing **that**).

Memory for factual knowledge and autobiographical events is collectively termed declarative memory. Declarative (also known as explicit) memory is of two types: episodic memory or memory for specific events (such as going to the shop this morning) and semantic memory or memory for abstract facts such as the capitals of countries.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015. pp. 510, 603.

Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1025.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 64.

The correct answer is: Procedural memory

Question 78

Not answered

Marked out of 1.00

A family has got a new dog. A 3 year old child in the family is afraid of dogs and would stay in the same room with the dog only if he could sit on his mother's lap and had his favourite ice cream. Select the learning theory principle employed here

Select one:

- Avoidance learning
- Extinction
- Operant conditioning
- Reciprocal inhibition
- Classical conditioning

Your answer is incorrect.

Explanation:

The modification of a behavioural pattern by the conditioning of responses that are incompatible with the response to be eliminated is called as reciprocal inhibition. The behavioral principle of **reciprocal inhibition** (Wolpe) states that anxiety and relaxation cannot coexist.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 908.

The correct answer is: Reciprocal inhibition

Question 79

Not answered

Marked out of 1.00

Identify the term that best denotes the feeling of familiarity that accompanies the return of stored material to consciousness;

Select one:

- Jamais vu
- Retrieval
- Dejavu
- Recognition
- Recollection

Your answer is incorrect.

Explanation:

Recognition is the act of process of perceiving or identifying information as matching or being the same as information that has been remembered. It is often contrasted with 'recall' which involves reproduction of remembered information without cues. Recognition is generally easier than recall.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: Recognition

Question 80

Not answered

Marked out of 1.00

Which of the following can reduce groupthink?

Select one:

- Strong chairperson
- Financial incentives
- Unplanned meetings of the group
- Seeking external opinion
- Pairing up group members

Your answer is incorrect.

Explanation:

Groupthink is a collective pattern of defensive avoidance, characteristic of group decision making in organisations in which group members develop rationalisations supporting shared illusions of their own infallibility and invulnerability with the organisation. Characteristic features include: an illusion of invulnerability, collective efforts to rationalise in order to discount warnings, an unquestioned belief in the group's inherent morality, stereotyped views of rivals and enemies, direct pressure on any member who dissents from the consensus view, self-censorship of disagreements with the apparent group consensus, a shared illusion of unanimity, and the emergence of self-appointed protectors of the group from adverse information that might undermine the existing complacency.

Various strategies that can reduce groupthink include appointing a 'devil's advocate', encouraging open debate, acknowledging the presence of groupthink, seeking external opinion, splitting the group into smaller units for discussion, holding last chance meetings to encourage challenges, and reserving leaders' opinions until after the group's discussion has been completed.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: Seeking external opinion

Question 81

Not answered

Marked out of 1.00

People prefer relationships that appear to offer an optimum cost benefit ratio and with expectations that our actions towards others will be reciprocated in some way. Which of the following theories suggest this?

Select one:

- Cannon Baird theory
- Functional attribution error
- James Lange theory
- Social Cognitive Theory
- Social exchange theory

Your answer is incorrect.

Explanation:

Social exchange theory is a theory of social interaction based on the proposition that people expect rewards and costs from a social exchange to be equitable. Most of our social behaviour is influenced by the expectations that our actions towards others will be reciprocated in some way and relationships are based on a mutual exchange of benefits. The US sociologist George Caspar Homans developed this theory.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: Social exchange theory

Question 82

Not answered

Marked out of 1.00

The ethics to determine goodness from acts and intentions behind them rather than their consequences is

Select one:

- Teleologic ethics
- Liberation ethics
- Utilitarian ethics
- Deontologic ethics
- Virtual ethics

Your answer is incorrect.

Explanation:

According to Immanuel Kant, a German philosopher, deontology is an ethical approach centered on rules and professional duties. Unlike many ethical theories, deontology does not focus on the consequences of individual actions. Personal emotions behind actions also do not matter with Kantian deontology because Kant believed humans do not always have rational control over their feelings. Instead, the intent behind chosen actions holds far more importance. Therefore, deontology proponents judge actions based on what people consider to be morally correct, regardless of actual consequences.

Consequentialist moral theories are teleological: they aim at some goal state and evaluate the morality of actions in terms of progress toward that state. The best-known version of consequentialism is utilitarianism. As most clearly stated by Mill a proponent of the utilitarianism (teleology), the basic principle of utilitarianism is: Actions are right to the degree that they tend to promote the greatest good for the greatest number.

The correct answer is: Deontologic ethics

Question 83

Not answered

Marked out of 1.00

A patient is next on waiting list to receive CBT. A new referral comes up which needs more urgent treatment. The psychotherapist refuses to displace the next patient on the list as this would be against his sense of duty and responsibility. Which ethical principle is he upholding?

Select one:

- Teleology
- Beneficience
- Deontic reasoning
- Non maleficence
- Autonomy

Your answer is incorrect.

Explanation:

Deontic reasoning is reasoning about obligation and duty, most often in relation to ethical or moral actions, but also in forms of non-ethical problem solving. Deontology and Teleology are two alternative higher-order ethical principles concerning medical practice. The term Deontology derives from the Greek 'Deon' for 'duty' indicating the centrality of rules in governing medical practice. Accordingly, rights and duties determine action and so it is also called as absolutism. The term Teleology derives its name from the Greek 'Teleon', meaning 'purpose' and the central concept is that rather than rights, people have interests, whether these are concerns, desires or needs. Accordingly, a broad judgment of benefits and harm determine medical practice.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: Deontic reasoning

Question 84

Not answered

Marked out of 1.00

Which of the following offers a guideline for principles of ethical research?

Select one:

- Hippocratic oath
- Mt. Sinai declaration
- Helsinki declaration
- Bristol code
- Warsaw pact

Your answer is incorrect.

Explanation:

The World Medical Association (WMA) has developed the Declaration of Helsinki as a statement of ethical principles for medical research involving human subjects, including research on identifiable human material and data. Adopted by the WMA's General Assembly in 1964, it is the most important set of guidelines about research on human participants. It both reflects and shapes the ethos of international research ethics. It is a living instrument and is reviewed and revised regularly. Its latest revision was in 2013.

Ref: https://www.wma.net/policies-post/wma-declaration-of-helsinki-ethical-principles-for-medical-research-involving-human-subjects/

Malik AY, Foster C. The revised Declaration of Helsinki: cosmetic or real change?. Journal of the Royal Society of Medicine. 2016 May;109(5):184-9.

The correct answer is: Helsinki declaration

Question 85

Not answered

Marked out of 1.00

A 40 year old man has chest pain and dizziness. His bloods and ECG are normal. He is advised to take medications, fat restricted diet, exercise. This is called

Select one:

- Illness behaviour
- Illness
- Sick role
- Worried well
- Depression

Your answer is incorrect.

Explanation:

Illness behaviour can be defined as any behaviour undertaken by an individual to relieve an experience of illness or to better define the meaning of the illness experience. Illness behaviours are considered to be the observable actions or reactions to perceived physical, emotional, or mental illness. These can include self-care or the use of healthcare services and may also include the interpretation and appraisal of symptoms, decisions to seek or delay health care, ongoing social and psychological coping strategies, adherence/concordance to recommended treatments and therapies, and the general ongoing uptake of available support services.

Ref: Murray J, Williams B. The Role of Images on Illness Behaviour: Interdisciplinary Theory, Evidence, and Ideas. Psychological Reports. 2020 Aug 5:1-23.

The correct answer is: Illness behaviour

Question 86

Not answered

Marked out of 1.00

Which of the following produced a written account on ethical aspects of medicine claiming that physicians have an obligation to protect both bodies and souls?

Select one:

- Burton
- Childress
- Hippocrates
- Charaka
- Ishaq bin Ali al-Ruhawi

Your answer is incorrect.

Explanation:

The oldest surviving Islamic work of medical ethics is the work of Ishaq bin Ali al-Ruhawi by the title of 'Adab al-Tabib' ('Practical Doctor's Ethics', 9th century) which included some of the ideas of Hippocrates and Galen. This work encouraged doctors to stay away from bad influences of people, to realise that good character is much more important than wealth, and to understand that wisdom is found in moral and perfect people.

In al-Ruhawi's opinion, doctors were responsible for the protection of both bodies and souls and one of the most important obligations of doctors is to be concerned about nurturing their own body and soul. He placed great emphasis on the interaction of the mental health and physical illness, though the mind is dealt mostly in spiritual sense. Interestingly, such an idea was not entirely novel at that time in the Islamic world. Another physician Abu Baker Al-Razi gave an earlier account of the physician's moral responsibilities in his volume Akhlaq Al-Tabib, "Medical Ethics," and the "Book of Spiritual Medicine," bringing spiritual aspects as important part of any medical care.

Ref: Zunic L et al. Medical ethics in the medieval Islamic sciences. Journal of research in pharmacy practice. 2014 Jul;3(3):75.

Muaygil RA. Demystifying bioethics: The past, present, and future of a flourishing discipline. Journal of Nature and Science of Medicine. 2020 Oct 1;3(4):334.

The correct answer is: Ishaq bin Ali al-Ruhawi

Question 87

Not answered

Marked out of 1.00

Which of the following model of reducing stigma demands equal rights and criminal responsibility for the mentally ill?

Select one:

- Disability inclusion model
- No fault approach
- Libertarian model
- Brain disease model
- Individual growth model

Your answer is incorrect.

The libertarian approach advocates equal rights with no discrimination - both positive and negative. It has the risk of inducing serial losses for the mentally ill at workplaces or courts.

The correct answer is: Libertarian model

Question 88

Not answered

Marked out of 1.00

Tearooom Trade Study is an example of ethically controversial research practices. Which of the following was the subject of investigation in this study?

Select one:

- Response to hepatitis vaccine
- Response to injected cancer cells
- Male sexual behaviours
- Natural history of syphilis
- Female contraception

Your answer is incorrect.

Explanation:

Laud Humphrey's *Tea Room Trade: Impersonal Sex in Public Places* is commonly presented in many sociological methods texts as an example of covert and deceptive research methods that endangered subjects without their consent. However, from the vantage point of beneficence as well as from a concern with social justice, Humphrey's tearoom study made significant positive contributions to the population he studied. His thorough fieldwork exploded the myth that the populations of men engaging in these casual sexual encounters were criminals, transients, or predatory paedophiles that communities needed to hunt down and criminally prosecute. Humphrey's work seriously questioned the received view that homosexual contact should be subjected to the retributive harms of imprisonment and ultimately created doubt in the dichotomous categories of gender that closely paralleled sexual behaviour.

Ref: Lenza M. Controversies surrounding Laud Humphreys' tearoom trade: An unsettling example of politics and power in methodological critiques. International Journal of Sociology and Social Policy. 2004.

The correct answer is: Male sexual behaviours

Question 89

Not answered

Marked out of 1.00

The nurses in your ward do not like a newly admitted patient. They are angry that she is trying to split them up by giving conflicting messages at ward rounds. Which of the following phenomenon is likely?

Select one:

- Introjection
- Identification
- Elimination
- Malignant alienation
- Transference

Your answer is incorrect.

Morgan (1979) coined the term malignant alienation to describe a process characterised by a progressive deterioration in the relationship between carers (staff in a ward) and a patient, including loss of sympathy and support from members of staff, who tended to construe these patients' behaviour as provocative, unreasonable, or overdependent. In some instances, such alienation may precede suicide/ attempted suicide of the patient (Excerpt from Watts & Morgan, Br J Psychiatry. 1994 Jan;164(1):11-5. http://www.ncbi.nlm.nih.gov/pubmed/8137091)

The correct answer is: Malignant alienation

Question 90

Not answered

Marked out of 1.00

Which one of the following is a feature of atypical Anorexia Nervosa?

Select one:

- Depressed mood
- Normal weight
- Binge eating
- Purging behaviour
- Normal menstruation

Your answer is incorrect.

Explanation:

In atypical anorexia nervosa (DSM-5), all the criteria for anorexia nervosa are met, except that despite significant weight loss, the individual's weight is within or above the normal range.

Ref: American Psychiatric Association. Diagnostic and statistical manual of mental disorders (DSM-5®). American Psychiatric Pub; 2013 May 22.

The correct answer is: Normal weight

Question 91

Not answered

Marked out of 1.00

Bleuler discarded the term dementia praecox primarily because

Select one:

- Mania was commonly coexistent with dementia praecox
- Not all cases began during adolescence
- Praecox affect was rarely seen in the early cases
- No treatment was available for dementia praecox
- No cognitive impairment was noted in the early cases

Your answer is incorrect.

Explanation:

In 1911, a book was published by Eugen Bleuler describing in detail asylum patients under his care who met clinical criteria for the psychotic disorder named Dementia Praecox by Emil Kraepelin. Bleuler followed Kraepelin and became the major figure to describe the characteristics of this syndrome in young adults, taking issue with Kraepelin on two points. He felt that cases of dementia praecox did not always arise in adolescence (praecox), nor did they always progress to intellectual dullness (dementia).

Ref: McGlashan TH. Eugen Bleuler: centennial anniversary of his 1911 publication of Dementia Praecox or the group of schizophrenias. Schizophrenia bulletin. 2011 Nov 1;37(6):1101-3.

The correct answer is: Not all cases began during adolescence

Question 92

Not answered

Marked out of 1.00

Mrs. Z arrives at work at least an hour earlier than anyone else to plan the day. She becomes visibly upset if the cleaners have moved anything on her desk and in her office, which is very tidy. This has led to some of her colleagues completely avoiding her office space. Her home and car are spotlessly clean. This is best described as

Select one:

- Obsessional personality trait
- Normal phenomenon
- Hoarding behaviour
- Obsessive compulsive disorder
- Compulsive behaviour

Your answer is incorrect.

Obsessive-compulsive personality disorder (OCPD) is a personality disorder characterized by excessive perfectionism and a need for control over one's environment. People with OCPD tend to be reliable and orderly, but also inflexible and unable to surrender control. This makes it hard for people with this condition to express their feelings, have close relationships and enjoy their successes. This lady is exhibiting some of the above traits.

The correct answer is: Obsessional personality trait

Question 93

Not answered

Marked out of 1.00

Which of the following is true with respect to adjustment disorder?

Select one:

- usually requires prolonged treatment
- correlates with the severity of the stressor
- occurs more often in males than in females
- occurs in all age groups
- it is a type of bereavement

Your answer is incorrect.

Although an adjustment disorder can occur at any age, it is more common at times in life when major transitions occur, such as adolescence, mid-life, and late-life (Mental Health: Adjustment Disorder - WebMD, http://www.webmd.com/mental-health/mental-health-adjustment-disorder).

The correct answer is: occurs in all age groups

Question 94

Not answered

Marked out of 1.00

With respect to models of diagnostic classification, which of the following is correct?

Select one:

- Ontological theories suggest that mental state does not exist by virtue of its abstract nature
- Fulford claims that psychiatric diagnoses are not value laden
- ICD-11 is entirely atheoretical
- Pheneticists stress on the natural existence of hierarchical structure among disease entities
- Categorical entities are not used in modern classification

Your answer is incorrect.

In contrast to the essentialist strategy that treats diseases as entities that exist in nature and form natural groups in essence, phenetic approach, (also called numerical taxonomy) shifts the emphasis to the systematic description of the observed disease presentations to enable manual classification. This approach treats all characters and attributes as having equal weight. While ICD and DSM are atheoretical in terms of assuming aetiology and causal models, they are not entirely atheoretical in descriptions used (e.g. dissociative disorder, conversion, etc. employ psychodynamic concepts).

The correct answer is: Ontological theories suggest that mental state does not exist by virtue of its abstract nature

Question 95

Not answered

Marked out of 1.00

Which of the following assessment is not a part of MMSE?

Select one:

- Repeating phrase
- Clock drawing
- Motor sequence
- Testing attention
- Copying polygons

Your answer is incorrect.

Explanation:

The MMSE is a 30-point cognitive test developed in the mid-1970s to provide a bedside assessment of a broad array of cognitive functions, including orientation, attention, memory, construction, and language. It does not include clock drawing.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 45.

The correct answer is: Clock drawing

Question 96

Not answered

Marked out of 1.00

Which of the following is a frontal lobe test?

Select one:

- Repertory grid
- Line bisection test
- Star cancellation test
- Cognitive estimates test
- Bar chart test

Your answer is incorrect.

Explanation:

Cognitive estimate testing assesses frontal lobe functioning. The patient is asked a question that requires abstract reasoning and cannot be answered by general knowledge alone (e.g. 'how many camels are there in the UK?').

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 88.

The correct answer is: Cognitive estimates test

Question 97

Not answered

Marked out of 1.00

Subtracting 7 out of 100 repeatedly (serial sevens test) is most useful to test one's

Select one:

- Attention
- Alertness
- Concentration
- Working memory
- Motivation

Your answer is incorrect.

Explanation:

Concentration can be tested by starting at 100 and counting backward by 7 (or 3). It can also be tested by saying the letters of the alphabet backward starting with Z or by naming the months of the year backward starting with December.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 11.

The correct answer is: Concentration

Question 98

Not answered

Marked out of 1.00

A 56-year-old woman is brought to the hospital with a stroke. A CT scan shows evidence of left occipital lobe damage. Which of the following is most likely to be seen?

Select one:

- Quadrantanopsia
- Cortical blindness
- Anosmia
- Bitemporal hemianopsia
- Apraxia

Your answer is incorrect.

Explanation:

'Cortical blindness' was first used by Gordon Holmes in 1918 to describe visual impairment of soldiers who were blind secondary to injuries to the occipital cortex. Cortical blindness is the total or partial loss of vision in a normal-appearing eye caused by damage to the occipital cortex of the brain. Cortical blindness can be acquired or congenital, and may also be transient in certain instances. Acquired cortical blindness is most often caused by loss of blood flow to the occipital cortex from either unilateral or bilateral posterior cerebral artery blockage due to ischemic stroke. In most cases, the patient may recover some of their vision.

The correct answer is: Cortical blindness

Question 99

Not answered

Marked out of 1.00

A boxer is knocked out by a blow to his lateral skull. Despite recovering after a few minutes, and being asymptomatic for the next 12 hours, he develops severe headache, vomiting and reduced consciousness. Which of the following is the most indicated investigation?

Select one:

- X ray skull
- SPECT scan
- EEG
- MRI brain
- CT scan brain

Your answer is incorrect.

Explanation:

The CT image is determined by the degree to which tissues absorb X-irradiation. This technique is particularly good at distinguishing between bones, blood vessels, and organs, so can well detect bone traumas, tumours, and haemorrhage. In a CT scan, CSF appears black, brain tissue appears grey, and the skull appears white. Blood appears bright white. As CT is quicker and relatively cheaper than MRI, it is far more commonly used in acute settings where medical stability still needs to be ascertained. In cases of suspected bony fracture and fresh hemorrhage, a CT scan is more useful than plain X-ray which cannot pick up the bleed, or MRI brain which is slower to obtain and may miss fresh bleeding.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 69.

Martinelli C, Shergill SS. Everything you wanted to know about neuroimaging and psychiatry, but were afraid to ask. BJPsych Advances. 2015 Jul;21(4):251-60.

The correct answer is: CT scan brain

Question 100

Not answered

Marked out of 1.00

Hyperhomocysteinemia is associated with increased risk for which of the following conditions?

Select one:

- Alcoholism
- Borderline personality disorder
- Panic disorder
- Self mutilation
- Dementia

Your answer is incorrect.

Explanation:

Elevated plasma total homocysteine is a modifiable risk factor for development of cognitive decline, dementia, and Alzheimer's disease in older persons. In a variety of clinical studies, the relative risk of dementia in elderly people for moderately raised homocysteine ranges from 1.1.5 to 2.5, and the Population Attributable risk ranges from 4.3 to 31%. Intervention trials in elderly with cognitive impairment show that homocysteine-lowering treatment with B vitamins markedly slows the rate of whole and regional brain atrophy and also slows cognitive decline. The findings are consistent with moderately raised plasma total homocysteine, which is common in the elderly, being one of the causes of age-related cognitive decline and dementia. Thus, the public health significance of raised tHcy in the elderly should not be underestimated, since it is easy, inexpensive, and safe to treat with B vitamins.

Ref: Smith AD et al. Homocysteine and dementia: an international consensus statement. Journal of Alzheimer's Disease. 2018 Jan 1;62(2):561-70.

The correct answer is: Dementia

Question 101

Not answered

Marked out of 1.00

Which of following is not commonly seen in nicotinic acid deficiency?

Select one:

- Fatigue
- Diarrhoea
- Dementia
- Dermatitis
- Mania

Your answer is incorrect.

Explanation:

Pellagra is a condition caused by low levels of niacin (nicotinic acid) or its precursor tryptophan. Niacin deficiency results in the condition of pellagra, which includes the triad or 'three D's': dermatitis, dementia, diarrhoea (related to mucosal atrophy), and can result in death. Similar to a sunburn, brown discolouration and lesions on the skin typically exposed to the sun such as hands, elbows, knees, and feet are present ('Casal's Necklace'). Initially, neurological changes such as anxiety, poor concentration, fatigue, and depression can manifest (due to reduced serotonin synthesis), but as pellagra advances, dementia and delirium may occur. Also, gastrointestinal complications, glossitis, cheilosis, stomatitis, nausea, vomiting, and diarrhoea (or less commonly constipation) may be present.

Ref: DeRight J. Vitamin Deficiencies. In Essential Neuropsychology: A Concise Handbook for Adult Practitioners 2022 (pp. 317-321). Springer, Cham.

The correct answer is: Mania

Question 102

Not answered

Marked out of 1.00

Which of the following best describes ideomotor apraxia?

Select one:

- Inability to coordinate and carry out facial and lip movements
- Inability to construct elements to a meaningful whole
- Difficulty in learning new information
- Inability to perform a multiple-step task
- Difficulty in copying skilled movements

Your answer is incorrect.

Explanation:

Ideomotor apraxia is characterised by the impaired ability to imitate unfamiliar actions. It is also called motor apraxia or ideokinetic apraxia. Ideomotor apraxia results from damage to the dominant parietal lobe. Abnormalities include (i) abnormal amplitude of motor movements (ii) body-part-as-object substitution, e.g. the patient uses his own finger to represent a toothbrush when asked to brush his teeth; (iii) abnormal orientation of body parts when attempting to perform an action.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: Difficulty in copying skilled movements

Question 103

Not answered

Marked out of 1.00

Which of the following is true regarding screening for alcohol problems?

Select one:

- Most patients give inaccurate drinking histories
- There are high rates of detection of alcohol problems in primary care and hospital settings
- Rates of detection are improved by asking routine alcohol-screening questions
- Diseases related to alcohol abuse respond poorly to early detection and intervention
- At-risk groups should not be specifically targeted for screening

Your answer is incorrect.

Explanation:

Diseases related to alcohol abuse are common, significant, and amenable to improvement by early detection and intervention. There are low rates of detection in primary care and hospital settings, which may be improved by increased vigilance, increased awareness of alcohol problems, awareness of routes of referral, asking routine alcohol screening questions, and paying special attention to at-risk groups. Many patients give reasonably accurate drinking histories if asked, although some may underestimate consumption.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 582.

The correct answer is: Rates of detection are improved by asking routine alcohol-screening questions

Question 104

Not answered

Marked out of 1.00

A patient with schizoaffective disorder is on both lithium and risperidone. He develops tremors as a side effect of his treatment. Which of the following will help detecting the offending agent?

Select one:

- Finger nose test
- Peripheral visual field
- Examination of tone of joints
- Frontal release sign
- Physical examination for tendon reflexes

Your answer is incorrect.

EPSE with risperidone can cause tremors (Parkinsonian) and rigidity, unlike Lithium tremor where one would not expect rigidity.

The correct answer is: Examination of tone of joints

Question 105

Not answered

Marked out of 1.00

When asked where do he lived when he was a child, a patient replied "Winchester. Manchester City was my favourite football team." The type of speech abnormality exhibited is

Select one:

- Clang association
- Pressured speech
- Tangentiality
- Coprolalia
- Circumstantiality

Your answer is incorrect.

Explanation:

This is an example of clang association. Clang association is an abnormality of speech where the connection between words is their sound, rather than their meaning. This may occur during manic flight of ideas.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 104.

The correct answer is: Clang association

Question 106

Not answered

Marked out of 1.00

Capgras syndrome (l'illusion des soies) is:

Select one:

- A form of negative autoscopy
- A pseudohallucination
- Also called the phantom mirror image
- A perceptual abnormality
- Delusional misidentification

Your answer is incorrect.

Explanation:

Capgras syndrome (l'illusion des sosies) is a type of delusional misidentification in which the patient believes that a person known to them has been replaced by a 'double' who is to all external appearances identical but is not the 'real person'.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 105.

The correct answer is: Delusional misidentification

Question 107

Not answered

Marked out of 1.00

A 20-year-old man is unable to express his problem coherently but appears very confused and feels that 'something is happening but not sure what it is'. This sense of perplexity and uncertainty that exists during a prodrome of psychosis may be linked most appropriately to which of the following experiences?

Select one:

- Negative symptoms
- Depression
- Delusional memory
- Delusional mood
- Autochthonous delusions

Your answer is incorrect.

Explanation:

Delusional mood is recalled as arising following a period when there is an abnormal mood state characterised by anticipatory anxiety, a sense of 'something is about to happen' and an increased sense of the significance of minor events. The development of the formed delusion may come as a relief to the patient in this situation.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 107.

The correct answer is: Delusional mood

Question 108

Not answered

Marked out of 1.00

A 23-year-old man notices the inscribed image of the Queen on a coin, and says "I know now, my mother is an alien". This is an example of

Select one:

- Ideas of reference
- Delusional mood
- Delusional memory
- Delusional misidentification
- Delusional perception

Your answer is incorrect.

Explanation:

A delusional perception is a primary delusion which is recalled as having arisen as a result of a perception (e.g. a patient who, on seeing two white cars pull up in front of his house, became convinced that he was therefore about to be wrongly accused of being a paedophile). The percept is a real external object, not a hallucinatory experience.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 107.

The correct answer is: Delusional perception

Question 109

Not answered

Marked out of 1.00

A 43-year-old woman, who was involved in a serious flight accident, felt at the time of the accident as if she had become unreal. For some time, she felt as if something had changed and that she was not herself. This is known as:

Select one:

- Nihilistic delusion
- Derealisation
- Suicidal thought
- Depersonalisation
- None of the above

Your answer is incorrect.

Explanation:

Depersonalisation is an unpleasant subjective experience where the patient feels as if they have become 'unreal'. It is a feeling that one is not oneself or that something has changed. Often this is associated with a feeling of unreality, so that the environment is experienced as flat, dull, and unreal (known as 'derealisation'). Many people who complain of depersonalisation also state that their capacity for feeling is diminished or absent. This is a subjective experience because to the outside observer there is no loss of ability to respond emotionally and appropriately to any given situation. Depersonalisation is not a delusion, and it should be distinguished from nihilistic delusions in which the patient denies that they exist or that they are alive, or that the work or other people exist. Depersonalisation is a non-specific symptom occurring in many psychiatric disorders, as well as in 'normal' people.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 108.

Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 10.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. pp. 81-82.

The correct answer is: Depersonalisation

Question 110

Not answered

Marked out of 1.00

One among the following statements is NOT true about depersonalisation?

Select one:

- Depersonalisation is easily distinguished from derealisation
- Depersonalisation is always subjective
- Depersonalisation is frequently accompanied by the symptom of derealisation
- Depersonalisation can be an experience of normal healthy people
- Affect is often involved in depersonalisation

Your answer is incorrect.

Explanation:

Depersonalisation is an unpleasant subjective experience where the patient feels as if they have become 'unreal'. It is a feeling that one is not oneself or that something has changed. Often this is associated with a feeling of unreality, so that the environment is experienced as flat, dull, and unreal (known as 'derealisation'). It is not necessarily easy to distinguish the two clinically. Many people who complain of depersonalisation also state that their capacity for feeling is diminished or absent. This is a subjective experience because to the outside observer there is no loss of ability to respond emotionally and appropriately to any given situation. Depersonalisation is not a delusion, and it should be distinguished from nihilistic delusions in which the patient denies that they exist or that they are alive, or that the work or other people exist. Depersonalisation is a non-specific symptom occurring in many psychiatric disorders, as well as in 'normal' people.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 108.

Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 10.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. pp. 81-82.

The correct answer is: Depersonalisation is easily distinguished from derealisation

Question 111

Not answered

Marked out of 1.00

Of various thought disorders classified by Andreasen, which of the following are more common in schizophrenia than mania?

Select one:

- Derailment
- Circumstantiality
- Neologisms
- Clanging
- Flight of ideas

Your answer is incorrect.

Explanation:

Nancy Andreasen created a scale in which the classic formal thought disorders are subdivided into three groups:

Communication disturbances: when the speaker does not meet the necessary requirements for the listener's understanding (poverty of content of speech, pressure of speech, distractible speech, tangentiality, derailment, stilted speech, echolalia, self-reference, circumstantiality, loss of goal, perseveration, and blocking)

Language disturbances: when the speaker violates the semantic and syntactic conventions (incoherence, clang association, neologisms, use of word approximations)

Thought disturbances: when only thinking alone seems affected (poverty of thought and illogically aberrant inferential processes)

Andreasen demonstrated that these psychopathological findings were not specific to schizophrenia and were also common in other mental disorders (e.g. mania). Of those listed, derailment (synonymous with loose associations - a symptom of schizophrenic thought disorder in which there is a total break in the chain of association between the meaning of thoughts) is more common in schizophrenia than mania.

Ref: Jerónimo J et al. Formal Thought Disorders–Historical Roots. Frontiers in psychiatry. 2018 Nov 9;9:572.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 108.

Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 10.

The correct answer is: Derailment

Question 112

Not answered

Marked out of 1.00

The phenomenon in which a subject perceives other people's heads to be as big as their actual body size can be termed as

Select one:

- Pareidolia
- Dysmegalopsia
- Lilliputian hallucinations
- Micropsia
- Macropsia

Your answer is incorrect.

Explanation:

Dysmegalopsia refers to a change in the perceived shape of an object. Some authors reserve the terms 'dysmegalopsia' to describe objects that are perceived to be larger (or smaller) on one side than the other, while others use the term generically to describe any change in perceived size.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 25.

The correct answer is: Dysmegalopsia

Question 113

Not answered

Marked out of 1.00

A patient repeats whatever the examiner says. He is exhibiting

Select one:

- Perseveration
- Logoclonia
- Parapraxis
- Stereotypy
- Echolalia

Your answer is incorrect.

Explanation:

Echolalia is the repetition of phrases or sentences spoken by others (e.g., the examiner). It is an echophenomenon that may be seen in catatonia. Other echophenomena are echopraxia (repeating what you do) and mirror movements (doing as you do at the same time).

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 109.

The correct answer is: Echolalia

Question 114

Not answered

Marked out of 1.00

The nature of speech produced in Broca aphasia is

Select one:

- Fluent and intelligible
- Impaired speech comprehension
- Fluent but unintelligible
- Effortful but agrammatism is seen
- Effortful but with grammatical use of language

Your answer is incorrect.

Explanation:

Broca aphasia (also called non-fluent or expressive aphasia) had traditionally been characterised by non-fluent speech and impaired repetition but intact auditory comprehension. It has long been thought to be associated with damage to Broca area (i.e. left inferior frontal convolution) or Brodmann area 44. Broca aphasia is characterised by retained use of nouns and verbs, but the patient may have lost all forms of pronouns, articles, and conjunctions, and hence may struggle to speak more than one word at a time even with enormous effort.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 48.

The correct answer is: Effortful but agrammatism is seen

Question 115

Not answered

Marked out of 1.00

Mr X was admitted to the psychiatric unit with a history of elated mood, overactivity, restlessness, grandiose delusions, disinhibited behaviour and was diagnosed with mania. A student nurse observed that he also experienced third person auditory hallucinations and wondered whether first rank symptoms could occur in manic patients. He wanted to clarify this with the treating psychiatric doctor in the team. Choose the most appropriate statement from the following:

Select one:

- First rank symptoms never occur in manic patients
- First rank symptoms may present in 10-25% of manic patients
- First rank symptoms are very rarely seen in manic patients
- First rank symptoms are present in all cases of mania with psychosis
- First rank symptoms are present in nearly half of all patients with mania

Your answer is incorrect.

Explanation:

First rank symptoms are a group of symptoms that have special significance in the diagnosis of schizophrenia. They are useful because they occur reasonably often in schizophrenia and more rarely in other disorders, and it is not too difficult to tell whether they are present or not. However, there is no symptom that is pathognomonic of schizophrenia. First rank symptoms can all be reported in other conditions (e.g. organic psychoses, manic illnesses). First rank symptoms are present in approximately 10-30% of patients with mania.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 110.

Channa A et al. First rank symptoms in Mania: An indistinct diagnostic strand. Acta Medica International. 2016 Jul 1;3(2):20.

The correct answer is: First rank symptoms may present in 10-25% of manic patients

Question 116

Not answered

Marked out of 1.00

Compulsory treatment cannot be justified under which of the following ethical principles?

Select one:

- Deontology
- Beneficence
- Autonomy
- Teleology
- Non maleficence

Your answer is incorrect.

Explanation:

A fundamental principle of medical care is that treatment of a patient should be with their consent. A patient has a right to decide for themselves which treatments to undergo and which treatments to refuse (autonomy). This right is retained, even when the refusal of treatment could result in death or significant deterioration in health. Treatment without consent is possible only in certain circumstances, constrained by appropriate laws.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 936.

The correct answer is: Autonomy

© 2023 SPMM Course Limited, UK.

Refund Policy | Terms & Conditions

