Started on	Sunday, 24 July 2022, 3:38 PM
State	Finished
Completed on	Sunday, 24 July 2022, 3:38 PM
Time taken	23 secs
Marks	0.00/189.00
Grade	**0.00** out of 100.00

Question **1**

Not answered

Marked out of 1.00

Flag question

The Doppleganger phenomenon is best described as a/an

Select one:

- Somatoform disorder
- Ideational disturbance
- Prerceptual disturbance
- Malingering phenomenon
- Body image disturbance

It refers to the awareness of another human being accompanying the self. Reported in psychosis and in severe sleep deprivation.

The correct answer is: Ideational disturbance

Question **2**

Not answered

Marked out of 1.00

Flag question

A 70-year-old patient with a diagnosis of chronic schizophrenia keeps touching her head as if she is saluting someone, even when no one is around. She denies 'hearing voices or seeing things'. Which of the following is she likely displaying?

Select one:

- Stereotypy
- Mannerism
- Echopraxia
- Chorea
- Hallucinatory behaviour

Explanation:

A mannerism is the abnormal and occasionally bizarre performance of a voluntary, goal-directed activity (e.g. a conspicuously dramatic manner of walking). Unusual, repeated performances of a goal-directed motor action or the maintenance of an unusual modification of an adaptive posture are considered mannerisms. Examples of this sign are unusual hand

movements while shaking hands, when greeting others, and during writing. Other examples may include peculiarities of dress, of hairstyle, and writing.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 114.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 90.

The correct answer is: Mannerism

Question **3**

Not answered

Marked out of 1.00

Flag question

Which of the following is a formal thought disturbance?

Select one:

- Thought insertion
- Obsessions
- Tangentiality
- Stuttering
- Delusions of persecution

Explanation:

Nancy Andreasen created a scale in which the classic formal thought disorders are subdivided into three groups:

- **Communication disturbances:** when the speaker does not meet the necessary requirements for the listener's understanding (poverty of content of speech, pressure of speech, distractible speech, tangentiality, derailment, stilted speech, echolalia, self-reference, circumstantiality, loss of goal, perseveration, and blocking)
- **Language disturbances:** when the speaker violates the semantic and syntactic conventions (incoherence, clang association, neologisms, use of word approximations)
- **Thought disturbances:** when only thinking alone seems affected (poverty of thought and illogically aberrant inferential processes)

Andreasen demonstrated that these psychopathological findings were not specific to schizophrenia and were also common in other mental disorders (e.g. mania).

Ref: Jerónimo J et al. Formal Thought Disorders–Historical Roots. Frontiers in psychiatry. 2018 Nov 9;9:572.

The correct answer is: Tangentiality

Question **4**

Not answered

Marked out of 1.00

Flag question

A depressed patient on the ward complained of total inability to experience pleasure. What is the term used to denote this type of pathological change in mood?

Select one:

- Apathy
- Affective flattening
- Affective blunting
- Anhedonia

- Incongruity of affect

Explanation:

Anhedonia is the feeling of absent or significantly diminished enjoyment of previously pleasurable activities. It is a core symptom of depressive illness and is also a negative symptom of schizophrenia.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 102.

The correct answer is: Anhedonia

Question **5**

Not answered

Marked out of 1.00

Flag question

An 82-year-old woman admitted to the psychiatric ward insists that she is already dead and should be buried, and that her viscera have stopped working and are rotting away. What is the probable diagnosis?

Select one:

- Couvade syndrome
- Briquet syndrome
- De Clerambault syndrome
- Cotard syndrome
- Hypochondriacal syndrome

Explanation:

Cotard syndrome is a presentation of psychotic depressive illness seen particularly in elderly people. There is a combination of severely depressed mood with nihilistic delusions and/or hypochondriacal delusions. The patient may state that they are already dead and should be buried, that their insides have stopped working and are rotting away, or that they have stopped existing altogether.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 105.

The correct answer is: Cotard syndrome

Question **6**

Not answered

Marked out of 1.00

Flag question

Apathy is characterised by

Select one:

- Lack of eye contact
- Lack of movements
- Lack of insight
- Lack of feeling
- Lack of speech

Explanation:

Apathy is a dulled emotional tone associated with detachment or indifference.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1136.

The correct answer is: Lack of feeling

Question **7**

Not answered

Marked out of 1.00

Flag question

Suggestibility is a prominent feature of

Select one:

- Perseveration
- Circumstantial thinking
- Tangentiality
- Over inclusive thinking
- Confabulation

The role of suggestibility is important in those who present with confabulation, pseudologia, retrospective falsification or false memory. Suggestible subjects accept statements from others, act upon their commands and deny evidence from the senses or from rational understanding that would contradict these statements.

The correct answer is: Confabulation

Question **8**

Not answered

Marked out of 1.00

Flag question

The patient would think 'I must put the kettle on', and after a pause of not more than one second would hear a voice say 'I must put the kettle on'. This is best described as

Select one:

- Thought insertion
- Thought diffusion
- Thought provocation
- Audible thoughts
- Thought broadcast

Explanation:

Audible thoughts are a form of auditory hallucination in which everything the patient thinks or speaks is repeated by the voices. It is also called thought echoing.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1136.

The correct answer is: Audible thoughts

Question **9**

A patient answers 'five' when asked how many days a week has. Which of the following is

Not answered

Marked out of 1.00

Flag question

true about the psychopathology exhibited by the patient?

Select one:

- It is done intentionally
- It is seen only in prison populations
- It is not seen in schizophrenia
- It is a disturbance of thought control
- It is a disturbance of thought form

Explanation:

Ganser syndrome is a disturbance of thought form characterised by approximate answers, disorientation, clouding of consciousness, hallucinations/pseudo-hallucinations, motor disturbance, anxiety or apathy, normal ADLs, and sudden resolution with amnesia for the period of illness. It is recognised to occur in a variety of psychiatric disorders, including schizophrenia, dissociative disorder, malingering, organic states, etc. In ICD-11, Ganser syndrome is included under ‘other specified dissociative disorders’.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 236.

Casey P, Kelly B. Fish’s clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 29.

Dieguez S. Ganser syndrome. Neurologic-Psychiatric Syndromes in Focus-Part II. 2018;42:1-22.

The correct answer is: It is a disturbance of thought form

Question **10**

Not answered

Marked out of 1.00

Flag question

A patient comes to the clinic and in a formal manner addresses you and the nurse as 'Hello sir, how are you? How is madam?' This is called

Select one:

- Stock word
- Neologism
- Scanning speech
- Stereotypy
- Stilted speech

Stilted speech - speech that is excessively stilted and formal. (e.g. 'The attorney comported himself indecorously'.)

The correct answer is: Stilted speech

Question **11**

Not answered

Marked out of

Which of the following is a thought disorder that predicts relapse in schizophrenia?

Select one:

1.00

Flag question

- Circumstantiality
- Tangentiality
- Concrete thinking
- Vorbeireden
- None of the listed options

FTDs seen in acute state have no prognostic value in predicting relapses.

The correct answer is: None of the listed options

Question **12**

Not answered

Marked out of 1.00

Flag question

Choose one incorrect statement about fugue states:

Select one:

- Consciousness is unimpaired
- It is a dissociative reaction to unbearable stress
- Fugue state generally has an insidious onset
- It is associated with amnesia
- Spontaneous resolution is a common outcome

Explanation:

Fugue is a dissociative reaction to unbearable stress. Following a severe external stressor, the affected individual develops sudden amnesia and may wander to a distant location. The amnesia will be for personal identity such as name, address, and history as well as for personal events, while at the same time the ability to perform complex behaviours is maintained. Consciousness is unimpaired. Recovery can be expected in time. Following resolution, there is amnesia for the events which occurred during the fugue.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 111, 869.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. pp. 64-65.

The correct answer is: Fugue state generally has an insidious onset

Question **13**

Not answered

Marked out of 1.00

Flag question

In which of the following forms does morbid jealousy NOT usually occur?

Select one:

- In dementia
- As a misidentification syndrome
- In depression
- As a delusion
- As an overvalued idea

Explanation:

Morbid jealousy (‘delusional jealousy’) is a delusional belief that one’s partner is being unfaithful. This can occur as part of a wider psychotic illness, secondary to organic brain damage, associated with alcohol dependence, or as a monosymptomatic delusional disorder (‘Othello syndrome’). It is not a misidentification syndrome.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 106.

The correct answer is: As a misidentification syndrome

Question **14**

Not answered

Marked out of 1.00

Flag question

In which of the following conditions the stimulus is perceived as corresponding object but the quality of the percept is altered?

Select one:

- Negative hallucinations
- Perceptual distortions
- Illusions
- Hallucinations
- Imagery

Distortions are changes in perception resulting from a change in the intensity and quality of the stimulus or the spatial form of the perception.

The correct answer is: Perceptual distortions

Question **15**

Not answered

Marked out of 1.00

Flag question

Erotomania has been reported in association with all EXCEPT

Select one:

- Paranoid schizophrenia
- Mania
- Delusional disorder
- Alcoholism
- Depersonalisation syndrome

DeClerambault's syndrome is a form of delusion of love (erotomania). The patient is usually a woman who believes that a celebrity / someone at a higher social status is in love with her.

The correct answer is: Depersonalisation syndrome

Question **16**

Not answered

Which one of the following phenomena can be described as ego-syntonic?

Marked out of 1.00

Flag question

Select one:

- Delusions
- Depersonalisation
- Derealization
- Both depersonalisation and derealization
- Obsessions

Explanation:

An experience is described as ego-syntonic if it is consistent with the goals and needs of the ego and/or consistent with the individual's ideal self-image; the reverse is the case for ego-dystonicity. In general, delusions are ego-syntonic, though as an individual with psychosis develops insight into their symptoms, delusions may seem increasingly ego-dystonic and distressing to them.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 47.

The correct answer is: Delusions

Question **17**

Not answered

Marked out of 1.00

Flag question

A patient with OCD has intrusive thoughts about the safety of his family. He fights these thoughts by often tapping his desk 5 times and also by praying mentally 5 times as fast as he can. The acts of praying are best termed as

Select one:

- Spiritual obsessions
- Delusions
- Normal behaviour
- Coping skills
- Compulsions

Explanation:

A compulsion is a behaviour or action that is recognised by the patient as unnecessary and purposeless, but that they cannot resist performing repeatedly. The drive to perform the action is recognised by the patient as their own, but it is associated with a subjective sense of need to perform the act, often in order to avoid the occurrence of an adverse event.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 104.

The correct answer is: Compulsions

Question **18**

Not answered

Marked out of 1.00

A middle-aged gentleman was noticed to be giggling inappropriately on hearing the news of his father's death. What is the term used to denote this type of pathological change in affect?

Select one:

Flag question

Affective blunting
- Apathy
- Affective flattening
- Anhedonia
- Incongruity of affect

Explanation:

Incongruity of affect refers to the objective impression that the displayed affect is not consistent with the current thoughts or actions (e.g. laughing while discussing traumatic experiences). It can occur in schizophrenia.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 113.

The correct answer is: Incongruity of affect

Question **19**

Not answered

Marked out of 1.00

Flag question

Delusions of motor control are thought to be related to

Select one:
- Loss of proprioceptive reflexes
- A form of motor neglect
- Paroxysmal spike activity in premotor area
- Failure to predict sensory feedback of one's own movements
- Inability to control execution of motor maps

According to this sensory-feedback model if something seems to be going wrong with the action, it is quite possible to correct for it on the basis of concurrent self-awareness. Delusions of motor control are thought to be caused by a failure to predict sensory feedback of own movements (One of the models used to explain delusions of control in patients with schizophrenia) [Ref: Gallagher, S. Psychopathology 2004; 37:8-19]

The correct answer is: Failure to predict sensory feedback of one's own movements

Question **20**

Not answered

Marked out of 1.00

Flag question

A voice in a flat monotone describes everything the patient is doing: 'She is peeling potatoes, she has now got hold of the peeler, she does not want that potato...' This is best described as

Select one:
- Running commentary
- Second person hallucinations
- Imperative hallucinations
- First person hallucinations
- Autochthonous perceptions

Explanation:

'Running commentary' is a type of third-person auditory hallucination that is a first-rank symptom of schizophrenia. The patient hears one or more voices providing a narrative of their current actions.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 117.

The correct answer is: Running commentary

Question **21**

Not answered

Marked out of 1.00

Flag question

Which of the following is true with regard to the mental state of ecstasy?

Select one:

- Associated with persecutory ideas
- Associated with somatoform pain state
- Cannot occur in the absence of pathological phenomenon
- Associated with altered boundary of self
- Some degree of altered consciousness is always seen

Explanation:

Ecstasy is a state of extreme wellbeing associated with a feeling of rapture, bliss, and grace. It can occur in the healthy population at times of profound religious experience or occasions of deep emotion such as following childbirth. The mind is usually occupied with a feeling of communion with God or some religious figure. There may be a feeling of being in tune with the whole of nature and at one with the universe. The person in ecstasy experiences the change in ego boundaries as voluntary.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 79.

The correct answer is: Associated with altered boundary of self

Question **22**

Not answered

Marked out of 1.00

Flag question

Changes in the shape of objects especially with a loss of symmetry is called

Select one:

- Eidetic imagery
- Macropsia
- Micropsia
- Dysmegalopsia
- Asterognosis

Explanation:

Dysmegalopsia refers to a change in the perceived shape of an object. Some authors reserve the terms 'dysmegalopsia' to describe objects that are perceived to be larger (or smaller) on

one side than the other, while others use the term generically to describe any change in perceived size. Yet others use the term ‘metamorphosia’ rather than dysmegalopsia to describe objects that are irregular in shape.

Ref: Casey P, Kelly B. Fish’s clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 25.

The correct answer is: Dysmegalopsia

Question **23**

Not answered

Marked out of 1.00

Flag question

Concrete thinking is usually tested by means of

Select one:

- Verbal fluency test
- Proverb testing
- Cognitive estimates test
- Luria test
- Sorting test

Explanation:

Concrete thinking is characterised by the inability to understand abstract concepts and metaphorical ideas, leading to a strictly literal form of speech and the inability to comprehend allusive language (as present in proverbs). Concrete thinking can be seen in schizophrenia and dementing illnesses.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 104.

The correct answer is: Proverb testing

Question **24**

Not answered

Marked out of 1.00

Flag question

You are interviewing a newly admitted patient in your ward. When asked what he had for breakfast, he says 'cereals'. Then when you asked 'what is your name?' he says 'cereals'. When you asked 'what is his date of birth?' he continues to say 'cereals'. The most likely symptom exhibited here is

Select one:

- Stuttering
- Stereotypy
- Automatism
- Perseveration
- Mannerism

Explanation:

Perseveration is characterised by continuing with a verbal response or action that was initially appropriate after it ceases to be apposite. It is associated with organic brain disease and is occasionally seen in schizophrenia.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 115.

The correct answer is: Perseveration

Question **25**

Not answered

Marked out of 1.00

Flag question

Which of the following is true with regard to primary delusions?

Select one:

- They are more common in chronic rather than in acute schizophrenia
- They are usually well elaborated
- The content is usually aligned to cultural and social norms
- Their origin can be traced back to a person's premorbid mental state
- By definition, they are not understandable

Explanation:

Primary delusions are the direct result of psychopathology, while secondary delusions can be understood as having arisen in response to other primary psychiatric conditions. Jaspers maintained that primary delusions were typical of un-understandable phenomena, being immediately experienced by the patients as absolutely self-evident like an intuition or a revelation and accompanied by abnormalities in the form such as radical transformation or emergence of thinking. He designated primary delusions as delusion proper, from which secondary delusions evolve via conscious thought processes.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 106.

Hayashi N et al. Delusion progression process from the perspective of patients with psychoses: A descriptive study based on the primary delusion concept of Karl Jaspers. Plos one. 2021 Apr 27;16(4):e0250766.

The correct answer is: By definition, they are not understandable

Question **26**

Not answered

Marked out of 1.00

Flag question

Vorbeireden is seen in

Select one:

- Couvade syndrome
- Tourette syndrome
- Ganser syndrome
- Cotard syndrome
- Asperger syndrome

Explanation:

Vorbeireden (also known as ‘vorbeigehen’ or ‘talking past the point’) is seen in Ganser syndrome. It refers to the production of ‘approximate answers’ in which the content of the patient’s replies to questions shows that they understand what has been asked but have responded by talking about an associated topic. Ganser syndrome is characterised by

approximate answers, disorientation, clouding of consciousness, hallucinations/pseudo-hallucinations, motor disturbance, anxiety or apathy, normal ADLs, and sudden resolution with amnesia for the period of illness. In ICD-11, it is included under 'other specified dissociative disorders'.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 119, 236.

Dieguez S. Ganser syndrome. Neurologic-Psychiatric Syndromes in Focus-Part II. 2018;42:1-22.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 57.

The correct answer is: Ganser syndrome

Question **27**

Not answered

Marked out of 1.00

Flag question

Deja vu and jamais vu are experiences of

Select one:

- Familiarity
- Volition
- Perception
- Behaviour
- Mood

Explanation:

Déjà vu is a sense that events being experienced for the first time have been experienced before. Jamais vu is the sensation that events or situations are unfamiliar, although they have been experienced before. Both déjà vu and jamais vu can be everyday experiences, but also non-specific symptoms of a number of disorders, including temporal lobe epilepsy, schizophrenia, and anxiety disorders. They are best described as states involving familiarity.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 105, 113.

The correct answer is: Familiarity

Question **28**

Not answered

Marked out of 1.00

Flag question

Kinaesthetic hallucination is reported in cases with

Select one:

- Amphetamine withdrawal
- Benzodiazepine withdrawal
- Cocaine withdrawal
- Alcohol withdrawal
- Cannabis withdrawal

Symptoms of benzodiazepine withdrawal include autonomic hyperactivity, increased tremor,

insomnia, nausea or vomiting, transient visual, tactile, or auditory hallucinations or illusions, psychomotor agitation. anxiety (most prominent), with grand mal seizures and kinaesthetic hallucinations reported in a few patients. Excerpt from Alcohol-Related Problems: Recognition and Intervention, http://www.aafp.org/afp/1999/0115/p361.html (accessed April 11, 2015).

The correct answer is: Benzodiazepine withdrawal

Question **29**

Not answered

Marked out of 1.00

Flag question

In most cases of dysmorphophobia, the affected mental state is associated with

Select one:

- Anxiety disorder
- Major mood disorder
- Personality disorder
- Psychotic disorder
- Eating disorder

Most cases of dysmorphophobia is associated with a major mood disorder. Most patients with Dysmorphophobia have other associated conditions like mood disorder, anxiety disorder and psychosis.

The correct answer is: Major mood disorder

Question **30**

Not answered

Marked out of 1.00

Flag question

Patients with chronic schizophrenia may lose the track of their age and may state an age different from their actual age. To be called as age disorientation, the difference in stated age and actual age must be at least

Select one:

- One year
- 10 years
- 5 years
- 20 years
- 3 years

Explanation:

Age disorientation is defined as misstating one's age by 5 or more years. It was initially described as 'perceptual distortion of temporal orientation' in early literature. The majority of age disoriented patients underreport their age (i.e. *time stood still),* with a small percentage overestimating their age.

Ref: Zhand N et al. Incidental finding of age and time disorientation in a clinical sample of patients with schizophrenia: Findings from a short term follow up survey regarding cannabis use. Psychiatry Research. 2020 Aug 1;290:113150.

The correct answer is: 5 years

Question **31**

Not answered

Marked out of 1.00

Flag question

A patient who is almost completely bald was observed to imitate combing his hair repeatedly but without a comb or brush. He is diagnosed to have schizophrenia. Which of the following is the most likely phenomenon observed?

Select one:

- Perseveration
- Stereotypy
- Mannerism
- Passivity
- Akathisia

Explanation:

A mannerism is the abnormal and occasionally bizarre performance of a voluntary, goal-directed activity (e.g. a conspicuously dramatic manner of walking).

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 114.

The correct answer is: Mannerism

Question **32**

Not answered

Marked out of 1.00

Flag question

Which of the following statements is not true with regard to depersonalisation?

Select one:

- Depersonalisation is rare in psychiatric practice
- Depersonalisation is often a difficult experience to describe
- Depersonalisation is a common experience of normal healthy people
- Depersonalisation is often accompanied by the symptom of derealization
- Affect is invariably involved in depersonalisation

Explanation:

Depersonalisation is an unpleasant subjective experience where the patient feels as if they have become ‘unreal’. It is a feeling that one is not oneself or that something has changed. Depersonalisation is a non-specific symptom occurring in many psychiatric disorders, as well as in ‘normal’ people.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 108.

Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 10.

The correct answer is: Depersonalisation is rare in psychiatric practice

Question **33**

Not answered

A patient gets recurrent intrusive thoughts about death. This sometimes involves her own death or death of her loved ones. She does not believe they are true, but she is afraid she

Marked out of 1.00

Flag question

might yield to them. The form of psychopathology exhibited here is

Select one:

- Obsessions
- Suicidal ideas
- Loss of control
- Delusions
- Homicidal ideas

Explanation:

An obsession is an idea, image, or impulse that is recognised by the patient as their own, but which is experienced as repetitive, intrusive, and distressing. The return of the obsession can be resisted for a time, at the expense of mounting anxiety. In some situations, the anxiety accompanying the obsessional thoughts can be relieved by associated compulsions.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 115.

The correct answer is: Obsessions

Question **34**

Not answered

Marked out of 1.00

Flag question

A sense of perplexity and uncertainty that exists during a prodrome of psychosis may be linked most appropriately to which of the following experiences?

Select one:

- Negative symptoms
- Delusional memory
- Autochthonous delusions
- Depression
- Delusional mood

Explanation:

Delusional mood is recalled as arising following a period when there is an abnormal mood state characterised by anticipatory anxiety, a sense of ‘something is about to happen’ and an increased sense of the significance of minor events. The development of the formed delusion may come as a relief to the patient in this situation.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 107.

The correct answer is: Delusional mood

Question **35**

Not answered

Marked out of 1.00

Epileptic automatism is associated with

Select one:

- Preserved consciousness

Flag question

- Loss of postural control
- Presence of aura
- Loss of muscle tone
- Duration lasting few seconds to few minutes

Explanation:

An automatism is a behaviour which is apparently conscious in nature, occurring in the absence of full consciousness (e.g. during a temporal lobe seizure). They typically last for several seconds to minutes. Common automatisms include oral automatisms (repetitive licking or smacking of the lips, swallowing, or kissing) and manual automatisms (fumbling with clothing, scratching, rubbing the abdomen, or fidgeting). Other physical manifestations are simple actions such as standing, walking, pacing, or even driving.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 103.

Kaufman DM, Geyer HL, Milstein MJ, Rosengard JL. Kaufman's Clinical Neurology for Psychiatrists, 9th Edition. 2023. p. 214.

The correct answer is: Duration lasting few seconds to few minutes

Question **36**

Not answered

Marked out of 1.00

Flag question

Which of the following statements about alcoholic hallucinosis is incorrect?

Select one:

- Patients with alcoholic hallucinosis have a clear sensorium
- These usually appear in persons abusing alcohol for a long time
- The hallucinations respond poorly to antipsychotics
- The hallucinations usually last less than a week, when patients believe in the hallucinations though afterwards they may realise the untrue nature.
- The most common hallucinations are unstructured sounds, or voices that may be characteristically malign and threatening

Explanation:

Alcoholic hallucinosis is a substance-induced psychotic illness that is a rare complication of **prolonged heavy alcohol use.** InICD-11**,** it is designated as "alcohol-induced psychotic disorder with hallucinations". It is characterised by experiencing **hallucinations** (usually auditory) **while sober.** It usually starts with **elemental hallucinations** (e.g. bangs or murmurings). With continued alcohol use they become **formed voices,** usually **derogatory. Unlike delirium tremens**, those with alcoholic hallucinosis have a **clear sensorium**. In 95% of patients, **symptoms rapidly resolve on ceasing alcohol consumption**, but rapidly recur on restarting drinking. In 5%, symptoms persist with abstinence. **Persisting symptoms** may be treated with antipsychotics.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 602.

The correct answer is: The hallucinations respond poorly to antipsychotics

Question **37**

Not answered

Marked out of 1.00

Flag question

A patient experiences some disturbing thoughts. He feels these thoughts originate elsewhere and reach him fully formed. He does not feel he owns these thoughts. Which of the following terms best describes the above?

Select one:

- Thought withdrawal
- Thought insertion
- Obsessions
- Thought broadcasting
- Thought blocking

Explanation:

Thought insertion is the delusional belief that thoughts are being placed in the patient's head from the outside. Thought insertion is a first-rank symptom of schizophrenia.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 118.

The correct answer is: Thought insertion

Question **38**

Not answered

Marked out of 1.00

Flag question

While assessing the psychopathology of a psychotic patient, you notice that a major thought gets replaced by a minor thought in his thinking process. This is called

Select one:

- substitution
- derailment
- drivelling
- fusion
- omission

Explanation:

Carl Schneider claimed that five features of formal thought disorders could be identified:

- Derailment: the thought slides on to a subsidiary thought
- Substitution: a major thought is substituted by a subsidiary one
- Omission: senseless omission of a thought or part of it
- Fusion: heterogenous elements of thought are interwoven with each other
- Drivelling: a disordered intermixture of constituent parts of one complex thought

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 55.

The correct answer is: substitution

Question **39**

Not answered

Marked out of 1.00

Flag question

A patient says 'I am constantly being followed by people I know, but I can't recognize them because they are always in disguise'. What symptom is she exhibiting?

Select one:

- Capgras delusion
- Fregoli delusion
- Intermetamorphosis
- Cotard delusion
- De Clerambault delusion

Explanation:

Fregoli syndrome is a type of delusional misidentification in which the patient believes that strangers have been replaced with familiar people. It may be believed that various people whom the subject meets are really the same person, in disguise. For example, a patient believed that her neighbour could change his appearance, clothes, and even his sex at will so as to spy on her.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 111.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 127.

The correct answer is: Fregoli delusion

Question **40**

Not answered

Marked out of 1.00

Flag question

A woman reports that she can see her recently deceased husband though she does not believe he could be alive. This is best described as a

Select one:

- Perceptual distortion
- Visual hallucination
- Affect illusion
- Pseudo-hallucination
- Pareidolia

A pseudo-hallucination is a false perception that is perceived as occurring as part of one's internal experience, not as part of the external world. It may be described as having an 'as if' quality or as being seen with the mind's eye. Pseudo-hallucinations are a type of mental image that, although clear and vivid, lack the substantiality of perceptions. Pseudo-hallucinations are seen in full consciousness, known to be not real perceptions, and are located not in objective space but in subjective space (for example, inside the head). Like true hallucinations, they are involuntary. Pseudo-hallucinations can occur in all modalities of sensation and are described in psychotic, organic, and drug-induced conditions, as well as occasionally in 'normal' individuals. (The hallucinations of deceased spouses commonly described by widows and widowers may have a form of a pseudo-hallucination.)

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 116.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 28.

The correct answer is: Pseudo-hallucination

Question **41**

Not answered

Marked out of 1.00

Flag question

Phantom mirror image is a synonymous term for which one of the following hallucinations?

Select one:

- Reflex hallucination
- Autoscopic hallucination
- Haptic hallucination
- Hypnagogic hallucination
- Teichoscopic hallucination

Explanation:

Autoscopy (also known as 'phantom mirror image') is the experience of seeing a visual hallucination or pseudo-hallucination of oneself. It is not just a visual hallucination because kinaesthetic and somatic sensation must also be present to give the subject the impression that the hallucination is oneself.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 103.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019.p. 36.

The correct answer is: Autoscopic hallucination

Question **42**

Not answered

Marked out of 1.00

Flag question

When one of the other residents pushed a salt cellar towards him, a patient knew that he must return home 'to greet the Pope who is visiting Ireland to see [his] family and reward them'. This is called

Select one:

- Partial delusion
- Delusional misinterpretation
- Referential delusion
- Grandiose delusion
- Delusional perception

Explanation:

A delusional perception is a primary delusion which is recalled as having arisen as a result of a perception (e.g. a patient who, on seeing two white cars pull up in front of his house, became convinced that he was therefore about to be wrongly accused of being a paedophile). The percept is a real external object, not a hallucinatory experience.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul

30. p. 107.

The correct answer is: Delusional perception

Question **43**

Not answered

Marked out of 1.00

Flag question

A patient on an adult psychiatric ward says that Dr. Smith has been his consultant for the past 15 years. Even though he recognises his face, he knows he is not Dr. Smith anymore but an imposter. What is this phenomenon called?

Select one:

- Capgras syndrome
- Fregoli syndrome
- Borognosia
- Prosopognosia
- Illusion of doubles

Explanation:

Capgras syndrome is a type of delusional misidentification in which the patient believes that a person known to them has been replaced by a 'double' who is to all external appearances identical, but is not the 'real person'. This syndrome can occur both in psychiatric disorders, mainly psychotic disorders (especially schizophrenia) as well as major affective disorders, and neurocognitive disorders (e.g., post-traumatic encephalopathy, epilepsy, cerebrovascular disease, dementia, and delirium/acute confusional state).

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 104.

Karakasi MV et al. In fear of the most loved ones. A comprehensive review on Capgras misidentification phenomenon and case report involving attempted murder under Capgras syndrome in a relapse of a schizophrenia spectrum disorder. Journal of forensic and legal medicine. 2019 Aug 1;66:8-24.

The correct answer is: Capgras syndrome

Question **44**

Not answered

Marked out of 1.00

Flag question

Which of the following is not included in the classification of Delusional Disorders in ICD 10?

Select one:

- Delusions of infestations
- Delusions of grandiosity
- Delusions of control
- Delusions of love
- Delusions of jealousy

Delusions of control are bizarre delusions seen often in schizophrenia-like disorders, and not in a persistent delusional disorder.

The correct answer is: Delusions of control

Question **45**

Not answered

Marked out of 1.00

Flag question

Pseudologia fantastica is most commonly associated with

Select one:

- Pick dementia
- Frontal lobe damage
- Obsessional personality
- Dissocial personality
- Borderline personality/Narcissistic personality

Pseudologia fantastica or fluent plausible lying (pathological lying) is seen in personality disorders of antisocial or hysterical type. It often carries a grandiose tinge and can be triggered by a personal crisis, such as facing legal proceedings. A person with pseudologia may believe their own stories and often there is a blur between fantasy and reality. When confronted, these individuals may admit to lying.

The correct answer is: Dissocial personality

Question **46**

Not answered

Marked out of 1.00

Flag question

Dereistic thinking is a term used to describe which one of the following?

Select one:

- Lateral thinking
- Abstract thinking
- Logical thinking
- Unrealistic thinking
- Goal directed thinking

Explanation:

Dereism is mental activity that follows a totally subjective and idiosyncratic system of logic. It is undirected fantasy thinking that fails to take the facts of reality or experience into consideration.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1138.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 43.

The correct answer is: Unrealistic thinking

Question **47**

Not answered

Marked out of 1.00

A 45-year-old man in a psychiatric inpatient unit is noted to move his right arm inwards while throwing his left arm outwards repeatedly while walking in the corridor. This is not noted when he is sitting in the lounge. He is exhibiting

Select one:

Flag question

- Parkinsonism
- Akathisia
- Dystonia
- Mannerism
- Stereotypy

Explanation:

A mannerism is the abnormal and occasionally bizarre performance of a voluntary, goal-directed activity (e.g. a conspicuously dramatic manner of walking).

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 114.

The correct answer is: Mannerism

Question **48**

Not answered

Marked out of 1.00

Flag question

Agitated depression is a well-recognized feature of

Select one:

- Depression in children
- Depression in young men
- Depression in young women
- Depression in the elderly
- Depression in middle aged women

Explanation:

Agitated depression is a combination of depressed mood and psychomotor agitation, contrasting with the more usual association of depressed mood with psychomotor retardation. It is a common presentation of depression in the elderly.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 102.

The correct answer is: Depression in the elderly

Question **49**

Not answered

Marked out of 1.00

Flag question

Which of the following is true concerning the psychopathology of morbid jealousy?

Select one:

- It is not seen as an isolated phenomenon
- The belief can be held with delusional fixity
- It is more common in women than men
- It is an obsessional state
- It intensifies on geographical separation of partners

Explanation:

Morbid jealousy (‘delusional jealousy’) is a fixed delusional belief that one’s partner is being unfaithful. This can occur as part of a wider psychotic illness, secondary to organic brain damage, associated with alcohol dependence, or as a monosymptomatic delusional disorder (‘Othello syndrome’). It is more common in men. Apart from delusions of infidelity, these patients tend not to show any other symptoms. Separation from the focus of delusional ideas is advised, as behaviour may progress to violence against the partner and even to murder.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 106, 230-234.

Casey P, Kelly B. Fish’s clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 51.

The correct answer is: The belief can be held with delusional fixity

Question **50**

Not answered

Marked out of 1.00

Flag question

Which one of the following is not a core feature of Ganser syndrome?

Select one:

- Visual hallucinations
- Somatic conversion features
- Clouding of consciousness
- Approximate answers or VORBEIGEHEN (to pass by)
- Pseudohallucinations

Ganser syndrome refers to the production of approximate answers. e.g. Question: "What is the capital of France?" - Answer: "London". It is occasionally associated with organic brain illness but is much more commonly seen as a form of malingering in those attempting to feign mental illness e.g. in prisoners awaiting trial.

The correct answer is: Visual hallucinations

Question **51**

Not answered

Marked out of 1.00

Flag question

The term anhedonia was coined by

Select one:

- Bleuler
- Ribot
- Kraepelin
- Sifneos
- Kretschmer

Explanation:

Anhedonia is an inability to enjoy anything in life or even get pleasure from everyday events (for example, being hugged by one’ children or waking up to a fine spring morning). ‘Anhedonia’ was coined by Ribot, a French psychologist, in 1896.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 77.

The correct answer is: Ribot

Question **52**

Not answered

Marked out of 1.00

Flag question

A patient repeats the last word that she uttered without any apparent purpose. This is called

Select one:

- Echolalia
- Logoclonia
- Perseveration
- Palilalia
- Stammering

Explanation:

Logoclonia refers to meaningless repetition of a syllable, especially an end syllable of a word. Palilalia and stuttering have disorders that are similar to those of logoclonia. Palilalia, however, refers to the repetition of an entire word, but not a syllable. Stuttering is the involuntary disruption or blocking of speech with sound repetition, especially the initial consonants of a word, and, unlike logoclonia, stuttering is accompanied by tense and effortful utterance.

Ref: Nakagawa Y et al. Logoclonia might be a Characteristic of Logopenic Variant Primary Progressive Aphasia at an Advanced Stage: Potential Mechanisms Underlying Logoclonia. Journal of Alzheimer's Disease. 2019 Jan 1;70(2):515-24.

The correct answer is: Palilalia

Question **53**

Not answered

Marked out of 1.00

Flag question

Common characteristics of schizophrenic auditory hallucinations include all except

Select one:

- Speaking in one's mother tongue
- Multiple voices
- Having different accents
- Present throughout the day incessantly
- Male voices

Explanation:

Auditory hallucinations in schizophrenia are often experienced as voices although they can also take the form of other non-verbal sounds. Many such voices can be reported by patients as being similar or just like hearing other people speak. Voice-hearers report on average 3 different voices. Voices may be male or female; however, male voices are more common than female voices. Hallucinated voices frequently speak with accents different to the region or social class of the patient. There is considerable variability in the frequency of auditory hallucinations in people with schizophrenia, typically ranging from once or twice weekly to

continuous.

Ref: Larøi F et al. The characteristic features of auditory verbal hallucinations in clinical and nonclinical groups: state-of-the-art overview and future directions. Schizophrenia bulletin. 2012 Jul 1;38(4):724-33.

The correct answer is: Present throughout the day incessantly

Question **54**

Not answered

Marked out of 1.00

Flag question

The ratio between number of different words used during a discourse and total number of spoken words is reduced in a patient with schizophrenia. What is the term for this?

Select one:

- Word count
- Cloze ratio
- Cohesion ratio
- Type-token ratio
- None of the listed options

Explanation:

The type-token ratio is a measure for lexical diversity. It is a ratio of the number of words used in the sample (# of types): all words in the sample (# of tokens).

Ref: de Boer JN et al. Language disturbances in schizophrenia: the relation with antipsychotic medication. NPJ schizophrenia. 2020 Sep 7;6(1):1-9.

The correct answer is: Type-token ratio

Question **55**

Not answered

Marked out of 1.00

Flag question

Which of the following is not a mixed affective state?

Select one:

- Depressive mania
- Inhibited mania
- Depression with poverty of thought
- Excited/agitated depression
- Manic stupor

In mixed states incongruity between motor features and mood state is found: mania with inhibition, depression with excitation etc.

The correct answer is: Depression with poverty of thought

Question **56**

Not answered

Marked out of

A woman states that she is often woken up when drifting to sleep by a voice saying a sentence or phrase that has no discoverable meaning. Which one of the following statements about her experience is true?

1.00

Flag question

Select one:

- It does not occur in organic states
- It can occur in the absence of a diagnosable mental illness
- EEG is likely to show beta rhythm
- Such experiences can occur only in the auditory modality
- It is a sign of nocturnal epilepsy

Explanation:

A hypnagogic hallucination is a transient false perception experienced while on the verge of falling asleep. The same phenomenon experienced while waking up is called a hypnopompic hallucination. These may have the characteristics of true or pseudo-hallucinations and are most commonly visual or auditory. While they are sometimes seen in narcolepsy or affective illnesses, they are not indicative of ill health and are frequently reported by healthy people.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 73, 112.

The correct answer is: It can occur in the absence of a diagnosable mental illness

Question **57**

Not answered

Marked out of 1.00

Flag question

A psychiatrist attempts to move a patient's arm. Before doing so, he instructs the patient to resist moving and not to let him manipulate his arm. But the patient continues to move his arm in the direction of the force. Once the psychiatrist removes the application of force, the patient's arm comes back to the original position. This phenomenon is called

Select one:

- Grimacing
- Posturing
- Negativism
- Mitgehen
- Automatic obedience

Mitgehen refers to a form of extreme cooperation in which the patient moves their body in the direction of the slightest pressure on the part of the examiner. For example, the doctor puts his forefinger under the patient's arm and presses gently, after that the arm moves upwards in the direction of the pressure. Once the pressure stops, the arm returns to its former position. Light pressure on the occiput of the patient, who is standing, leads to bending of the neck, flexing of the trunk and, if the pressure continues, the patient may fall forward.

The correct answer is: Mitgehen

Question **58**

Not answered

Marked out of 1.00

The most common type of seizure with an aura is

Select one:

- Pseudoseizures

Flag question

- Grand mal seizure
- Absence seizure
- Complex partial seizure
- Parietal seizure

Explanation:

Preictal events (auras) in complex partial seizures include autonomic sensations (e.g., fullness in the stomach, blushing, and changes in respiration); cognitive sensations (e.g., déjà vu, jamais vu, forced thinking, dreamy states); affective states (e.g., fear, panic, depression, elation); and classically, automatisms (e.g., lip smacking, rubbing, chewing).

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 251.

The correct answer is: Complex partial seizure

Question **59**

Not answered

Marked out of 1.00

Flag question

Which one among the following is not a motor symptom of schizophrenia?

Select one:

- Cataplexy
- Psychological pillow
- Negativism
- Ambitendency
- Catalepsy

Explanation:

Cataplexy is a symptom of narcolepsy in which there is sudden loss of muscle tone, leading to collapse. This usually occurs following emotional stress. Recognised motor symptoms of schizophrenia include catatonia, catalepsy, negativism, ambitendency, psychological pillow, mannerism, stereotypy, echopraxia, mitmachen, mitgehen and automatic obedience.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 104.

The correct answer is: Cataplexy

Question **60**

Not answered

Marked out of 1.00

Flag question

In formal thought disorder, asyndesis refers to

Select one:

- Lack of genuine causal links
- Lack of logical arguments
- Lack of wide vocabulary
- Lack of information
- Lack of adjectives

Explanation:

Asyndesis (also known as loosening of associations) is a symptom of formal thought disorder in which there is a lack of meaningful connection between sequential ideas. Cameron (1944), used the term 'asyndesis' to describe the lack of adequate connections or causal links between successive thoughts.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 103, 113.

The correct answer is: Lack of genuine causal links

Question **61**

Not answered

Marked out of 1.00

Flag question

Which of the following is associated with Ganser syndrome?

Select one:

- Automatisms
- Amnesia for behaviour
- Confabulation
- Anhedonia
- Accurate answers

Explanation:

Ganser syndrome is characterised by approximate answers, disorientation, clouding of consciousness, hallucinations/pseudo-hallucinations, motor disturbance, anxiety or apathy, normal ADLs, and sudden resolution with amnesia for the period of illness. In ICD-11, it is included under ‘other specified dissociative disorders’.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 236.

Dieguez S. Ganser syndrome. Neurologic-Psychiatric Syndromes in Focus-Part II. 2018;42:1-22.

The correct answer is: Amnesia for behaviour

Question **62**

Not answered

Marked out of 1.00

Flag question

A woman wakes up at 6am every morning, does 1 hour of vacuuming, prepares her list for the day and arrives at work 30 minutes before the start time. At work, she is always ahead of her schedule and always plans things well in advance. She expects her junior colleagues to be at her standard. What is the correct term to describe this scenario?

Select one:

- Compulsion
- Obsessive thinking
- Obsessional traits
- OCD
- Normal phenomena

Explanation:

One approach to separating 'normal' from 'pathological' requires that either distress or functional impairment be present in order to make a diagnosis. While DSM requires distress or functional impairment, ICD requires both. Unless there is some evidence of this, it must be considered as normal.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019.pp. 14-15.

The correct answer is: Normal phenomena

Question **63**

Not answered

Marked out of 1.00

Flag question

Automatism is a recognised phenomenon seen most commonly in patients with

Select one:

- Borderline personality disorder
- Hysterical dissociation
- Bipolar disorder
- Schizophrenia
- Temporal lobe epilepsy

Explanation:

An automatism is a behaviour which is apparently conscious in nature, occurring in the absence of full consciousness (e.g. during a temporal lobe seizure).

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 103.

The correct answer is: Temporal lobe epilepsy

Question **64**

Not answered

Marked out of 1.00

Flag question

Which of the following is NOT a feature associated with alexithymia?

Select one:

- Increase in abstract thinking ability
- Reduced symbolic thinking
- Difficulties in differentiating body sensations and emotional states
- Diminution of fantasy
- Difficulties in recognising one's own feelings

Explanation:

Patients with alexithymia know very little about their own feelings and, in most instances, are unable to link them with memories, fantasies, or specific situations. Salient features of alexithymia include:

- Difficulty identifying feelings

- Difficulty distinguishing between feelings and the bodily sensations of emotional arousal
- Difficulty describing feelings to other people
- Constricted imaginal processes, as evidenced by a paucity of fantasy
- A stimulus-bound, externally-oriented cognitive style

Ref: Goerlich KS. The multifaceted nature of alexithymia–a neuroscientific perspective. Frontiers in psychology. 2018 Aug 29;9:1614.

The correct answer is: Increase in abstract thinking ability

Question **65**

Not answered

Marked out of 1.00

Flag question

Which of the following is the commonest cause of stupor in psychiatric units?

Select one:

- Mania
- Epilepsy
- Dissociation
- Depression
- Stroke

Explanation:

Stupor is a state of more or less complete loss of activity where there is no reaction to external stimuli. It can be regarded as an extreme form of hypokinesia. Stupor may occur in states of shock, dissociative or conversion disorders, depression, psychosis, catatonia, and organic brain disease. The most common cause of stupor in psychiatric units is depression.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 98.

The correct answer is: Depression

Question **66**

Not answered

Marked out of 1.00

Flag question

A dominating but solitary belief that is neither delusional nor obsessional but influences one's life to a notable extent is called

Select one:

- Compulsion
- Stereotype
- Rumination
- Cognitive distortion
- Over-valued idea

Explanation:

An over-valued idea is a form of abnormal belief that is reasonable and understandable in itself, but which comes to unreasonably dominate the patient's life. It is a thought that, because of the associated feeling tone, takes precedence over all other ideas and maintains

this precedence permanently or for a long period of time. Over-valued ideas tend to be less fixed than delusions and tend to have some degree of basis in reality.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 115.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 48.

The correct answer is: Over-valued idea

Question **67**

Not answered

Marked out of 1.00

Flag question

Which of the following is NOT a first rank symptom of schizophrenia?

Select one:

- Voices giving repeated feedback as and when the patient does an act
- Voices commanding the patient
- Voices discussing among themselves
- Voices echoing patient's own thoughts
- Voices giving commentary on the patient

Explanation:

A command hallucination is an auditory hallucination of a commanding voice, instructing the patient towards a particular action. It is also known as a teleological hallucination and is not a first-rank symptom of schizophrenia. In first rank symptoms, patients may hear voices commenting on what they are doing, discussing it between themselves, or reporting the patient's own thoughts back to them.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 78-79, 104.

The correct answer is: Voices commanding the patient

Question **68**

Not answered

Marked out of 1.00

Flag question

Circumstantial speech is seen in all EXCEPT

Select one:

- Intellectual disability
- Mania
- Obsessive-compulsive personality disorder
- Dementia
- Broca aphasia

Explanation:

Circumstantial thinking is a disorder of the form of thought where irrelevant details and digressions overwhelm the direction of the thought process. This abnormality may be reflected in the resultant speech and can be seen in a number of conditions including mania and obsessive-compulsive personality disorder. In contrast, Broca aphasia (also called non-

fluent aphasia) is characterised by non-fluent speech and somewhat impaired repetition.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 104

Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 48.

The correct answer is: Broca aphasia

Question **69**

Not answered

Marked out of 1.00

Flag question

Dysmorphophobia is typically a/an

Select one:

- Delusional idea
- Cognitive distortion
- Cultural idea
- Obsessional idea
- Overvalued idea

Explanation:

Dysmorphophobia is a type of over-valued idea where the patient believes that one aspect of their body is abnormal or conspicuously deformed.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 109.

The correct answer is: Overvalued idea

Question **70**

Not answered

Marked out of 1.00

Flag question

Which of the following are the two stages in a delusional perception?

Select one:

- Normal perception and misinterpretation
- Delusional mood and delusional significance
- Normal perception and delusional significance
- Recall of abnormal memory and delusional significance
- Abnormal perception and delusional significance

Explanation:

A delusional perception is a primary delusion which is recalled as having arisen as a result of a perception (e.g. a patient who, on seeing two white cars pull up in front of his house, became convinced that he was therefore about to be wrongly accused of being a paedophile). The percept is a real external object, not a hallucinatory experience. A delusional perception is the attribution of a new meaning, usually in the sense of self-reference, to a normally perceived object. Schneider emphasised the importance of a delusional perception's 'two memberedness', as there is a link from the perceived object to the subject's perception of this object, and a second link to the new significance of the perception.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 107.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. pp. 48-49.

The correct answer is: Normal perception and delusional significance

Question **71**

Not answered

Marked out of 1.00

Flag question

Psychomotor retardation is a characteristic feature of

Select one:

- Dementia
- Schizophrenia
- Borderline personality disorder
- Depression
- Bulimia

Explanation:

Psychomotor retardation is characterised by decreased spontaneous movement and slowness in instigating and completing voluntary movement. It is usually associated with a subjective sense of actions being more of an effort and with subjective retardation of thought. It occurs in moderate to severe depressive illness.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 117.

The correct answer is: Depression

Question **72**

Not answered

Marked out of 1.00

Flag question

Musical hallucinations has been reported in all of the following except

Select one:

- Temporal lobe lesions
- Cochlear defects
- Schizophrenia
- Dominant parietal lobe lesions
- Sensory deprivation

Hearing loss or deafness is a very important factor associated with musical hallucinations. The hallucinatory experiences are more intense when the surrounding noise is low. Brain imaging studies of patients with musical hallucinations reveal a dysfunctional temporal cortex, especially on the left more than the right hemisphere.

The correct answer is: Dominant parietal lobe lesions

Question

73

Not answered

Marked out of 1.00

Flag question

Which one of the following is not a negative symptom of schizophrenia?

Select one:

- Anhedonia
- Attentional impairment
- Affective flattening or blunting
- Alexithymia
- Avolition

The prominent negative symptoms include affective flattening or blunting, attentional impairment, avolition-apathy (lack of initiative), anhedonia, asociality, alogia (lack of speech output).

The correct answer is: Alexithymia

Question **74**

Not answered

Marked out of 1.00

Flag question

Which one among the following is not a first rank symptom of schizophrenia?

Select one:

- Thought insertion
- Thought withdrawal
- Thought broadcasting
- Delusional perception
- Thought blocking

Thought blocking is not a first-rank symptom. First rank symptoms of schizophrenia are useful because they occur reasonably often in schizophrenia and more rarely in other disorders, and it is not too difficult to tell whether they are present or not. First rank symptoms can be categorised as follows:

- Auditory hallucinations (voices heard arguing, thought echo, running commentary)
- Delusions of thought interference (thought insertion, thought withdrawal, thought broadcasting)
- Delusions of control (passivity of affect, passivity of impulse, passivity of volitions, somatic passivity)
- Delusional perception (a primary delusion of any content that is reported by the patient as having arisen following the experience of a normal perception)

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 110.

The correct answer is: Thought blocking

Question **75**

Not answered

A hospital manager is presenting data to a trust board. He says 'murder of 2 hospitals' instead of 'merger of 2 hospitals'. This is an example of

Marked out of 1.00

Flag question

Select one:

- stock word
- logoclonia
- paragrammatism
- neologism
- parapraxis

Explanation:

Parapraxis is a faulty act, such as a slip of the tongue or the misplacement of an article. Freud ascribed parapraxes to unconscious motives.

Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1142.

The correct answer is: parapraxis

Question **76**

Not answered

Marked out of 1.00

Flag question

Which of the following correctly describes a hypnagogic hallucination?

Select one:

- Hallucinations on waking up from sleep
- Hallucinations when falling asleep
- Hallucinations that occur only in day time sleep
- Hallucinations that form a part of dreaming
- Hallucinations that induce a sleep-like state

Explanation:

A hypnagogic hallucination is a transient false perception experienced while on the verge of falling asleep. These may have the characteristics of true or pseudo-hallucinations and are most commonly visual or auditory. While they are sometimes seen in narcolepsy or affective illnesses, they are not indicative of ill health and are frequently reported by healthy people.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 73, 112.

The correct answer is: Hallucinations when falling asleep

Question **77**

Not answered

Marked out of 1.00

Flag question

A 48-year-old man who exhibits decreased speech stated that his thoughts were 'taken from me years ago by a parish council'. The most appropriate psychopathology described here is

Select one:

- Crowding of thoughts
- Retrospective falsification
- Thought diffusion
-

Delusional memory

Thought withdrawal

Explanation:

Thought withdrawal is the delusion that one's thoughts are being removed from one's mind by other people or forces. Schneider gave the description of a man who was experiencing thought withdrawal in the above format.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1144.

The correct answer is: Thought withdrawal

Question **78**

Not answered

Marked out of 1.00

Flag question

Psychopathology of near death experiences include all of the following except

Select one:

- Motor perseveration
- Autoscopic experience
- Depersonalisation
- Transcendental experience
- Out of body experience

Explanation:

Near-death experiences are often strikingly similar. They often involve an out-of-body experience of viewing one's own body (autoscopy) and overhearing conversations, feelings of peace, hearing a distant voice, entering a dark tunnel, leaving the body behind, meeting dead loved ones, witnessing beings of light, returning to life to complete unfinished business, and deep sadness on leaving this new dimension. Motor perseveration is not a part of this experience.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 848.

The correct answer is: Motor perseveration

Question **79**

Not answered

Marked out of 1.00

Flag question

Echolalia is characterised by

Select one:

- Automatic repetition of one's own speech
- Automatic repetition of the actions of others
- Automatic repetition of one's own actions
- Virtual absence of movement and speech but preserved consciousness
- Automatic repetition of another person's speech

Explanation:

Echolalia is the repetition of phrases or sentences spoken by others (e.g., the examiner). Echopraxia refers to the automatic repetition of visually perceived actions of others. Akinetic mutism is the virtual absence of movement and speech in the presence of full consciousness.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 109.

The correct answer is: Automatic repetition of another person's speech

Question **80**

Not answered

Marked out of 1.00

Flag question

When the word 'pink' is considered, linked words such as 'Barbie' and 'girls' get activated in brain. This is called

Select one:

- Syntax
- Indirect priming
- Secondary priming
- Direct priming
- Cohesion

Reaction times are decreased in response to words preceded by semantically related words ('cat'-'dog') in comparison with words preceded by semantically unrelated words ('bus'-'dog'), this is known as semantic priming effect. "When a concept node is activated by corresponding word stimulus, this activation then spreads through the network to connected nodes, falling off with decreasing relatedness. The degree to which one concept activates another is thus presumably proportional to their relatedness in the semantic network". Large priming effects for indirectly related words (those related only through at least one other concept, such as 'cat' and 'cheese,' mediated by 'mouse') have been observed in schizophrenia patients with thought disorder. This is called indirect priming. [Excerpts from Kiang et al., Am J Psychiatry 2008; 165:74-81]

The correct answer is: Direct priming

Question **81**

Not answered

Marked out of 1.00

Flag question

The phenomenon of deja vu is

Select one:

- Inappropriate familiarity to a new place or event
- Transient loss of sense of familiarity to a well accustomed place
- Inappropriate familiarity with a new thought
- Loss of sense of familiarity to a well known person
- Mixture of meaningless words

Explanation:

Déjà vu is a sense that events being experienced for the first time have been experienced

before. It is not strictly a disturbance of memory, but a problem with familiarity of places and events. It comprises the feeling of having experienced a current event in the past, although it has no basis in fact. The converse, jamais vu, is the knowledge that an event has been experienced before but is not presently associated with the appropriate feelings of familiarity. Deja entendu, the feeling of auditory recognition, and deja pense, a new thought recognised as having previously occurred, are related to deja vu, being different only in the modality of experience. These can be experienced by normal subjects as well as among those with temporal lobe epilepsy.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 105.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 69.

The correct answer is: Inappropriate familiarity to a new place or event

Question **82**

Not answered

Marked out of 1.00

Flag question

Astasia-abasia has been observed in

Select one:

- Dissociative fugue
- Conversion disorder
- Depressive pseudodementia
- Phobic anxiety disorder
- Schizophrenia

Explanation:

Astasia-abasia is a gait disturbance that has been seen in conversion disorder. It is a wildly ataxic, staggering gait accompanied by gross, irregular, jerky truncal movements and thrashing and waving arm movements. Patients with the symptoms rarely fall; if they do, they are generally not injured.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 455.

The correct answer is: Conversion disorder

Question **83**

Not answered

Marked out of 1.00

Flag question

A middle-aged woman suffers a panic attack and believes she is going to die. Which term could be used to describe this psychopathology?

Select one:

- Schemas
- Overvalued idea
- Obsessional rumination
- Thought monitoring
- Outcome fear

Outcome fear is a diagnostic feature of panic disorder. The bias in information processing that leads to outcome fear is termed as catastrophic misinterpretation.

The correct answer is: Outcome fear

Question **84**

Not answered

Marked out of 1.00

Flag question

According to Jaspers which of the following is true about primary delusions?

Select one:

- They occur before other symptoms
- They are un-understandable in terms of other mental experiences
- They are specific for schizophrenia
- They have prognostic significance in schizophrenia
- They are resistant to treatment

Explanation:

Jaspers maintained that primary delusions were typical of un-understandable phenomena, being immediately experienced by the patients as absolutely self-evident like an intuition or a revelation and accompanied by abnormalities in the form such as radical transformation or emergence of thinking. He designated primary delusions as delusion proper, from which secondary delusions evolve via conscious thought processes.

Ref: Hayashi N et al. Delusion progression process from the perspective of patients with psychoses: A descriptive study based on the primary delusion concept of Karl Jaspers. Plos one. 2021 Apr 27;16(4):e0250766.

The correct answer is: They are un-understandable in terms of other mental experiences

Question **85**

Not answered

Marked out of 1.00

Flag question

A 66 yr old patient diagnosed with schizophrenia is observed to have rigidity at rest but carries out voluntary movements normally. Which of the following is true?

Select one:

- He has extrapyramidal rigidity from antipsychotics
- He is a malingerer
- He has spasticity due to cerebrovascular pathology
- He does not have catatonia
- He is exhibiting catatonia

Explanation:

Catatonia is characterised by increased resting muscle tone that is not present on active or passive movement (in contrast to the rigidity associated with Parkinson disease and extra-pyramidal side effects).

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 104.

The correct answer is: He is exhibiting catatonia

Question **86**

Not answered

Marked out of 1.00

Flag question

All of the following are accepted as neologisms except

Select one:

- Personal jargon invented to describe the schizophrenic experience
- A phrase constructed with incorrect grammar
- A new word incorrectly built up with origins clearly understandable
- An accepted word used in an unusual way acquiring a new meaning
- Completely new word, the derivation of which cannot be understood

Explanation:

Neologisms may be new words that are constructed by the patient or ordinary words that are used in a new way. The term 'neologism' is usually applied to new word formations produced by individuals with schizophrenia. When an individual with schizophrenia produces a new word, it may be completely new and its derivation cannot be understood; it may be a distortion of another word; or it may be a word that has been incorrectly constructed by the faulty use of the accepted rules of word formation.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 57.

The correct answer is: A phrase constructed with incorrect grammar

Question **87**

Not answered

Marked out of 1.00

Flag question

Which of the following is characterised by patients maintaining unusual postures for long time with no resistance to a passive movement?

Select one:

- Cataplexy
- Advertence
- Spasticity
- Mitgehen
- Catalepsy

Explanation:

Mitmachen is a motor symptom of schizophrenia where the patient's limbs can be moved without resistance to any position. The limbs return to their resting state once the examiner lets go, in contrast with catalepsy (or waxy flexibility) where the limbs remain in their set positions for prolonged periods. Mitgehen is an extreme form of mitmachen where the patient's limbs can be moved to any position by very slight or fingertip pressure.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 114.

The correct answer is: Catalepsy

Question **88**

Not answered

Marked out of 1.00

Flag question

Overinclusive thinking is tested by using a

Select one:

- Verbal fluency test
- Proverb testing
- Similarities test
- Sorting test
- Cognitive estimates test

Explanation:

Abstract attitude is a term introduced by German psychiatrist Kurt Goldstein to denote the ability to use conceptual categories in order to classify things according to their attributes and to think symbolically rather than concretely. Goldstein's sorting test requires the respondent to sort items into categories according to colour, form, and material. It can be used to test overinclusion.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015. pp. 4, 318.

The correct answer is: Sorting test

Question **89**

Not answered

Marked out of 1.00

Flag question

On seeing a cat crossing the road, a cab driver starts thinking that the MI5 are about to arrest him soon. This is an example of

Select one:

- Delusional perception
- Somatic passivity
- Delusional misinterpretation
- Thought insertion
- Delusional memory

Explanation:

A delusional perception is a primary delusion which is recalled as having arisen as a result of a perception (e.g. a patient who, on seeing two white cars pull up in front of his house, became convinced that he was therefore about to be wrongly accused of being a paedophile). The percept is a real external object, not a hallucinatory experience.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 107.

The correct answer is: Delusional perception

Question **90**

Not answered

Marked out of 1.00

A patient can hear voices whenever the noise of water running through a tap is heard. This is called

Select one:

Flag question

- Reverse hallucination
- Synaesthesia
- Reflex hallucination
- Functional hallucination
- Extracampine hallucination

Explanation:

In a functional hallucination, an auditory stimulus causes a hallucination, but the stimulus as well as the hallucination is experienced. In other words, the hallucination requires the presence of another real sensation. (For example, a patient with schizophrenia first heard the voice of God as her clock ticked; later she heard voices coming from the running tap and voices coming from the chirruping of the birds.) Patients can distinguish both features from each other and crucially, the hallucination does not occur without the stimulus.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 35.

The correct answer is: Functional hallucination

Question **91**

Not answered

Marked out of 1.00

Flag question

Which one of the following statements regarding deja vu and jamais vu is false?

Select one:

- Deja vu is associated with inappropriate familiarity
- Deja vu may occur in normal individuals
- Deja vu can occur in schizophrenia
- Jamais vu is the sense that events being experienced for the first time have been experienced before
- Deja vu occurs more often in TLE with a right-sided focus

Explanation:

Déjà vu is a sense that events being experienced for the first time have been experienced before. Jamais vu is the sensation that events or situations are unfamiliar, although they have been experienced before. Both déjà vu and jamais vu can be everyday experiences, but also non-specific symptoms of a number of disorders, including temporal lobe epilepsy, schizophrenia, and anxiety disorders.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 105, 113.

The correct answer is: Jamais vu is the sense that events being experienced for the first time have been experienced before

Question **92**

Not answered

Marked out of 1.00

Which of the following is true regarding pseudo-hallucinations?

Select one:

- Cannot be seen in normal people

Flag question

- Seen in Ganser syndrome
- Insight is not retained
- Associated with amnesia for the object
- Occurs in external outer space

Explanation:

A pseudo-hallucination is a false perception that is perceived as occurring as part of one's internal experience, not as part of the external world. It may be described as having an 'as if' quality or as being seen with the mind's eye. Pseudo-hallucinations are a type of mental image that, although clear and vivid, lack the substantiality of perceptions. Pseudo-hallucinations are seen in full consciousness, known to be not real perceptions, and are located not in objective space but in subjective space (for example, inside the head). Like true hallucinations, they are involuntary. Pseudo-hallucinations can occur in all modalities of sensation and are described in psychotic, organic, and drug-induced conditions, as well as occasionally in 'normal' individuals. (The hallucinations of deceased spouses commonly described by widows and widowers may have a form of a pseudo-hallucination.)

Ganser syndrome is characterised by approximate answers, disorientation, clouding of consciousness, hallucinations/pseudo-hallucinations, motor disturbance, anxiety or apathy, normal ADLs, and sudden resolution with amnesia for the period of illness. In ICD-11, it is included under 'other specified dissociative disorders'.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 116, 236.

Dieguez S. Ganser syndrome. Neurologic-Psychiatric Syndromes in Focus-Part II. 2018;42:1-22.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 28.

The correct answer is: Seen in Ganser syndrome

Question **93**

Not answered

Marked out of 1.00

Flag question

Which one of the following is NOT an elementary hallucination?

Select one:

- Seeing flashes of light
- Hearing multiple buzzing sounds
- Unstructured flakes of images in a delirious patient
- Voices repeating the word 'go'
- Single tone of unclear sound

Explanation:

Hallucinations may be elementary and unformed, and experienced as simple noises, bells, or undifferentiated whispers. Elemental hallucinations are of very simple form (e.g. flashes of light or unstructured noises such as clicks and bangs). Elementary auditory hallucinations can occur in organic states and schizophrenia.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 30.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 109.

The correct answer is: Voices repeating the word 'go'

Question **94**

Not answered

Marked out of 1.00

Flag question

Which of the following is NOT a catatonic sign?

Select one:

- Posturing
- Flexibilitas cerea
- Mitgehen
- Negativism
- Essential tremor

Explanation:

Catatonia is a syndrome of primarily psychomotor disturbances characterised by the occurrence of several different symptoms including stupor; catalepsy/waxy flexibility/flexibilitas cerea; mutism; negativism; posturing; mannerisms; stereotypies; psychomotor agitation; grimacing; echolalia; echopraxia; ambitendency (alternating mixture of automatic obedience and negativism).

Essential tremor (6-12Hz) occurs at rest, with action and postural. It is most noticeable symmetrically in the upper limbs. It is not a catatonic sign.

Ref: Gaebel W et al. Changes from ICD-10 to ICD-11 and future directions in psychiatric classification. Dialogues in clinical neuroscience. 2020 Mar;22(1):7.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 132.

The correct answer is: Essential tremor

Question **95**

Not answered

Marked out of 1.00

Flag question

Obsessional slowness is common in which of the following type of obsessions?

Select one:

- Washing obsessions
- Obsession of symmetry
- Obsessions of contamination
- Cleaning obsessions
- Religious obsessions

An obsessive need for symmetry and exactness and a compulsion to order and arrange often results in slowness.

The correct answer is: Obsession of symmetry

Question **96**

Not answered

Marked out of 1.00

Flag question

A 78-year-old man hospitalized after a fall, repeatedly sees small angels flying around his head. Which of the following is least likely to be true in this case?

Select one:

- This phenomenon can occur due to anticholinergic toxicity
- This phenomenon is reported in delirium tremens
- The perception has occurred without appropriate stimulus
- This phenomenon is associated with organic disorders of the brain
- Presence of intact consciousness suggests a functional disorder than an organic mental illness

Explanation:

An hallucination is an internal percept without a corresponding external object. Auditory hallucinations, particularly of voices, are characteristic of schizophrenic illness, whjle visual hallucinations are characteristic of organic states. All varieties of visual hallucinations are found in organic states, but small animals and insects are most often hallucinated in delirium.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 112.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 32.

The correct answer is: Presence of intact consciousness suggests a functional disorder than an organic mental illness

Question **97**

Not answered

Marked out of 1.00

Flag question

Eidetic imagery is best classified as a

Select one:

- Vivid visual memory
- Pseudohallucination
- Affect illusion
- Hallucination
- Complete illusion

Explanation:

Eidetic imagery is a particular type of exceptionally vivid visual memory. It is not a hallucination. It is more common in children than adults.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 109.

The correct answer is: Vivid visual memory

Question **98**

A special ability of memory in which visual images are drawn from memory accurately at will

Not answered

Marked out of 1.00

Flag question

and described as if being perceived currently is noted in some people. Which of the following is true about this?

Select one:

- It is common in depression
- Common in the elderly
- It is a type of hallucination
- Seen in some religious practices
- Always part of pathological experience

Eidetic images represent visual memories of almost hallucinatory vividness that are found in disorders due to substance misuse, especially hallucinogenic agents such as LSD and mescaline. It is seen more commonly in children and those involved in group religious practices.

The correct answer is: Seen in some religious practices

Question **99**

Not answered

Marked out of 1.00

Flag question

Schnauzkrampf is associated with

Select one:

- Dementia
- Schizophrenia
- Bipolar disorder
- Depression
- None of the listed options

Explanation:

The muscle tension in catatonic stupor (most commonly seen in schizophrenia) is usually increased in the muscles of the forehead and the masseters. The lips may be thrust forward in a tubular manner known as 'snout spasm' (*Schnauzkrampf).* Although this is obviously a disorder of expression, it is best regarded as a stereotyped posture.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019.pp. 88, 98.

The correct answer is: Schizophrenia

Question **100**

Not answered

Marked out of 1.00

Flag question

Of various thought disorders classified by Andreasen, which of the following is more common in schizophrenia than mania?

Select one:

- Circumstantiality
- Neologisms
- Flight of ideas

- Clanging
- Derailment

Explanation:

Nancy Andreasen created a scale in which the classic formal thought disorders are subdivided into three groups:

- **Communication disturbances:** when the speaker does not meet the necessary requirements for the listener's understanding (poverty of content of speech, pressure of speech, distractible speech, tangentiality, derailment, stilted speech, echolalia, self-reference, circumstantiality, loss of goal, perseveration, and blocking)
- **Language disturbances:** when the speaker violates the semantic and syntactic conventions (incoherence, clang association, neologisms, use of word approximations)
- **Thought disturbances:** when only thinking alone seems affected (poverty of thought and illogically aberrant inferential processes)

Andreasen demonstrated that these psychopathological findings were not specific to schizophrenia and were also common in other mental disorders (e.g. mania). Of those listed, derailment (synonymous with loose associations - a symptom of schizophrenic thought disorder in which there is a total break in the chain of association between the meaning of thoughts) is more common in schizophrenia than mania.

Ref: Jerónimo J et al. Formal Thought Disorders–Historical Roots. Frontiers in psychiatry. 2018 Nov 9;9:572.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 108.

Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 10.

The correct answer is: Derailment

Question **101**

Not answered

Marked out of 1.00

Flag question

Which one of the following statements about delusional mood is incorrect?

Select one:

- Delusional mood is same as delusional atmosphere
- The patient feels profoundly apprehensive and uncomfortable
- Delusional mood is often followed by a fully formed delusion
- Delusional mood is characteristically associated with perplexity
- Delusional mood is a first rank symptom of schizophrenia

Explanation:

Delusional mood (also called delusional atmosphere) is a primary delusion, but not a first-rank symptom. It is recalled as arising following a period when there is an abnormal mood state characterised by anticipatory anxiety, a sense of 'something is about to happen' and an increased sense of the significance of minor events. The development of the formed delusion may come as a relief to the patient in this situation.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 106-107.

The correct answer is: Delusional mood is a first rank symptom of schizophrenia

Question **102**

Not answered

Marked out of 1.00

Flag question

A 33-year-old patient with psychotic depression spilt all the contents of a boiling sauce pan by inverting it upside down. When asked about this event, she said 'It came to me from the Polish Embassy. It was nothing to do with me; they wanted it so I picked up the pan and poured it.' She is describing

Select one:

- Made act
- Made affect
- Delusional perception
- Perseveration
- Made impulse

She owns up the action, but not the impulse behind it. Hence, this is made impulse, not made act (Excerpt from TABLE 1. Examples of positive symptoms of schizophrenia, http://www.nature.com/nrn/journal/v10/n1/fig_tab/nrn2536_T1.html (accessed April 11, 2015).).

The correct answer is: Made impulse

Question **103**

Not answered

Marked out of 1.00

Flag question

Which one of the following statements about illusions is false?

Select one:

- Illusions can be brought on by sensory deprivation
- Visual illusions may be influenced by the prevailing emotional tone
- Both affect and complete illusions resolve on closer attention
- Illusions occur when the perception threshold is increased
- Illusions can be normal experiences

Explanation:

An illusion is a type of false perception in which the perception of a real-world object is combined with internal imagery to produce a false internal percept. Three types are recognised: affect, completion, and pareidolic illusions. In affect illusion, there is a combination of a heightened emotion and misperception. Completion illusions rely on our brain's tendency to 'fill in' presumed missing parts of an object to produce a meaningful percept and are the basis for many types of optical illusions. Both these types of illusions resolve on closer attention. Pareidolic illusions are meaningful percepts produced when experiencing a poorly defined stimulus (e.g. seeing faces in a fire or clouds). Illusions in themselves are not indicative of psychopathology since they can occur in the absence of psychiatric disorder.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 112-113.

th

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4 Edition. 2019. p. 26.

The correct answer is: Illusions occur when the perception threshold is increased

Question **104**

Not answered

Marked out of 1.00

Flag question

The commonest psychiatric diagnosis reported in patients with Capgras syndrome is

Select one:

- Psychotic depression
- Schizophrenia
- Alzheimer dementia
- Depression
- Bipolar disorder

Explanation:

Capgras syndrome is a type of delusional misidentification in which the patient believes that a person known to them has been replaced by a 'double' who is to all external appearances identical but is not the 'real person'. This syndrome can occur both in psychiatric disorders, mainly psychotic disorders (especially schizophrenia) as well as major affective disorders, and neurocognitive disorders (e.g., post-traumatic encephalopathy, epilepsy, cerebrovascular disease, dementia, and delirium/acute confusional state).

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 104.

Karakasi MV et al. In fear of the most loved ones. A comprehensive review on Capgras misidentification phenomenon and case report involving attempted murder under Capgras syndrome in a relapse of a schizophrenia spectrum disorder. Journal of forensic and legal medicine. 2019 Aug 1;66:8-24.

The correct answer is: Schizophrenia

Question **105**

Not answered

Marked out of 1.00

Flag question

A patient, who lost his right arm amputated following a crush injury, suffers from recurrent sensations arising out of the lost limb. This experience is a

Select one:

- Delusion
- Body image disturbance
- Hallucination
- Pseudohallucination
- Somatisation

In most phantom limbs the phenomenon is produced by peripheral and central disorders. Hence it is improper to term this as a hallucination (which in strict sense must occur in the absence of a stimulus). The phantom limb does not necessarily correspond to the previous image of the limb in that it may be shorter or consist only of the distal portion so that the

phantom hand arises from the shoulder. It is best described as a body image disturbance.

The correct answer is: Body image disturbance

Question **106**

Not answered

Marked out of 1.00

Flag question

Depersonalisation is also described as a/an

Select one:

- 'what if' phenomenon
- 'why me' phenomenon
- 'if not' phenomenon
- 'what next' phenomenon
- 'as if' phenomenon

Explanation:

Depersonalisation is an unpleasant subjective experience where the patient feels **as if** they have become ‘unreal’. It is a feeling that one is not oneself or that something has changed. Depersonalisation is a non-specific symptom occurring in many psychiatric disorders, as well as in ‘normal’ people.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 108.

Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 10.

The correct answer is: 'as if' phenomenon

Question **107**

Not answered

Marked out of 1.00

Flag question

Crowding of thoughts occurs mainly in

Select one:

- Schizophrenia
- Dementia
- Mania
- Learning disability
- Depression

Crowding of thought occurs in schizophrenia. Here the patient describes his thoughts as being passively concentrated and compressed in his head. The associations are experienced as being excessive in amount, too fast, inexplicable and outside the person's control. Experientially, this is different from a manic flight of ideas.

The correct answer is: Schizophrenia

Question **108**

Not answered

Refraining from speech and making no attempt at spoken communication despite being fully conscious is called

Marked out of 1.00

Flag question

Select one:

- Mutism
- Stuttering
- Stammering
- Neologisms
- Vorbereiden

Explanation:

Mutism is the absence of speech without impairment of consciousness. Mutism may occur in children with a range of emotional or psychiatric disorders and in adults with hysteria, depression, schizophrenia, or organic brain disorders.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 115.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 56.

The correct answer is: Mutism

Question **109**

Not answered

Marked out of 1.00

Flag question

A 9-year-old girl can see the colour white associated with the number 7 and see the colour purple whenever she comes across the number 9. This is called

Select one:

- Functional hallucination
- Reflex hallucination
- Reverse hallucination
- Synaesthesia
- Extracampine hallucination

Explanation:

Synaesthesia is the experience of a stimulus in one sense modality producing a sensory experience in another (e.g. tasting sounds or hearing colours). Reflex hallucinations are a morbid form of synaesthesia. For example, a patient felt a pain in her head (somatic hallucination) when she heard other people sneeze (the stimulus) and was convinced that sneezing caused the pain.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 118.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 36.

The correct answer is: Synaesthesia

Question **110**

A patient says 'X-rays sent by my neighbour enter the back of my neck, where the skin

Not answered

Marked out of 1.00

Flag question

tingles and feels warm, they pass down the back in a hot wavy current about six inches to the sides of my hip bone.' This is best described as

Select one:

- Made volition
- Made impulse
- Synaesthesia
- Somatic passivity
- Kinaesthetic hallucination

Explanation:

Delusions of control are a group of delusions which are also known as passivity phenomenon or delusions of body passivity. They are considered first-rank symptoms of schizophrenia. The core feature is the delusional belief that one is no longer in sole control of one's own body. The individual delusions are that one is being forced by some external to feel emotions (passivity of affect or 'made affect'), to desire to do things (passivity of impulse or 'made impulse'), to perform actions (passivity of volition or 'made act'), or to experience bodily sensations (somatic passivity).

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 107.

The correct answer is: Somatic passivity

Question **111**

Not answered

Marked out of 1.00

Flag question

Which of the following is not a culture-bound disorder?

Select one:

- Bouffee delirante
- Koro
- Frigophobia
- Windigo
- Dysmorphophobia

Explanation:

Dysmorphophobia is a type of over-valued idea where the patient believes that one aspect of their body is abnormal or conspicuously deformed. It is not a culture-bound syndrome.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 109.

Tseng WS. From peculiar psychiatric disorders through culture-bound syndromes to culture-related specific syndromes. Transcultural psychiatry. 2006 Dec;43(4):554-76.

The correct answer is: Dysmorphophobia

Question **112**

"The thoughts of Elvis come into my mind. He treats my mind like a screen and flashes his

Not answered

Marked out of 1.00

Flag question

thoughts on to it like you flash a picture." This patient is experiencing

Select one:

- Thought blocking
- Thought insertion
- Thought broadcasting
- Obsessions
- Thought withdrawal

Explanation:

Thought insertion is the delusional belief that thoughts are being placed in the patient's head from the outside. Thought insertion is a first-rank symptom of schizophrenia.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 118.

The correct answer is: Thought insertion

Question **113**

Not answered

Marked out of 1.00

Flag question

A 42 yr old man is at a haste to catch the last train for the day to London. In his rush, he reads 'Swindon' as London and boards the wrong train. Which of the following has taken place?

Select one:

- Hallucination
- Pareidolic illusion
- Imagery
- Completion illusion
- Affect illusion

Explanation:

An illusion is a type of false perception in which the perception of a real-world object is combined with internal imagery to produce a false internal percept. Three types are recognised: affect, completion, and pareidolic illusions. In affect illusion, there is a combination of a heightened emotion and misperception. Completion illusions rely on our brain's tendency to 'fill in' presumed missing parts of an object to produce a meaningful percept and are the basis for many types of optical illusions. Both these types of illusions resolve on closer attention. Pareidolic illusions are meaningful percepts produced when experiencing a poorly defined stimulus (e.g. seeing faces in a fire or clouds).

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 112-113.

The correct answer is: Affect illusion

Question **114**

A patient complains that people come to know what she is thinking by looking into her eyes. This can be described as

Not answered

Marked out of 1.00

Flag question

Select one:

- Overvalued idea
- Thought broadcast
- Thought alienation
- Delusion of reference
- Thought withdrawal

A patient with delusions of reference believes that other people are talking about her, slandering or spying on her. At times, it may be difficult to ascertain whether the patient has delusions alone or if she has self-referential hallucinations that are associated with the delusions.

The correct answer is: Delusion of reference

Question **115**

Not answered

Marked out of 1.00

Flag question

A 35 year old man in your inpatient unit keeps his ears blocked with cotton and tape. He is disturbed by two persons discussing between themselves about every movement that he makes. They also discuss his emotions and plans. The most likely description is

Select one:

- Referential delusions
- 2nd person hallucinations
- 3rd person hallucinations
- Thought broadcast
- 1st person hallucinations

Explanation:

'Running commentary' is a type of third-person auditory hallucination that is a first-rank symptom of schizophrenia. The patient hears one or more voices providing a narrative of their current actions.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 117.

The correct answer is: 3rd person hallucinations

Question **116**

Not answered

Marked out of 1.00

Flag question

Ekbom syndrome is least likely to be related to

Select one:

- Cocaine use
- Dementia
- Autism
- Schizophrenia

- Alcohol withdrawal

Explanation:

Ekbom syndrome is a monosymptomatic delusional disorder where the core delusion is a delusion of infestation. Patients believe that their skin is infected with multiple tiny, mite-like animals. It is also seen in acute confusional states (particularly secondary to drug or alcohol withdrawal), in schizophrenia, in dementing illnesses, and as delusional elaboration of tactile hallucinatory experiences.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 107, 109.

The correct answer is: Autism

Question **117**

Not answered

Marked out of 1.00

Flag question

Illusions that are created out of vague sensory percepts by an admixture of fantasy thinking and vivid visual imagery are called

Select one:

- Pseudo-hallucinations
- Imagery
- Pareidolia
- Fantasy
- True hallucinations

Explanation:

Pareidolic illusions are meaningful percepts produced when experiencing a poorly defined stimulus (e.g. seeing faces in a fire or clouds). In pareidolia, vivid illusions occur without the patient making any effort. These illusions are the result of excessive fantasy thinking and a vivid visual imagery. They cannot therefore be explained as the result of affect or mindset, and so they differ from the ordinary illusion.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 113.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 27.

The correct answer is: Pareidolia

Question **118**

Not answered

Marked out of 1.00

Flag question

A 32-year-old shift worker presents with a history of headaches occurring in the mornings for a few months. His GP has been treating him for a diagnosis of tension headache. Which of the following features suggest a different pathology?

Select one:

- Pain relieved by analgesics
- The localisation of pain is vague
- Associated with low mood and anxiety

- Postural worsening of pain
- Pain worse with stress

Postural change may indicate intracranial pathology.

The correct answer is: Postural worsening of pain

Question **119**

Not answered

Marked out of 1.00

Flag question

Choose one of the following statements that is true about autochthonous delusions.

Select one:

- Autochthonous delusion is a type of secondary delusion
- Autochthonous delusions and wahneinfall are equivalent terms
- Autochthonous delusions may give rise to secondary delusions
- Retrospective delusions are the same as autochthonous delusions
- Autochthonous delusions arise as a result of auditory hallucinations

Explanation:

An autochthonous delusion is a primary delusion which appears to arise fully formed in the patient's mind without explanation. Primary delusions are the direct result of psychopathology, while secondary delusions can be understood as having arisen in response to other primary psychiatric conditions. If the patient is asked to recall the point when they became aware of the delusion and its significance to them, they may report that it arose 'out of the blue', which is an autochthonous delusion. Primary delusions such as autochthonous delusions may give rise to many secondary delusions that get tightly systematised later. For example, a patient who suddenly gets an autochthonous idea that Martians are invading his office may then start connecting normal behaviours of his colleagues to Martians and may form a systematised belief complex involving cameras, alien sign languages, UFOs etc.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 103, 106.

The correct answer is: Autochthonous delusions may give rise to secondary delusions

Question **120**

Not answered

Marked out of 1.00

Flag question

Which of the following factors predict a high persistence of delusional ideas?

Select one:

- Younger patients
- Having a diagnosis of schizophrenia
- Married patients
- Having a delusion of grandiosity
- Having single primary delusion

A diagnosis of schizophrenia is the best predictor of persistence of delusional beliefs.

The correct answer is: Having a diagnosis of schizophrenia

Question **121**

Not answered

Marked out of 1.00

Flag question

A patient who is on antipsychotic drugs is seen pacing around the ward but when questioned he described no subjective discomfort or restlessness in legs. Which of the following best describes his condition?

Select one:

- Chorea
- Agitation
- Akathisia
- Passivity
- Dyskinesia

Explanation:

Psychomotor agitation is a combination of psychic anxiety and excess and purposeless motor activity. It is a symptom common to many mental illnesses and is found in 'normal' individuals in response to stress.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 117.

The correct answer is: Agitation

Question **122**

Not answered

Marked out of 1.00

Flag question

Which one of the following statements about delusional perception is false?

Select one:

- It refers to a type of secondary delusion
- A normal perception is interpreted with delusional meaning and has immense personal significance
- It may be preceded by delusional mood
- It refers to a delusion that follows a normal perception
- It is one of the first rank symptoms of schizophrenia

Explanation:

A delusional perception is a primary delusion of any content that is reported by the patient as having arisen following the experience of a normal perception. If the patient is asked to recall the point when they became aware of the delusion and its significance to them, they will report that the delusional perception arose on seeing a normal percept (the percept is a real external object, not a hallucinatory experience). Delusional perception is a first-rank symptom of schizophrenia.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 78, 106-107.

The correct answer is: It refers to a type of secondary delusion

Question **123**

Not answered

Marked out of 1.00

Flag question

Reduced symbolic thinking is a feature of

Select one:

- Ambivalence
- Avolition
- Apathy
- Alexithymia
- Anhedonia

Explanation:

Patients with alexithymia know very little about their own feelings and, in most instances, are unable to link them with memories, fantasies, or specific situations. Salient features of alexithymia include:

- Difficulty identifying feelings
- Difficulty distinguishing between feelings and the bodily sensations of emotional arousal
- Difficulty describing feelings to other people
- Constricted imaginal processes, as evidenced by a paucity of fantasy
- A stimulus-bound, externally-oriented cognitive style

Ref: Goerlich KS. The multifaceted nature of alexithymia–a neuroscientific perspective. Frontiers in psychology. 2018 Aug 29;9:1614.

The correct answer is: Alexithymia

Question **124**

Not answered

Marked out of 1.00

Flag question

Alexithymia is common in those with

Select one:

- Psychosomatic disorders
- Dementia
- Psychotic disorders
- Mood disorders
- Personality disorders

Explanation:

Alexithymia is the inability to describe one’s subjective emotional experiences verbally. It may be a personality characteristic but is also associated with somatisation.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 102.

The correct answer is: Psychosomatic disorders

Question **125**

Not answered

Marked out of 1.00

Flag question

Which of the following statements with respect to pseudohallucinations is true?
Pseudohallucinations

Select one:

- Are under the control of subject's volition
- Are located in the inner subjective space
- Are experienced as real and concrete
- Are typically located in the outer objective space
- Are pathognomonic of personality disorders

Explanation:

A pseudo-hallucination is a false perception that is perceived as occurring as part of one's internal experience, not as part of the external world. It may be described as having an 'as if' quality or as being seen with the mind's eye. Pseudo-hallucinations are a type of mental image that, although clear and vivid, lack the substantiality of perceptions. Pseudo-hallucinations are seen in full consciousness, known to be not real perceptions, and are located not in objective space but in subjective space (for example, inside the head). Like true hallucinations, they are involuntary. Pseudo-hallucinations can occur in all modalities of sensation and are described in psychotic, organic, and drug-induced conditions, as well as occasionally in 'normal' individuals. (The hallucinations of deceased spouses commonly described by widows and widowers may have a form of a pseudo-hallucination.)

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 116.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 28.

The correct answer is: Are located in the inner subjective space

Question **126**

Not answered

Marked out of 1.00

Flag question

Stereotypy is characterised by

Select one:

- Mimicking examiner's movements and speech
- Maintenance of unchanged facial expression
- Repetitive, non-goal-directed motor activity
- Spontaneous maintenance of limb posture
- Odd purposeful movements

Explanation:

Stereotypy is a repetitive and bizarre movement which is not goal-directed (in contrast to mannerism, which is goal-directed). A stereotypy may have delusional significance to the patient and is sometimes seen in schizophrenia.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 118.

The correct answer is: Repetitive, non-goal-directed motor activity

Question **127**

Not answered

Marked out of 1.00

Flag question

A patient complaints of odd sensations such as 'electric heating' emanating from her knees. This can be described as

Select one:

- First rank symptom
- Somatic delusion
- Overvalued idea
- Somatic passivity
- Somatic hallucination

Explanation:

Somatic (or tactile) hallucinations are hallucinations of touch. Some patients experience cold winds blowing on them, sensations of heat, electric shocks, and sexual sensations.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 33.

The correct answer is: Somatic hallucination

Question **128**

Not answered

Marked out of 1.00

Flag question

Elderly women who develop Ekbom's delusion most commonly visit which of the following medical services before being referred to psychiatry?

Select one:

- Endocrinology
- Rheumatology
- Neurology
- Ophthalmology
- Dermatology

Explanation:

In psychiatry, Ekbom syndrome is a monosymptomatic delusional disorder where the core delusion is a delusion of infestation. Patients believe that their skin is infected with multiple tiny, mite-like animals. It is also seen in acute confusional states (particularly secondary to drug or alcohol withdrawal), in schizophrenia, in dementing illnesses, and as delusional elaboration of tactile hallucinatory experiences. NOTE: The term Ekbom syndrome has also been used to describe Restless Legs Syndrome.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 107, 109.

The correct answer is: Dermatology

Question **129**

Regarding pure word deafness, which one of the following statements is false?

Not answered

Marked out of 1.00

Flag question

Select one:

- Here a patient can speak fluently
- The patient cannot understand speech, even though hearing is unimpaired for other sounds
- It is also called as subcortical auditory dysphasia
- The patient can recognize the meaning of words
- The patient can read and write correctly

Explanation:

Pure word deafness (also called subcortical auditory dysphasia) denotes the inability to understand speech in the face of spared reading comprehension, writing, and speech production. Hearing remains intact for voices and other sounds, but there is an inability to recognise speech.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022.

Maffei C et al. Pure word deafness following left temporal damage: Behavioral and neuroanatomical evidence from a new case. Cortex. 2017 Dec 1;97:240-54.

The correct answer is: The patient can recognize the meaning of words

Question **130**

Not answered

Marked out of 1.00

Flag question

What does the term paranoia literally mean?

Select one:

- 'Grandiosity'
- 'Self reference'
- 'On suspicion'
- 'Besides mind'
- 'Inside mind'

Explanation:

Etymologically, ‘paranoia’ joins ‘para’ (beside) with ‘nous’ (mind), denoting an experience beside-the-mind. Paranoia is Greek for ‘by the side of the mind’.

Ref: Liebert, R.J. Beside-the-mind: an unsettling, reparative reading of paranoia. *Subjectivity* **10,** 123–145 (2017). https://doi.org/10.1057/s41286-016-0015-5

Casey P, Kelly B. Fish’s clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019.

The correct answer is: 'Besides mind'

Question **131**

Not answered

Marked out of 1.00

Who first considered the distinction between 'understanding' and 'explaining' as a core skill in psychiatric practice?

Select one:

Flag question

- Jaspers
- Durkheim
- Kraepelin
- Kretschmer
- Schneider

Explanation:

Karl Jaspers laid down the methodology of the phenomenological approach for describing in-depth patients' subjective experiences (phenomena). He defined methods for understanding such experiences and distinguished 'understanding' as a component that is different from 'explaining' an observed phenomenon.

Ref: Hayashi N et al. Delusion progression process from the perspective of patients with psychoses: A descriptive study based on the primary delusion concept of Karl Jaspers. Plos one. 2021 Apr 27;16(4):e0250766.

The correct answer is: Jaspers

Question **132**

Not answered

Marked out of 1.00

Flag question

Teichopsic hallucinations are typical of

Select one:

- Migraine
- Temporal lobe epilepsy
- Lacunar infarcts
- Sleep deprivation
- Parietal lobe lesions

Explanation:

Migraine with aura, previously termed 'classic migraine', affects only about 20% of migraine patients. The aura, which can represent almost any symptom of cortex or brain dysfunction, typically precedes the headache by up to 1 hour or accompanies its onset. The so-called teichopsia (from the Greek for town wall and vision, as coined by Airy in 1870), a slowly expanding luminous horseshoe-shaped zigzag perception, constitutes the most typical aura pattern.

Ref: Kaufman DM, Geyer HL, Milstein MJ, Rosengard JL. Kaufman's Clinical Neurology for Psychiatrists, 9th Edition. 2023. p. 185.

Vincent MB. Vision and migraine. Headache: The Journal of Head and Face Pain. 2015 Apr;55(4):595-9.

The correct answer is: Migraine

Question **133**

Not answered

Marked out of

Hallucinations in depressive psychosis are usually

Select one:

1.00

Flag question

- Gustatory
- Auditory
- Olfactory
- Visual
- Tactile

Explanation:

Auditory hallucinations are the most frequent type of hallucination in patients with psychotic major depression.

Ref: Gaudiano BA et al. Prevalence and clinical characteristics of psychotic versus nonpsychotic major depression in a general psychiatric outpatient clinic. Depression and anxiety. 2009 Jan;26(1):54-64.

The correct answer is: Auditory

Question **134**

Not answered

Marked out of 1.00

Flag question

Choose the incorrect association from the following list

Select one:

- Flight of ideas - mania
- Thought blocking - schizophrenia
- Circumstantiality - obsessive compulsive disorder
- Perseveration - frontal lobe syndrome
- Inhibition of thinking - depression

Explanation:

Circumstantial thinking is a disorder of the form of thought where irrelevant details and digressions overwhelm the direction of the thought process. This abnormality may be reflected in the resultant speech and can be seen in a number of conditions including mania and obsessive-compulsive personality disorder (not OCD).

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 104.

The correct answer is: Circumstantiality - obsessive compulsive disorder

Question **135**

Not answered

Marked out of 1.00

Flag question

The organic states associated with autoscopic hallucinations of seeing oneself in external space include all except

Select one:

- Parietal tumours
- Toxic infective states
- Temporoparietal lesions
- Epilepsy

- Frontal lobe dementia

Explanation:

Autoscopy (also known as 'phantom mirror image') is the experience of seeing a visual hallucination or pseudo-hallucination of oneself. The cause of autoscopic phenomenon is unknown, but a biological hypothesis is that abnormal, episodic activity in areas of the temporoparietal lobes is involved with the sense of self, perhaps combined with abnormal activity in parts of the visual cortex. It is an uncommon symptom reported in epilepsy, focal lesions/tumours (particularly in the parieto-occipital region), toxic infective states, schizophrenia, depression, anxiety, and dissociative disorders.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 103.

Anzellotti F et al. Autoscopic phenomena: case report and review of literature. Behavioral and Brain Functions. 2011 Dec;7(1):1-1.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 36.

The correct answer is: Frontal lobe dementia

Question **136**

Not answered

Marked out of 1.00

Flag question

Which of the following is not a type of sensory distortion?

Select one:

- Hyperacusis
- Hyperaesthesia
- Macropsia
- Dysmegalopsia
- Pseudo-hallucinations

Explanation:

Sensory distortions are changes in the perceived intensity or quality of a real external stimulus. They are associated with organic conditions and with drug ingestion or withdrawal. Examples include hyperacusis (experiencing sounds as abnormally loud), micropsia ('wrong end of the telescope' effect, perceiving objects as smaller and further away), macropsia (objects appears larger than they really are), dysmegalopsia (size and shape of objects is misperceived, sometimes called the 'Alice in Wonderland' effect), and hyperaesthesia (increased sensitivity to tactile stimulation).

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 117-118.

Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. pp. 1138-1141.

The correct answer is: Pseudo-hallucinations

Question **137**

Not answered

Which of the following is NOT a delusional misidentification syndrome?

Marked out of 1.00

Flag question

Select one:

- Fregoli syndrome
- Cotard syndrome
- Capgras syndrome
- Reduplicative paramnesia
- Syndrome of subjective doubles

Explanation:

Delusional misidentification is a delusional belief that certain individuals are not who they externally appear to be. Types of delusional misidentification syndromes include Capgras, Fregoli, Intermetamorphosis, Subjective doubles, Autoscopic, Reverse subjective double, and Reverse Fregoli.

Cotard syndrome is a presentation of psychotic depressive illness seen particularly in elderly people. There is a combination of severely depressed mood with nihilistic delusions and/or hypochondriacal delusions. The patient may state that they are already dead and should be buried, that their insides have stopped working and are rotting away, or that they have stopped existing altogether. It is not a delusional misidentification syndrome.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 105-106, 240.

The correct answer is: Cotard syndrome

Question **138**

Not answered

Marked out of 1.00

Flag question

Which one of the following is not a characteristic feature of transsexualism?

Select one:

- It is characterized by normal anatomical sex.
- The diagnosis is usually made after puberty.
- It is a delusional belief of mistaken sexual identity
- Distress is due to the inappropriateness of one's anatomical sex to perceived gender
- Subjects show a marked preoccupation with the wish to undergo surgical correction

Transsexualism occurs as an overvalued idea - not a delusion.

The correct answer is: It is a delusional belief of mistaken sexual identity

Question **139**

Not answered

Marked out of 1.00

Flag question

A 78-year-old patient is observed to be lying when asked what he had for breakfast or who visited him the day before. He is not embarrassed by this and does not appear to be jocular with regard to the lies; the lies are usually trivial bringing no harm to anyone. Which of the following is true?

Select one:

- He is having memory difficulties
- He is experiencing Ganser syndrome

- He is playing practical jokes intentionally
- He is exhibiting pseudologia fantastica
- He has an antisocial personality

Explanation:

Confabulation is the process of describing plausibly false memories for a period for which the patient has amnesia. Confabulation occurs in Korsakoff's syndrome, in dementing illnesses, and following alcoholic palimpsest.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 105.

The correct answer is: He is having memory difficulties

Question **140**

Not answered

Marked out of 1.00

Flag question

'It is not me who is unhappy, but they are projecting unhappiness into my brain. They also project upon me laughter for no reason.' This is called

Select one:

- Made impulse
- Made act
- Blunted affect
- Somatic passivity
- Made affect

Explanation:

Delusions of control are a group of delusions which are also known as passivity phenomenon or delusions of body passivity. They are considered first-rank symptoms of schizophrenia. The core feature is the delusional belief that one is no longer in sole control of one's own body. The individual delusions are that one is being forced by some external to feel emotions (passivity of affect or 'made affect'), to desire to do things (passivity of impulse or 'made impulse'), to perform actions (passivity of volition or 'made act'), or to experience bodily sensations (somatic passivity).

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 107.

The correct answer is: Made affect

Question **141**

Not answered

Marked out of 1.00

Flag question

A 34-year-old man with schizophrenia resists moving his limbs during a neurological examination. Despite repeated reassurances from the examining doctor, he resists yielding his limbs for passive manipulations. He is exhibiting

Select one:

- Mannerism
- Automatism
- Negativism

- Obstruction
- Borderline traits

Explanation:

Negativism is when the patient resists carrying out the examiner's instructions and his attempts to move or direct the limbs. It can be found in catatonia, severe intellectual disability, and dementia. Negativism may be passive when all interference is resisted and orders are not carried out, or it may be manifest as active or command negativism when the patient does the exact opposite of what they are asked to do, in a reflex way.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 115.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 94.

The correct answer is: Negativism

Question **142**

Not answered

Marked out of 1.00

Flag question

A patient at your clinic reports, 'When I heard the railway station master's whistle, I knew that there was a plot to attack me.' This is called

Select one:

- Referential delusion
- Secondary delusion
- Delusional mood
- Delusional perception
- Partial delusion

Explanation:

A delusional perception is a primary delusion which is recalled as having arisen as a result of a perception (e.g. a patient who, on seeing two white cars pull up in front of his house, became convinced that he was therefore about to be wrongly accused of being a paedophile). The percept is a real external object, not a hallucinatory experience.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 107.

The correct answer is: Delusional perception

Question **143**

Not answered

Marked out of 1.00

Flag question

A patient repeatedly refers to computers as 'mathscreen'. His speech exhibits numerous other disturbances. This can be termed as

Select one:

- Stammering
- Parapraxis
- Stereotypy

- Stock words
- Metonymy

Explanation:

Metonymy is a speech disturbance common in schizophrenia in which the affected person uses a word or phrase that is related to the proper one but is not the one ordinarily used (for example, the patient refers to losing the 'piece of string' of the conversation rather than the 'thread' of the conversation).

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1141.

The correct answer is: Metonymy

Question **144**

Not answered

Marked out of 1.00

Flag question

Which of the following is not a type of formal thought disorder described by Carl Schneider?

Select one:

- Derailment
- Fusion
- Substitution
- Omission
- Overinclusion

Explanation:

Carl Schneider claimed that five features of formal thought disorders could be identified:

- Derailment: the thought slides on to a subsidiary thought
- Substitution: a major thought is substituted by a subsidiary one
- Omission: senseless omission of a thought or part of it
- Fusion: heterogenous elements of thought are interwoven with each other
- Drivelling: a disordered intermixture of constituent parts of one complex thought

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 55.

The correct answer is: Overinclusion

Question **145**

Not answered

Marked out of 1.00

Flag question

Emotional lability is least likely to be due to

Select one:

- Frontal lobe damage
- Pseudo bulbar palsy
- Bipolar disorder
- Temporal lobe atrophy

- Delirium

Explanation:

Bilateral injury/atrophy of the temporal lobes can result in a syndrome similar to Kluver-Bucy syndrome, characterised by hypersexuality, placidity, a tendency to explore the environment with the mouth, inability to recognise the emotional significance of visual stimuli, and continually shifting attention. Complete experimental ablation of the temporal lobes appears to produce a uniform, bland reaction to the environment, possibly because of an inability to access memories.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. pp. 933-934.

The correct answer is: Temporal lobe atrophy

Question **146**

Not answered

Marked out of 1.00

Flag question

The commonest psychiatric diagnosis reported in patients with autoscopic hallucinations is

Select one:

- Depression
- Hysterical dissociative states
- Schizophrenia
- Dementia
- Bipolar disorder

It is also seen in normal people when they have extreme fatigue, where it is usually accompanied by impaired consciousness. Sims states that depression is the commonest psychiatric cause of autoscopy.

The correct answer is: Depression

Question **147**

Not answered

Marked out of 1.00

Flag question

A 7-year-old girl speaks very clearly and fluently with her friends at school but becomes mute when at home. This is called

Select one:

- Akinetic mutism
- Poverty of content of speech
- Elective mutism
- Poverty of speech
- Hysterical mutism

Elective mutism may occur in children who refuse to speak to certain people; for example, the child may be mute at school but speak at home.

The correct answer is: Elective mutism

Question **148**

Not answered

Marked out of 1.00

Flag question

A woman believes that the tennis player Roger Federer is in love with her. She has never seen him in person but believes that he is delivering romantic e-mails to her inbox in various disguised names. She is exhibiting

Select one:

- De Clerambault delusion
- Cotard delusion
- Morbid jealousy
- Capgras delusion
- Fregoli delusion

Explanation:

De Clerambault syndrome is a form of delusion of love. The patient, usually female, believes that another, higher status individual is in love with them and may make efforts to contact that person.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 105, 230.

The correct answer is: De Clerambault delusion

Question **149**

Not answered

Marked out of 1.00

Flag question

Which one of the following is not true about Folie a deux?

Select one:

- The person with the primary psychosis is usually the active member
- In folie, a deux the associate usually experiences a primary delusion
- The associate is usually in some way disadvantaged and dependent
- The commonest relationship among sufferers is that of twin sisters
- The delusions are usually persecutory in nature

Folie a deux is a term derived from French words meaning 'madness shared by two'. An individual develops a delusion similar to another individual who already has an established delusion; usually the two individuals have a close relationship (often sisters). In DSM-5, this is termed as Shared Psychotic Disorder.

Note: Folie a trios - madness shared by three. Folie a quatre - madness shared by four. Folie a famille - madness shared by the family. Folie a plusieurs - madness shared by many.

The correct answer is: In folie, a deux the associate usually experiences a primary delusion

Question **150**

Not answered

Marked out of

Which of the following syndromes and delusional types are not correctly associated?

Select one:

1.00

Flag question

- Morbid jealousy - delusions of marital infidelity
- De Clerambault syndrome - delusions of love
- Cotard syndrome - nihilistic delusions
- Ekbom syndrome - delusions of infestation
- Couvade syndrome - delusion of pregnancy in postpartum mothers

Explanation:

Othello syndrome (morbid jealousy): a monosymptomatic delusional disorder where the core delusion has the content of delusional jealousy

De Clerambault syndrome: a form of delusion of love in which the patient, usually female, believes that another, higher-status individual is in love with them

Cotard syndrome: a presentation of psychotic depressive illness seen particularly in elderly people where there is a combination of severely depressed mood with nihilistic delusions and/or hypochondriacal delusions

Ekbom syndrome: a monosymptomatic delusional disorder where the core delusion is a delusion of infestation

Couvade syndrome: a conversion symptom seen in partners of expectant mothers during their pregnancy; symptoms vary, but mimic pregnancy symptoms; NOT delusional in nature – the affected individual does not believe that they are pregnant

The correct answer is: Couvade syndrome - delusion of pregnancy in postpartum mothers

Question **151**

Not answered

Marked out of 1.00

Flag question

Which one of the following is a type of abnormal induced movement?

Select one:

- Tics
- Chorea
- Stereotypy
- Perseveration
- Tremors

Explanation:

Some abnormal induced movements can be regarded as the result of undue compliance on the part of the patient, while others may be interpreted as indicating rejection of the environment. For example, in automatic obedience the patient carries out every instruction regardless of the consequence. While stereotypy is a spontaneous abnormal movement, perseveration is an induced movement because it is a senseless repetition of a goal-directed action that has already served its purpose.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. pp. 92-93.

The correct answer is: Perseveration

Question **152**

Not answered

Marked out of 1.00

Flag question

La belle indifference is typically seen in

Select one:

- Multiple sclerosis
- Schizophrenia
- Conversion disorder
- Dissociative amnesia
- Dissociative fugue

Explanation:

(La) belle indifference is a surprising lack of concern for, or denial of, apparently severe functional disability. It is part of classical descriptions of hysteria and continues to be associated with operational descriptions of conversion disorder. It is also seen in medical illnesses (e.g. following a stroke) and is a rare and non-specific symptom of no diagnostic value.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 103.

The correct answer is: Conversion disorder

Question **153**

Not answered

Marked out of 1.00

Flag question

In pure word blindness which of the following is true?

Select one:

- Speech fluency is diminished
- Speech comprehension is lost
- Cannot write spontaneously
- Reading comprehension is impaired
- Writing comprehension is impaired

Explanation:

Pure word blindness (also known as ‘pure alexia’, ‘alexia without agraphia’, and ‘agnostic alexia’) is characterised by various degrees of impaired word reading with no deficits in the production or comprehension of oral language, and a preserved ability to write spontaneously or on dictation.

Ref: Martinaud O. Visual agnosia and focal brain injury. Revue neurologique. 2017 Jul 1;173(7-8):451-60.

The correct answer is: Reading comprehension is impaired

Question **154**

Not answered

Marked out of 1.00

A patient withdrawing from alcohol sees small Chinese soldiers marching on his carpet. This phenomenon is called

Select one:

Flag question

- Pseudohallucination
- Affective illusion
- Macropsia
- Micropsia
- Lilliputian hallucination

Explanation:

Lilliputian hallucinations are a type of visual hallucination in which the subject sees miniature people or animals. They are associated with organic states, particularly delirium tremens.

The correct answer is: Lilliputian hallucination

Question **155**

Not answered

Marked out of 1.00

Flag question

In logoclonia, the subject keeps repeating which of the following?

Select one:

- Last syllable
- Last word
- Examiner's question
- Recently hear word
- Last sentence

Explanation:

Logoclonia refers to meaningless repetition of a syllable, especially an end syllable of a word. It was first reported among individuals with dementia by Kraepelin in 1910 who described is as a severe linguistic dysfunction observed in the advanced stages of early-onset Alzheimer disease. Logoclonia has been described by several neurologists since then and has often been linked to Alzheimer disease. Palilalia and stuttering have disorders that are similar to those of logoclonia and have been associated with Parkinson disease and progressive supranuclear palsy. Palilalia, however, refers to the repetition of an entire word, but not a syllable. Stuttering is the involuntary disruption or blocking of speech with sound repetition, especially the initial consonants of a word, and, unlike logoclonia, stuttering is accompanied by tense and effortful utterance.

Ref: Nakagawa Y et al. Logoclonia might be a Characteristic of Logopenic Variant Primary Progressive Aphasia at an Advanced Stage: Potential Mechanisms Underlying Logoclonia. Journal of Alzheimer's Disease. 2019 Jan 1;70(2):515-24.

The correct answer is: Last syllable

Question **156**

Not answered

Marked out of 1.00

Flag question

Which of the following is true concerning depersonalisation?

Select one:

- It is experienced as an unpleasant state
- The subject feels that the world around him is unreal

- Multiple personalities emerge during the experience
- It is associated with approximate answers
- The subject cannot recall events later

Explanation:

Depersonalisation is an unpleasant subjective experience where the patient feels as if they have become ‘unreal’. It is a feeling that one is not oneself or that something has changed. Depersonalisation is a non-specific symptom occurring in many psychiatric disorders, as well as in ‘normal’ people.

Derealisation is an unpleasant subjective experience where the patient feels as if the world has become unreal. It is a feeling that one’s environment has changed in some strange way that is difficult to describe. Like depersonalisation, derealisation is also a non-specific symptom of a number of disorders.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 108.

Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 10.

The correct answer is: It is experienced as an unpleasant state

Question **157**

Not answered

Marked out of 1.00

Flag question

Flashback phenomenon is commonly reported after ingesting which of the following substances?

Select one:

- Cocaine
- Psychotropic medication
- Nicotine
- Hallucinogen
- Psychostimulant

Explanation:

Long after ingesting a hallucinogen, a person can experience a flashback of hallucinogenic symptoms. Flashbacks are spontaneous, transitory recurrences of the substance-induced experience. Most flashbacks are episodes of visual distortion, geometric hallucinations, hallucinations of sounds or voices, false perceptions of movement in peripheral fields, flashes of colour, trails of images from moving objects, positive afterimages and halos, macropsia, micropsia, time expansion, physical symptoms, or relived intense emotion.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. pp. 322-323.

The correct answer is: Hallucinogen

Question **158**

Not answered

Marked out of

Coenesthetic hallucinations are a type of

Select one:

1.00

Flag question

- Visual hallucination
- Olfactory hallucination
- Eidetic imagery
- Visceral hallucination
- Delusional belief

Explanation:

Coenesthesia is an awareness of one's own body. Somatic hallucinations with peculiar visceral, bodily, or tactile perceptions in the absence of sensorial input are coenesthetic hallucinations. Such hallucinations of pain and deep sensations are 'visceral' hallucinations. Some patients may complain of twisting and tearing pains. These may be very bizarre when the patient complains that their organs are being torn out or the flesh ripped away from the body.

Ref: Nasser A et al. A case of coenesthetic hallucinations treated with low-frequency repetitive transcranial magnetic stimulation. Brain Stimulation: Basic, Translational, and Clinical Research in Neuromodulation. 2017 Jul 1;10(4):845-6.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 34.

The correct answer is: Visceral hallucination

Question **159**

Not answered

Marked out of 1.00

Flag question

Overvalued ideas are NOT noted in the core symptoms of which one of the following disorder?

Select one:

- Transsexualism
- Anorexia nervosa
- Body dysmorphic disorder
- PTSD
- Morbid jealousy

PTSD is an anxiety disorder; overvalued ideas are not described in the diagnostic criteria of PTSD

The correct answer is: PTSD

Question **160**

Not answered

Marked out of 1.00

Flag question

One-way amnesia is associated with

Select one:

- Histrionic personality disorder
- Schizotypal personality diso
- Multiple personality disorder
- Borderline personality disorder

- Anankastic personality disorder

In multiple personality disorder, one-way amnesia is reported. One-way amnesia refers to the phenomenon where in the personality A knows B's existence, while B is not aware of A's life.

The correct answer is: Multiple personality disorder

Question **161**

Not answered

Marked out of 1.00

Flag question

In spite of being instructed not to pull his shirt up and show his abdomen, a patient does the same whenever a doctor comes to speak to him. Which of the following symptom is he exhibiting?

Select one:

- Automatism
- Echopraxia
- Negativism
- Mannerism
- Automatic obedience

In automatic obedience the patient carries out every instruction regardless of the consequence. To demonstrate this, Emil Kraepelin would ask the patient to put out their tongue and he would prick it with a pin; patients with automatic obedience continued to put their tongue out when asked to, although every time they did so their tongue was pricked

The correct answer is: Automatic obedience

Question **162**

Not answered

Marked out of 1.00

Flag question

Which of the following is false with regard to pseudohallucinations?

Select one:

- Patient may have insight about the absence of reality
- May occur in a subjective space (e.g. within the head)
- They may accompany true hallucinations
- Pathognomonic of borderline personality disorder
- Patient does not usually search for the bearer of the voice

Explanations:

Pseudo-hallucinations are a type of mental image that, although clear and vivid, lack the substantiality of perceptions. They are experienced in full consciousness, known to be not real perceptions, and are located not in objective space but subjective space (for example, inside the head). Pseudo-hallucinations can occur in all modalities of sensation and are described in psychotic, organic, and drug-induced conditions, as well as occasionally in 'normal' individuals.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th

Edition. 2019. p. 28.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 116.

The correct answer is: Pathognomonic of borderline personality disorder

Question **163**

Not answered

Marked out of 1.00

Flag question

Auditory hallucinations with clear consciousness are least likely to be due to

Select one:

- Temporal lobe epilepsy
- Alzheimer disease
- Cocaine abuse
- Alcohol abuse
- Schizophrenia

Alcoholic hallucinosis, any functional psychiatric disorder, and early dementia can cause auditory hallucinations with clear consciousness.

The correct answer is: Temporal lobe epilepsy

Question **164**

Not answered

Marked out of 1.00

Flag question

Which of the following is a thought disorder where figure-ground differentiation apparently fails?

Select one:

- Neologism
- Concrete thinking
- Tangentiality
- Circumstantiality
- Vorbeireden

Circumstantiality occurs when thinking proceeds slowly with many unnecessary and trivial details, but finally the point is reached. The goal of thinking is never completely lost and thinking proceeds towards it by an intricate and convoluted path, and is often due to the loss of figure-ground differentiation in the thought process (From Fish's Psychopathology. 3rd ed).

The correct answer is: Circumstantiality

Question **165**

Not answered

Marked out of 1.00

Flag

A man shouts 'Biance' every time he sees a pretty woman. Which of the following best describes this phenomenon?

Select one:

- Compulsive act

question

- Obsessive image
- Obsessive impulse
- Mental compulsion
- Rumination

Explanation:

A compulsion is a behaviour or action that is recognised by the patient as unnecessary and purposeless, but that they cannot resist performing repeatedly. It is an obsessional motor act. Compulsive acts may not be preceded by an identifiable obsession, especially in disorders other than OCD such as autism, dementia, Tourette, etc. where they can be seen without preceding obsessions.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 104.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 46.

The correct answer is: Compulsive act

Question **166**

Not answered

Marked out of 1.00

Flag question

A 64-year-old patient admitted in a stroke ward bursts out to laughter or tears within minutes with no control over these emotions. What is the psychopathology seen in this case?

Select one:

- Depression
- Mania
- Mixed affective state
- Grief reaction
- Emotional incontinence

Explanation:

Emotionalism (or emotional incontinence) is characterised by an increase in emotional behaviour (crying or, less commonly, laughing) following minimal provoking stimuli. Patients may persistently cry or laugh in unexpected situations.

Ref: Bowen A et al. Royal College of Physicians 2016 National clinical guideline for stroke. RCP.

The correct answer is: Emotional incontinence

Question **167**

Not answered

Marked out of 1.00

Flag question

A male patient with a diagnosis of schizophrenia says 'I had a hysterectomy at age 3 and since then I became a man'. This can be described as which of the following symptoms?

Select one:

- Delusional misinterpretation
- Delusional memory

- Pseudologia fantastica
- Delusional perception
- Confabulation

Explanation:

A delusional memory is a primary delusion that is recalled as arising as a result of a memory.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 106.

The correct answer is: Delusional memory

Question **168**

Not answered

Marked out of 1.00

Flag question

Which one of the following statements about first rank symptoms of schizophrenia is NOT true?

Select one:

- They carry prognostic significance
- They are also seen in organic psychoses
- These are a group of symptoms which are useful in the diagnosis of schizophrenia
- They are also seen in affective psychoses
- They are neither pathognomonic nor specific to schizophrenia

Explanation:

First rank symptoms of schizophrenia are a group of symptoms that have special significance in the diagnosis of schizophrenia. They are useful because they occur reasonably often in schizophrenia and more rarely in other disorders, and it is not too difficult to tell whether they are present or not. However, there is no symptom that is pathognomonic of schizophrenia. First rank symptoms can all be reported in other conditions (e.g. organic psychoses, manic illnesses). They do not give a guide to severity or prognosis of illness, and they may not occur at all in a patient who undoubtedly has schizophrenia.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 78.

The correct answer is: They carry prognostic significance

Question **169**

Not answered

Marked out of 1.00

Flag question

Which one of the following is a type of catatonic phenomenon?

Select one:

- Aphonia
- Waxy flexibility
- Dyskinesia
- Oculogyric crisis
- Akathisia

Explanation:

Catatonia is a syndrome of primarily psychomotor disturbances characterised by the occurrence of several different symptoms including stupor; catalepsy/waxy flexibility/flexibilitas cerea; mutism; negativism; posturing; mannerisms; stereotypies; psychomotor agitation; grimacing; echolalia; echopraxia; ambitendency (alternating mixture of automatic obedience and negativism).

Waxy flexibility (also known as catalepsy or flexibilitas cerea) is a rare motor symptom in which the patient's limbs can be passively moved to any posture, which will then be held for a prolonged period of time.

Ref: Gaebel W et al. Changes from ICD-10 to ICD-11 and future directions in psychiatric classification. Dialogues in clinical neuroscience. 2020 Mar;22(1):7.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 104.

The correct answer is: Waxy flexibility

Question **170**

Not answered

Marked out of 1.00

Flag question

A chronic schizophrenia patient snouts his lips and maintains this posture for a long time even when he is active doing other work. Which of the following best describes the above sign?

Select one:

- Catalepsy
- Snout spasm
- Cataplexy
- Posturing
- Mannerism

In catatonia the lips may be thrust forward in a tubular manner known as 'snout spasm' (or schnauzkrampf) and although this is obviously a disorder of expression it is best regarded as a stereotyped posture.

The correct answer is: Snout spasm

Question **171**

Not answered

Marked out of 1.00

Flag question

Which of the following is true about first rank symptoms of schizophrenia?

Select one:

- They are essential for diagnosing schizophrenia
- They represent disturbance of ego-boundary
- They are not seen in other psychiatric disorders
- They are specific for schizophrenia
- They emphasize on content rather than form

Explanation:

First rank symptoms (FRS) of schizophrenia are a group of symptoms that have special significance in the diagnosis of schizophrenia. They are useful because they occur reasonably often in schizophrenia and more rarely in other disorders, and it is not too difficult to tell whether they are present or not. However, there is no symptom that is pathognomonic of schizophrenia. FRS can all be reported in other conditions (e.g. organic psychoses, manic illnesses). They do not give a guide to severity or prognosis of illness, and they may not occur at all in a patient who undoubtedly has schizophrenia. The original psychoanalytic interpretation of FRS is that the boundary between the ego and the surrounding world breaks down, leading to thought and volition being alienated.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 78.

The correct answer is: They represent disturbance of ego-boundary

Question **172**

Not answered

Marked out of 1.00

Flag question

A man with psychotic symptoms said: 'When I try to think, my head gets full of noise; it is like as if my brain were in an uproar with my thoughts'. The most appropriate psychopathology is

Select one:

- Thought diffusion
- Thought broadcast
- Thought provocation
- Thought echo
- Crowding of thoughts

Schneider described a man with schizophrenia who had the experience of his thoughts being spoken aloud, i.e. Thought Echo in the above fashion.Schizophr Bull, January 1, 2008; 34(1): 137 - 154

The correct answer is: Thought echo

Question **173**

Not answered

Marked out of 1.00

Flag question

'If I think of anything, at once those opposite to me know it and it is embarrassing'. This is best termed as

Select one:

- Referential delusion
- Delusional misinterpretation
- Thought diffusion
- Thought blocking
- Thought echo

Explanation:

Thought broadcasting (also known as ‘thought diffusion’) is the delusional belief that one’s thoughts are accessible directly to others. It is a first-rank symptom of schizophrenia. This

description here was given by a shopkeeper with schizophrenia and was reported by Schneider himself.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 118.

The correct answer is: Thought diffusion

Question **174**

Not answered

Marked out of 1.00

Flag question

Which of the following can differentiate depersonalisation seen in normal persons and those with a depersonalisation-related psychiatric problem?

Select one:

- No difference is notable between the two groups
- Shorter duration in the latter
- Intense affective change is seen in the latter
- Feelings of detachment are not seen in the former
- Sense of time is not altered in the former

Explanation:

Depersonalisation in psychiatric illness is characterised as distressing and anxiety-provoking.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 406.

The correct answer is: Intense affective change is seen in the latter

Question **175**

Not answered

Marked out of 1.00

Flag question

A 35 yr old woman is on sodium valproate for many years. She is having lack of sleep following a holiday and tells her husband 'Life is not a game, every idiot has to earn a name, so work hard for fame, if not you will end up feeling shame'. She is exhibiting

Select one:

- Rhyming
- Metonymy
- Punning
- Pressured speech
- Neologisms

Explanation:

An example of flight of ideas associated with rhymes comes from a manic patient described by Fish. She was asked where she lived and she replied: 'Birmingham, Kingstanding; see the king he's standing, king, king, sing, sing, bird on the wing, wing, wing on the bird, bird, turd, turd.'

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 44.

The correct answer is: Rhyming

Question **176**

Not answered

Marked out of 1.00

Flag question

A partially sighted, 77-year-old lady reports seeing Peter Rabbit images in front of her eye. She has no psychotic features. Which of the following is true about the above symptom?

Select one:

- Her consciousness may be altered
- It cannot occur unless she is completely blind in at least one eye
- The rabbit will not disappear on sleeping
- She may retain insight
- The objects seen are usually static

Explanation:

Charles Bonnet syndrome (phantom visual images) is a condition in which complex visual hallucinations occur in the absence of any psychopathology and in clear consciousness. It is associated with either central or peripheral reduction in vision. The images may be static or in motion. They may be enjoyable or distressing. Patients may retain insight.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. pp. 29, 124.

The correct answer is: She may retain insight

Question **177**

Not answered

Marked out of 1.00

Flag question

Which of the following is a term used to describe the phenomenon of hearing one's own thoughts aloud?

Select one:

- Mitmachen
- Mitgehen
- Gegenhalten
- Gedankenlautwerden
- Word salad

Explanation:

Gedankenlautwerden (also known as 'thought echo' or 'echo de la pensee') is the experience of an auditory hallucination in which the content is the individual's current thoughts. It is a first-rank symptom of schizophrenia.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 118.

The correct answer is: Gedankenlautwerden

Question **178**

Which one of the following is not a type of primary delusion?

Not answered

Marked out of 1.00

Flag question

Select one:

- Delusional atmosphere
- Delusional memory
- Delusional misidentification
- Delusional perception
- Autochthonous delusions

The 4 types of primary delusions include delusional mood, delusional perception, delusional memory and autochthonous delusions

The correct answer is: Delusional misidentification

Question **179**

Not answered

Marked out of 1.00

Flag question

Which of the following cannot be considered as a component of descriptive psychopathology?

Select one:

- Defence mechanisms
- Understanding patient's experience
- Using terms to describe behaviour
- Establishing rapport with patient
- Empathetic approach

Explanation:

Descriptive psychopathology includes close observation of the patient's behaviour and empathic exploration of their subjective experience. The latter is called phenomenology. Defence mechanisms are best described under psychodynamic (explanatory) psychopathology.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 100.

The correct answer is: Defence mechanisms

Question **180**

Not answered

Marked out of 1.00

Flag question

"I am thinking about my mother, and suddenly my thoughts are sucked out of my mind by a phrenological vacuum extractor, and there is nothing in my mind." This is best described as

Select one:

- Thought blocking
- Thought withdrawal
- Thought insertion
- Thought broadcasting
- Thought echo

Explanation:

Thought withdrawal is the delusion that one's thoughts are being removed from one's mind by other people or forces.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1144.

The correct answer is: Thought withdrawal

Question **181**

Not answered

Marked out of 1.00

Flag question

Which of the following best describes paraschemazia?

Select one:

- Mixture of meaningless words
- Distortion of body image
- Inappropriate familiarity in a new place
- Distorted awareness of time
- Loss of emotional control following brain injury

Explanation:

Paraschemazia or distortion of body image is described as a feeling that parts of the body are distorted or twisted or separated from the rest of the body. This can occur in association with hallucinogenic use, with an epileptic aura, and with migraine on rare occasions.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019.p. 39.

The correct answer is: Distortion of body image

Question **182**

Not answered

Marked out of 1.00

Flag question

Retardation of thinking seen in depressive disorder is appropriately classified as which one of the following?

Select one:

- Disturbed form of thought
- Disturbed content of thought
- Disturbed control of speech
- Disturbed thought agency
- Disturbed stream of thought

Explanation:

With inhibition or slowing of thinking in depression, the train of thought is slowed down and the number of ideas and mental images that present themselves is decreased. This is a disturbance in the stream of thought.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th

Edition. 2019. p. 44.

The correct answer is: Disturbed stream of thought

Question **183**

Not answered

Marked out of 1.00

Flag question

'My thoughts leave my head in a type of mental ticker-tape. Everyone around me has only to pass the tape through their mind, and they know my thoughts.' This patient is describing

Select one:

- Thought broadcasting
- Thought withdrawal
- Thought echo
- Thought blocking
- Thought insertion

Explanation:

Thought broadcasting (also known as 'thought diffusion') is the delusional belief that one's thoughts are accessible directly to others. It is a first-rank symptom of schizophrenia.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 118.

The correct answer is: Thought broadcasting

Question **184**

Not answered

Marked out of 1.00

Flag question

Inability to verbalize one's strong emotions is termed as

Select one:

- Ambivalence
- Apathy
- Anhedonia
- Alexithymia
- Avolition

Explanation:

Alexithymia is the inability to describe one's subjective emotional experiences verbally. It may be a personality characteristic but is also associated with somatisation.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 102.

The correct answer is: Alexithymia

Question **185**

Not answered

Which of the following neurological disturbances closely mimic schizophrenic speech disturbance?

Marked out of 1.00

Flag question

Select one:

- Anomia
- Motor aphasia
- Jargon aphasia
- Non fluent aphasia
- Conduction aphasia

Explanation:

Jargon aphasia is a form of fluent aphasia characterised by a copious flow of unintelligible speech. Jargon-aphasic speech may be subdivided into semantic jargon, in which ordinary words are strung together to form unintelligible utterances, and phonemic jargon involving neologisms. A neologism is a made-up word or normal word used in an idiosyncratic way. Neologisms may be found in schizophrenic speech.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015. p. 396.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 115.

The correct answer is: Jargon aphasia

Question **186**

Not answered

Marked out of 1.00

Flag question

A 35-year-old lady in your inpatient unit covers her head with a veil. She claims that other people can come to know what she thinks. The most likely description is

Select one:

- Thought insertion
- Thought broadcasting
- Thought withdrawal
- Thought blocking
- Obsessions

Explanation:

Thought broadcasting (also known as 'thought diffusion') is the delusional belief that one's thoughts are accessible directly to others. It is a first-rank symptom of schizophrenia.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 118.

The correct answer is: Thought broadcasting

Question **187**

Not answered

Marked out of 1.00

Flag

A constellation of unchanging facial expression, decreased spontaneous movements, poverty of expressive gestures, poor eye contact and affective non-responsivity is termed as

Select one:

- Lability of affect

question

- Incongruity of affect
- Flattening of affect
- Reactive affect
- Blunting of affect

Explanation:

Flattening of affect is the diminution of the normal range of emotional experience and is a negative symptom of schizophrenia.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 111.

The correct answer is: Flattening of affect

Question **188**

Not answered

Marked out of 1.00

Flag question

Which of the following is not a catatonic sign?

Select one:

- Catalepsy
- Posturing
- Negativism
- Ambivalence
- Ambitendency

Explanation:

Catatonia is a syndrome of primarily psychomotor disturbances characterised by the occurrence of several different symptoms including stupor; catalepsy/waxy flexibility/flexibilitas cerea; mutism; negativism; posturing; mannerisms; stereotypies; psychomotor agitation; grimacing; echolalia; echopraxia; ambitendency (alternating mixture of automatic obedience and negativism).

Ref: Gaebel W et al. Changes from ICD-10 to ICD-11 and future directions in psychiatric classification. Dialogues in clinical neuroscience. 2020 Mar;22(1):7.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 102.

The correct answer is: Ambivalence

Question **189**

Not answered

Marked out of 1.00

Flag question

Which of the following is most likely to be seen in an 18-year-old girl with hysterical aphonia?

Select one:

- Vocal cord movements are restricted
- Patient loses ability to speak aloud but whispers
- Speech is absent
- There is difficulty in articulation

- Patient loses ability to cough

Dysarthria: There is difficulty in articulation, which occurs in lesions of the brain stem, schizophrenia and cerebellar disorder.

Dysphonia: Impairment of ability to vocalize with some hoarseness of voice, which occurs in the paralysis of the 9th cranial nerve and disease of vocal cords.

Aphonia: Patient loses ability to vocalise and whispers, occurs in conversion disorders. Aphonia is defined as the absence of vocalization or phonation and differs from mutism, which is the absence of speech. Hysterical aphonia is characterized by a normal whisper and cough. Examination with a laryngoscope reveals normal vocal cord movement with respiration.

The correct answer is: Patient loses ability to speak aloud but whispers

Finish review

PREVIOUS ACTIVITY
Clinical Examination

NEXT ACTIVITY
Dynamic Psychopathology

Jump to...

Drop here!

© 2022 SPMM Course Limited, UK.

Refund Policy | Terms & Conditions