Started on	Saturday, 23 July 2022, 11:21 AM
State	Finished
Completed on	Saturday, 23 July 2022, 11:21 AM
Time taken	8 secs
Grade	**0.00** out of 17.00 (**0**%)

Question **1**

Not answered

Marked out of 1.00

Flag question

Which of the following is a qualifier for the diagnoses under the disruptive and dissocial disorders grouping?

Select one:

- Emotional lability
- Childhood vs adult onset
- Criminal behaviour
- Limited prosocial emotions
- Self-injury

Your answer is incorrect.

The disruptive and dissocial disorders diagnostic grouping includes oppositional defiant disorder and conduct disorder. They can both receive a qualifier of 'limited prosocial emotions' that indicates a poorer long-term prognosis. Oppositional defiant disorder can receive a qualifier of 'with chronic irritability and anger', while conduct disorder can receive a qualifier of 'childhood vs adolescent onset'.

The correct answer is: Limited prosocial emotions

Question **2**

Not answered

Marked out of 1.00

Flag question

Which of the following is a qualifier, an additional diagnostic descriptor, that can be used for disorders under the obsessive-compulsive and related disorders grouping?

Select one:

- Prominent checking behaviours
- Without obsessive ideation
- Poor to absent insight
- Treatment resistant
- Without compulsive behaviours

Your answer is incorrect.

The poor to absent insight qualifier can be used when an individual's obsessive cognitions

are so intense as to be almost delusional in nature, but they are consistent with the existing obsessive-compulsive or related disorder diagnosis.

The correct answer is: Poor to absent insight

Question **3**

Not answered

Marked out of 1.00

Flag question

Which of the following disorders is included in the new diagnostic grouping of disorders of bodily distress and bodily experience in ICD-11?

Select one:

- Trichotillomania
- Body integrity dysphoria
- Hypochondriasis
- Body dysmorphic disorder
- Excoriation disorder

Your answer is incorrect.

The diagnostic grouping of disorders of bodily distress and bodily experience includes bodily distress disorder, which broadly replaces somatoform disorders and neurasthenia, and body integrity dysphoria, which is characterised by the desire to have a certain disability. The other diagnoses lie within the obsessive-compulsive and related disorders, with trichotillomania and excoriation disorder being within the body-focused repetitive behaviour disorders subgrouping.

The correct answer is: Body integrity dysphoria

Question **4**

Not answered

Marked out of 1.00

Flag question

Which of the following diagnostic groupings includes selective mutism in ICD-11?

Select one:

- Anxiety and fear-related disorders
- Neurodevelopmental disorders
- Dissociative disorders
- Obsessive-compulsive and related disorders
- Disorders specifically associated with stress

Your answer is incorrect.

Anxiety and fear-related disorders includes selective mutism and separation anxiety. Diagnoses are categorised by the focus of the apprehension for the individual, with this being diffuse in generalised anxiety disorder.

The correct answer is: Anxiety and fear-related disorders

Question **5**

Not answered

Body integrity dysphoria is a new inclusion in ICD-11. Which of the following is characteristic of this new disorder?

Marked out of 1.00

Flag question

Select one:

- The feeling that a body part is not truly part of one's body
- The sense that one's body is not real
- The desire to have a specific disability
- The sense that one's body is not functioning correctly
- The feeling that a certain body part is not under one's control

Your answer is incorrect.

Body integrity dysphoria is characterised by the desire to have a particular disability and to feel discomforted by not having this disability. It is usually thought to have onset during childhood or adolescence.

The correct answer is: The desire to have a specific disability

Question **6**

Not answered

Marked out of 1.00

Flag question

In ICD-11, which of the following diagnoses is still recognised in the schizophrenia and primary psychotic disorders diagnostic grouping?

Select one:

- Hebephrenic schizophrenia
- Schizoaffective disorder
- Simple schizophrenia
- Paranoid schizophrenia
- Residual schizophrenia

Your answer is incorrect.

The subtypes of schizophrenia have been eliminated from ICD-11, with dimensional descriptors used instead to provide information about the nature of a patient's symptoms, with the following symptom groups being used: positive, negative, depressive, manic, psychomotor, and cognitive symptoms. The same applies to other primary psychotic disorders. Schizoaffective disorder continues to be recognised in ICD-11.

The correct answer is: Schizoaffective disorder

Question **7**

Not answered

Marked out of 1.00

Flag question

Which of the following is a qualifier, an additional diagnostic descriptor, for depressive episodes in ICD-11?

Select one:

- Severe disturbance of functioning
- Catatonia
- Suicidal ideation
- Melancholic features

- Acute onset

Your answer is incorrect.

The qualifiers for depressive episodes in ICD-11 are melancholic features, anxiety symptoms, panic attacks, and seasonal pattern. Catatonia is contained within its own diagnostic grouping

The correct answer is: Melancholic features

Question **8**

Not answered

Marked out of 1.00

Flag question

Which of the following is a new disorder in ICD-11 under the obsessive-compulsive and related disorders grouping?

Select one:

- Body integrity dysphoria
- Avoidant/restrictive food intake disorder
- Olfactory reference disorder
- Gaming disorder
- Bodily distress disorder

Your answer is incorrect.

Olfactory reference disorder is characterised by the persistent belief that one is emitting an offensive body odour or breath, which others cannot perceive. Individuals perform repeated and excessive behaviours such as checking for the odour, seeking reassurance, trying to eliminate or cover up the smell, and avoiding social situations. Bodily distress disorder and body integrity dysphoria are grouped under disorders of bodily distress and bodily experience. Bodily distress disorder involves persistent bodily sensations that are distressing to the individual. Body integrity dysphoria is a rare disorder characterised by the persistent desire to have a specific physical disability. Avoidant/restrictive food intake disorder is categorised under feeding and eating disorders and replaces feeding disorder of infancy or childhood and refers to individuals who engage in abnormal behaviour around eating, not motivated by cognitions around body image, which leads to inadequate intake. Gaming disorder is categorised under disorders due to substance use and addictive behaviours.

The correct answer is: Olfactory reference disorder

Question **9**

Not answered

Marked out of 1.00

Flag question

Malingering is the falsification or exaggeration of illness, either physical or mental, for the purpose of secondary gain. Under which diagnostic grouping is it placed in ICD-11?

Select one:

- Factitious disorders
- Impulse control disorders
- Factors influencing health status or contact with health services
- Personality disorders
- Disruptive behaviour and dissocial disorders

Your answer is incorrect.

Malingering is not categorised as a mental health condition under ICD-11, although is mentioned under factors influencing health status or contact with health status so it can be accurately recorded when it is part of an individual's presentation.

The correct answer is: Factors influencing health status or contact with health services

Question **10**

Not answered

Marked out of 1.00

Flag question

The core concept of the paraphilic disorders in ICD-11 is that they involve sexual arousal that focuses upon non-consenting others. In which of the following scenarios would solitary acts or those involving consenting others meet the criteria for diagnosis?

Select one:

- If the individual takes on the role of another person as part of the behaviour
- If the individual is very concerned about their partner rejecting them due to their behaviour
- If the individual performs the acts outside of a private space
- If the behaviour is potentially dangerous
- If the individual would be embarrassed if their behaviour was discovered

Your answer is incorrect.

There is a category under paraphilic disorders which can be used if an individual's solitary acts or those with consenting others are associated with thoughts, fantasies, urges or behaviours which are very distressing (but not as a consequence of rejection or feared rejection of the arousal pattern by others) or confer direct risk of injury or death.

The correct answer is: If the behaviour is potentially dangerous

Question **11**

Not answered

Marked out of 1.00

Flag question

Which of the following is a change in the diagnostic criteria for anorexia nervosa in ICD-11?

Select one:

- Endocrine disturbance not necessary for diagnosis ✓
- Cross-listed with obsessive-compulsive and related disorders ✗
- Threshold BMI for diagnosis is now 18.5kg/m2 ✗ 18.
- Fat phobia necessary for diagnosis ✗
- Specific behaviours for weight loss not mentioned mentioned

Your answer is incorrect.

Endocrine disturbance is no longer necessary for a diagnosis of anorexia nervosa in ICD-11. A significant proportion of diagnoses of atypical anorexia nervosa were patients who met the other criteria for the full disorder but did not yet have endocrine disturbance; this change is intended to eliminate this issue. The threshold BMI for anorexia nervosa has been raised from 17.5kg/m2 but to 18kg/m2 rather than 18.5kg/m2. Fat phobia is no longer necessary for

the diagnosis. Qualifiers are used in ICD-11 to describe the behaviours associated with the diagnosis, such as restriction, and bingeing and purging.

The correct answer is: Endocrine disturbance not necessary for diagnosis

Question **12**

Not answered

Marked out of 1.00

Flag question

Which of the following is a new disorder in ICD-11, found under the obsessive-compulsive and related disorders diagnostic grouping?

Select one:

- Gaming disorder
- Compulsive sexual behaviour disorder
- Complex PTSD
- Excoriation disorder
- Hypochondriasis

Your answer is incorrect.

Excoriation disorder is a new diagnosis under the obsessive-compulsive and related disorders diagnostic grouping. It is characterised by repetitive picking at the skin, leading to lesions, and repeated but unsuccessful attempts to reduce the behaviour. Complex PTSD is a new disorder under the disorders specifically related to stress grouping. Compulsive sexual behaviour disorder is a new disorder under disorders of impulse control. Hypochondriasis was already an established diagnosis in ICD-10. Gaming disorder is a new disorder under disorders due to substance use and addictive behaviours.

The correct answer is: Excoriation disorder

Question **13**

Not answered

Marked out of 1.00

Flag question

How many groupings of diagnoses are there in the new ICD-11 chapter on mental, behavioural and neurodevelopmental disorders?

Select one:

- 11
- 21
- 10
- 26
- 18

Your answer is incorrect.

There are 21 groupings in the ICD-11 chapter focusing on mental health disorders, which uses an alphanumeric ordering system. They are:
• Neurodevelopmental disorders
• Schizophrenia and other primary psychotic disorders
• Catatonia
• Mood disorders
• Anxiety and fear-related disorders

• Obsessive-compulsive and related disorders
• Disorders specifically associated with stress
• Dissociative disorders
• Feeding and eating disorders
• Elimination disorders
• Disorders of bodily distress and bodily experience
• Disorders due to substance use and addictive behaviours
• Impulse control disorders
• Disruptive behaviour and dissocial disorders
• Personality disorders
• Paraphilic disorders
• Factitious disorders
• Neurocognitive disorders
• Mental and behavioural disorders associated with pregnancy, childbirth and the puerperium
• Psychological and behavioural factors affecting disorders or diseases classified elsewhere
• Secondary mental or behavioural syndromes associated with disorders or diseases classified elsewhereThere are 21 groupings in the ICD-11 chapter focusing on mental health disorders, which uses an alphanumeric ordering system. They are:
• Neurodevelopmental disorders
• Schizophrenia and other primary psychotic disorders
• Catatonia
• Mood disorders
• Anxiety and fear-related disorders
• Obsessive-compulsive and related disorders
• Disorders specifically associated with stress
• Dissociative disorders
• Feeding and eating disorders
• Elimination disorders
• Disorders of bodily distress and bodily experience
• Disorders due to substance use and addictive behaviours
• Impulse control disorders
• Disruptive behaviour and dissocial disorders
• Personality disorders
• Paraphilic disorders
• Factitious disorders
• Neurocognitive disorders
• Mental and behavioural disorders associated with pregnancy, childbirth and the puerperium
• Psychological and behavioural factors affecting disorders or diseases classified elsewhere
• Secondary mental or behavioural syndromes associated with disorders or diseases classified elsewhere

The correct answer is: 21

Question **14**

Not answered

Marked out of 1.00

Flag question

Which of the following concepts is reflective of a change in ICD-11 (from ICD-10)?

Select one:

- Removal of minimum symptom count for all disorders
- Introduction of gender incongruence as mental health disorder
- All disorders can be diagnosed at any age
- Cultural considerations have been removed
- Use of dimensional approach for all disorders

Your answer is incorrect.

ICD-11 takes a lifespan approach to diagnosis, permitting diagnoses which were usually associated with childhood or adolescence, such as attachment or conduct disorder, to be diagnosed at any age. It specifically considers the cultural background of a patient in diagnosis. A dimensional approach has been taken in personality and primary psychotic disorders, as well as for depressive episodes in mood disorders. Minimum symptom counts have been removed from diagnostic criteria for most disorders, but not major depression. Gender incongruence, replacing gender identity disorder, has been moved to the sexual health chapter of ICD-11 as it is not a mental health disorder

The correct answer is: All disorders can be diagnosed at any age

Question **15**

Not answered

Marked out of 1.00

Flag question

Which of the following is one of the maladaptive traits that can be used to further describe a diagnosis of personality disorder in ICD-11?

Select one:

- Mood variability
- Anxiousness
- Dissociality
- Affective lability
- Impulsivity

Your answer is incorrect.

The maladaptive traits that can be used to describe a diagnosis of personality disorder in ICD-11 are: negative affectivity, detachment, dissociality, disinhibition and anankastia. A further qualifier of 'borderline pattern' can also be used.

The correct answer is: Dissociality

Question **16**

Not answered

Marked out of 1.00

Flag question

Which of the following is a new mental health disorder under ICD-11?

Select one:

- Acute stress reaction
- Hyperkinetic disorder
- Prolonged grief disorder
- Trichotillomania
- Sadomasochistic disorder

Your answer is incorrect.

Prolonged grief disorder is a new mental health disorder under ICD-11 categorised under disorders specifically associated with stress. It describes a persistent and disabling grief reaction, which lasts longer than 6 months and is out of keeping with what is expected for the cultural context. 'Trichotillomania' was in ICD-10, as was hyperkinetic disorder, which has now been re-categorised as ADHD. Sadomasochism and other consensual sexual practices

are no longer generally considered to be pathological. 'Acute stress reaction' is no longer seen as a mental health condition, but as a normal reaction to extreme stress. It is categorised under factors influencing health status or contact with health services.

The correct answer is: Prolonged grief disorder

Question **17**

Not answered

Marked out of 1.00

Flag question

In ICD-11, which of the following groupings includes a dimensional element for the diagnoses within it?

Select one:

- Catatonia
- Disorders due to substance use and addictive behaviours
- Schizophrenia and other primary psychotic disorders
- Dissociative disorders
- Anxiety and fear-related disorders

Your answer is incorrect.

'Personality disorders', 'schizophrenia and other primary psychotic disorders', and depressive episodes in 'mood disorders' each include significant dimensional elements to enable clearer descriptions of the symptomatology experienced by patients. In schizophrenia and other primary psychotic disorders, this includes: positive symptoms; negative symptoms; depressive mood symptoms; manic mood symptoms; psychomotor symptoms; and cognitive symptoms. Neurodevelopmental disorders also include qualifiers for a diagnosis of autism spectrum disorder to enable a description of the extent of impairment of intellectual functioning and language abilities.

The correct answer is: Schizophrenia and other primary psychotic disorders

Finish review

PREVIOUS ACTIVITY
NEW 105Qs Fresh Practice Questions (2020-2021) Set 6

NEXT ACTIVITY
Basic Psychology

Jump to...

Drop here!

© 2022 SPMM Course Limited, UK.

Refund Policy | Terms & Conditions