Started on	Sunday, 24 July 2022, 3:40 PM
State	Finished
Completed on	Sunday, 24 July 2022, 3:40 PM
Time taken	8 secs
Marks	0.00/43.00
Grade	**0.00** out of 100.00

Question **1**

Not answered

Marked out of 1.00

Flag question

Which of the following is a gold standard test for intelligence?

Select one:

- Auditory Verbal Learning Test
- National Adult Reading Test
- Quick Test
- Wechsler Adult Intelligence Scale
- Wisconsin Card Sorting Test

Explanation:

The Wechsler Adult Intelligence Scales have represented the traditional gold standard in intellectual assessment for many years, based on carefully developed normative standards.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 52.

The correct answer is: Wechsler Adult Intelligence Scale

Question **2**

Not answered

Marked out of 1.00

Flag question

Which of the following is not a type of validity?

Select one:

- Face validity
- Construct validity
- Ecological validity
- Interrater validity
- Content validity

Explanation:

Inter-rater reliability (not validity) is a measure of agreement between two or more observers evaluating the same subjects using the same information.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 30.

The correct answer is: Interrater validity

Question **3**

Not answered

Marked out of 1.00

Flag question

Which of the following rating scales for depression used in adults is well validated in children aged 14 and older?

Select one:

- Clifton Assessment Procedure
- Beck Depression Inventory
- Cornell Depression Scale
- Wechsler Assessment Scale
- Edinburgh Depression Scale

Explanation:

The Beck Depression Inventory (BDI) was developed in the early 1960s to rate depression severity, with a focus on behavioural and cognitive dimensions of depression. It is a self-rated questionnaire containing 21 statements, with four possible responses for each. The current version has added more coverage of somatic symptoms and covers the most recent 2 weeks. Because it is a self-report measure, it can be used to screen for major depression (in individuals aged 13 and older). A score of >17 indicates moderate and >30 indicates severe depression.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 39.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 97.

The correct answer is: Beck Depression Inventory

Question **4**

Not answered

Marked out of 1.00

Flag question

Which of the following is not an alcohol assessment tool?

Select one:

- MAST
- AUDIT
- CAGE
- SCOFF
- CIWA

Explanation:

The SCOFF questions are useful as a screening tool for eating disorders in primary care.

SCOFF questions are:

- Do you make yourself **Sick** because you feel uncomfortably full?
- Do you worry you have lost **Control** over how much you eat?
- Have you recently lost more than **One** stone in a 3-month period?
- Do you believe yourself to be **Fat** when others say you are too thin?

- Would you say that **Food** dominates your life?

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 418.

The correct answer is: SCOFF

Question **5**

Not answered

Marked out of 1.00

Flag question

The SF-36 scale measures which of the following?

Select one:

- Occupational functioning
- Stress level at work
- Developmental psychopathology
- Health-related quality of life
- Social readjustment

Explanation:

The 36-Item Short Form Health Survey questionnaire (SF-36) is a very popular instrument for evaluating Health-Related Quality of Life. The SF-36 measures eight scales: physical functioning, role physical, bodily pain, general health, vitality, social functioning, role emotional, and mental health.

Ref: Lins L, Carvalho FM. SF-36 total score as a single measure of health-related quality of life: Scoping review. SAGE open medicine. 2016 Sep 30;4:2050312116671725.

The correct answer is: Health-related quality of life

Question **6**

Not answered

Marked out of 1.00

Flag question

The maximum possible score from General Health Questionnaire (28 items version) is

Select one:

- 84
- 28
- 12
- 42
- 72

Explanation:

The General Health Questionnaire (GHQ) is a self-rated questionnaire used as a screening instrument for the presence of psychiatric illness. The patient is asked to report the presence of a list of symptoms in the preceding weeks. Four versions are available, using 12, 28 (max score = 84), 30, and 60 items.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 96.

The correct answer is: 84

Question **7**

Not answered

Marked out of 1.00

Flag question

Choose the correct statement about Brief Psychiatric Rating Scale?

Select one:

- It is an assessment of global psychiatric symptomatology
- Each item is rated on a 6-point scale
- It focuses on negative symptoms
- It is a 30-item scale
- It is self administered

Explanation:

The Brief Psychiatric Rating Scale (BPRS) was developed in the late 1960s as a short clinician-rated scale (based on observation) for measuring the severity of psychiatric symptomatology. It consists of 18 or 24 items, each rated on a 7-8 point scale. It covers a broad range of areas, including thought disturbance, emotional withdrawal and retardation, anxiety and depression, and hostility and suspiciousness. It has been used extensively for decades as an outcome measure in treatment studies of schizophrenia. Reliability of the BPRS is good to excellent when raters are experienced, but this is difficult to achieve without substantial training.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. pp. 34-38.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 97.

The correct answer is: It is an assessment of global psychiatric symptomatology

Question **8**

Not answered

Marked out of 1.00

Flag question

Which of the following is the most important aspect of construct validity for a test of cognitive impairment in dementia?

Select one:

- The scores depend on educational attainment of the subjects
- It predicts the likelihood of progression to end stage dementia
- It includes most of the components of cognition that are commonly affected by dementia
- Scores are not affected by taking medications
- Experts agree with the choice of sub-items for the global scale

Explanation:

In psychometrics, construct validity is the extent to which a test measures a specified construct or hypothetical construct, determined by interpreting the psychological meaning of the test scores and testing implications of this interpretation. If our scale includes most of the components of cognition that are commonly affected by dementia, we can be assured of a good degree of construct validity.

Ref: Colman AM. Oxford Dictionary of Psychology, 4 th Edition. 2015.

The correct answer is: It includes most of the components of cognition that are commonly affected by dementia

Question **9**

Not answered

Marked out of 1.00

Flag question

Which of the following is a useful screening tool for early detection of an eating disorder?

Select one:

- Prochaska questionnaire
- AUDIT
- CAMCOG questionnaire
- SCOFF questionnaire
- CAGE questionnaire

Explanation:

The SCOFF questions are useful as a screening tool for eating disorders in primary care.

SCOFF questions are:

- Do you make yourself **Sick** because you feel uncomfortably full?
- Do you worry you have lost **Control** over how much you eat?
- Have you recently lost more than **One** stone in a 3-month period?
- Do you believe yourself to be **Fat** when others say you are too thin?
- Would you say that **Food** dominates your life?

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 418.

The correct answer is: SCOFF questionnaire

Question **10**

Not answered

Marked out of 1.00

Flag question

Which of the following is a self-rated questionnaire used as a screening instrument for the presence of psychiatric illness?

Select one:

- Global Assessment of Functioning Scale
- Quality of Life Interview
- Structured Clinical Interview
- General Health Questionnaire
- Diagnostic Interview Schedule

Explanation:

The General Health Questionnaire (GHQ) is a self-rated questionnaire used as a screening instrument for the presence of psychiatric illness. The patient is asked to report the presence of a list of symptoms in the preceding weeks. Four versions are available, using 12, 28 (max score = 84), 30, and 60 items.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 96.

The correct answer is: General Health Questionnaire

Question **11**

You want to screen for depression in adolescents of age between 14 and 18 years. Which is the best

Not answered

Marked out of 1.00

Flag question

screening tool?

Select one:

- Hamilton depression rating scale
- Child behavioural checklist
- Child and adolescent functional assessment scale
- Beck depression inventory
- SCID (Structured clinical interview for DSM-5)

Explanation:

The Beck Depression Inventory (BDI) was developed in the early 1960s to rate depression severity, with a focus on behavioural and cognitive dimensions of depression. It is a self-rated questionnaire containing 21 statements, with four possible responses for each. The current version has added more coverage of somatic symptoms and covers the most recent 2 weeks. Because it is a self-report measure, it can be used to screen for major depression (in individuals aged 13 and older). A score of >17 indicates moderate and >30 indicates severe depression.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 39.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 97.

The correct answer is: Beck depression inventory

Question **12**

Not answered

Marked out of 1.00

Flag question

Mr X is a 56-year-old gentleman assessed in the outpatient clinic. He is presenting with features of alcohol withdrawal. The patient is asked to complete the (SAWS) Short Alcohol Withdrawal Scale. The nurse wants to know the score above which he might require pharmacotherapy for withdrawal. Choose the correct answer from the following options.

Select one:

- 8
- 10
- 15
- 12
- 6

Explanation:

The Short Alcohol Withdrawal Scale is a tool to assess the severity of alcohol withdrawal. Patients indicate how they have felt in the previous 24 hours. A cut-off of 12 points is used where mild withdrawal is indicated by <12 points, and moderate to severe withdrawal (requiring intervention) is indicated by 12 points or greater.

Ref: Muncie Jr HL et al. Outpatient management of alcohol withdrawal syndrome. American family physician. 2013 Nov 1;88(9):589-95.

The correct answer is: 12

Question **13**

Not answered

Marked out of 1.00

Flag question

A tool designed to test cognitive functions has subtests that measure cognitive abilities well. This test can be said to have good

Select one:

- Concurrent validity
- Construct validity
- Predictive validity
- Face validity
- Incremental validity

Explanation:

In psychometrics, construct validity is the extent to which a test measures a specified construct or hypothetical construct, determined by interpreting the psychological meaning of the test scores and testing implications of this interpretation.

Ref: Colman AM. Oxford Dictionary of Psychology, 4 th Edition. 2015.

The correct answer is: Construct validity

Question **14**

Not answered

Marked out of 1.00

Flag question

The Clifton Assessment Procedure is used in which of the following situations?

Select one:

- Residential care for the elderly
- Schizophrenia rehabilitation units
- Children in foster care
- Care homes for the learning disabled
- Prisons with high mental health burden

CAPE (Pattie and Gilleard, 1979) is intended to assess level of disability and estimate need for care. It consists of a short cognitive scale and a behavioural rating scale. The latter has four sub-scales: physical disability, apathy, communication difficulties and social disturbance. It is quick and easy to administer, and is widely used by professional staff and care workers especially at elderly residential care settings

The correct answer is: Residential care for the elderly

Question **15**

Not answered

Marked out of 1.00

Flag question

Which of the following scales is used for measuring drug-induced movement abnormalities in general and tardive dyskinesia in particular?

Select one:

- AIMS scale
- Barnes Akathisia scale
- Simpson Angus scale

- Lane scale
- LUNSERS scale

Explanation:

The Abnormal Involuntary Movement Scale (AIMS) is a 12-item clinician-administered scale for assessing the severity of antipsychotic side effects, including tardive dyskinesia.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 97.

Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 597.

The correct answer is: AIMS scale

Question **16**

Not answered

Marked out of 1.00

Flag question

Which of the following technique is best utilised minimally if possible in those with high suggestibility?

Select one:

- Open questions
- Structured questionnaire
- Validation
- Summarising techniques
- Facilitating comments

Structured interviews have demonstrated a high degree of reliability, validity, and legal defensibility. But in highly suggestible patients, rigid interview questions are bound to induce responses that are in agreement with the tone of the question, leading to inaccurate answers.

The correct answer is: Structured questionnaire

Question **17**

Not answered

Marked out of 1.00

Flag question

The term test-retest reliability refers to

Select one:

- The correlation between the 1st and 2nd test of a number of subjects
- The correlation between the 1st and 2nd test in the same subject
- The correlation between the 1st and 2nd test carried out by different raters
- The agreement between 2 raters
- The correlation between ratings before and after treatment

Explanation:

Test-retest reliability is a measure of the ability of a psychological testing instrument to yield the same result for a single subject at 2 different test periods. Test-retest evaluations measure reliability only to the extent that the same subject's true condition remains stable in the time interval.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 30.

The correct answer is: The correlation between the 1st and 2nd test in the same subject

Question **18**

Not answered

Marked out of 1.00

Flag question

Which of the following rating scale for depression has more cognitive items than the others?

Select one:

- Young rating scale
- Hamilton rating Scale
- Beck depression inventory
- General Health Questionnaire
- Montgomery Asberg scale

Explanation:

The Beck Depression Inventory (BDI) was developed in the early 1960s to rate depression severity, with a focus on behavioural and cognitive dimensions of depression. It is a self-rated questionnaire containing 21 statements, with four possible responses for each. The current version has added more coverage of somatic symptoms and covers the most recent 2 weeks. Because it is a self-report measure, it can be used to screen for major depression (in individuals aged 13 and older). A score of >17 indicates moderate and >30 indicates severe depression.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 39.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 97.

The correct answer is: Beck depression inventory

Question **19**

Not answered

Marked out of 1.00

Flag question

What is the most reliable instrument to make a clinical diagnosis according to DSM criteria?

Select one:

- AIMS
- SCID
- SCAN
- HAMD
- MMPI

Explanation:

The Structured Clinical Interview for the DSM (SCID) is one of the most used diagnostic instruments in clinical research worldwide. The latest version of the instrument is called ‘SCID-5’. SCID is a clinician-administered semi-structured interview for use in patients in whom a psychiatric diagnosis is suspected. It is primarily used in research with trained interviewers.

Ref: Osório FL et al. Clinical validity and intrarater and test–retest reliability of the Structured Clinical Interview for DSM-5–Clinician Version (SCID-5-CV). Psychiatry and clinical neurosciences. 2019 Dec;73(12):754-60.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 96.

The correct answer is: SCID

Question **20**

Not answered

Marked out of 1.00

Flag question

With respect to Young Mania Rating Scale, which of the following is least likely to be correct?

Select one:

- It is sensitive to change
- It does not have items on suicidality
- Psychotic symptoms are not included in the scale
- It has 11 items
- It is self-rated

Explanation:

The Young Mania Rating Scale (YMRS) has 11 items and is based on the patient's subjective report of their condition over the previous 48 hours. Additional information is based upon clinical observations made during the course of the clinical interview. There are four items that are graded on a scale from 0 to 8 (irritability, speech, thought content, and disruptive/aggressive behaviour), while the remaining seven items are graded on a scale of 0 to 4. Mania is indicated by a score of 12, but many clinical studies require YMRS > 20 to determine a manic state.

Ref: Di Giacomo E et al. Unblending borderline personality and bipolar disorders. Journal of psychiatric research. 2017 Aug 1;91:90-7.

The correct answer is: It is self-rated

Question **21**

Not answered

Marked out of 1.00

Flag question

Which is the most commonly used screening tool to diagnose dementia in older people?

Select one:

- CAMDEX
- Addenbrooke cognitive examination
- Abbreviated mental test score
- Mini-mental state examination
- Mini-COG

MMSE: Cut off 24/30, takes 5-10 minutes to complete and is a standard baseline test. It is the most widely used cognitive test in old age psychiatry and has been validated in a variety of populations. It is good for screening global cognitive dysfunctions as opposed to focal cognitive dysfunctions. It covers the following areas of cognition such as orientation, memory, concentration, language, praxis and Gnosis

The correct answer is: Mini-mental state examination

Question **22**

Not answered

Marked out of

Which of the following rating scales retain the aspects of a clinical examination during consultation?

Select one:

1.00

Flag question

- HoNOS
- CATEGO
- SANS
- PSE
- BPRS

PSE retains the aspects of a mental state examination

The correct answer is: PSE

Question **23**

Not answered

Marked out of 1.00

Flag question

Which of the following scales of depression is most sensitive to change?

Select one:

- Visual analogue scale
- Beck depression inventory
- Zung self-rating scale
- Montgomery-Asberg depression rating scale
- Hospital anxiety and depression scale

Montgomery-Asberg depression rating scale is most sensitive to change. It has a ten items version, which requires a clinical interview like Hamilton depression rating scale.

The correct answer is: Montgomery-Asberg depression rating scale

Question **24**

Not answered

Marked out of 1.00

Flag question

All of the following are projective tests except

Select one:

- Rorschach test
- Sentence Completion Test
- HAM-D
- Draw A Person Test
- Thematic Apperception Test

Explanation:

Projective tests of personality include the Rorschach test, Thematic Apperception Test (TAT), Sentence Completion Test, Holtzman Inkblot Technique (TIT), Figure drawing (typically human forms but can involve houses or other forms) and Make-a-Picture Story (MAPS). HAM-D is not a projective test.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 59.

The correct answer is: HAM-D

Question **25**

Not answered

Marked out of 1.00

Flag question

Which of the following refers to the extent to which performance on a test fits with theoretical schema about the test attributes?

Select one:

- Criterion validity
- Convergent validity
- Construct validity
- Face validity
- Concurrent validity

Explanation:

In psychometrics, construct validity is the extent to which a test measures a specified construct or hypothetical construct, determined by interpreting the psychological meaning of the test scores and testing implications of this interpretation.

Ref: Colman AM. Oxford Dictionary of Psychology, 4 th Edition. 2015.

The correct answer is: Construct validity

Question **26**

Not answered

Marked out of 1.00

Flag question

Considering the Beck Depression Inventory, which of the following is true?

Select one:

- Refers to symptoms over the last five days
- It has 21 items
- Maximum score is 42
- Has no item on suicidal ideas
- Strictly a clinician-rated scale

Explanation:

The Beck Depression Inventory (BDI) was developed in the early 1960s to rate depression severity, with a focus on behavioural and cognitive dimensions of depression. It is a self-rated questionnaire containing 21 statements, with four possible responses for each. The current version has added more coverage of somatic symptoms and covers the most recent 2 weeks. Because it is a self-report measure, it can be used to screen for major depression (in individuals aged 13 and older). A score of >17 indicates moderate and >30 indicates severe depression.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 39.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 97.

The correct answer is: It has 21 items

Question **27**

Not answered

A 22-year-old man is admitted with troublesome voices discussing his sexuality. He also believes they influence his actions and control his thoughts. Which of the following instruments can be used to

Marked out of 1.00

Flag question

monitor his symptoms?

Select one:

- Schedule for Assessment of Negative Symptoms
- Brief Psychiatric Rating Scale
- Bristol Assessment Scale
- Thought Language Index
- General Health Questionnaire

Explanation:

The Brief Psychiatric Rating Scale (BPRS) was developed in the late 1960s as a short clinician-rated scale (based on observation) for measuring the severity of psychiatric symptomatology. It consists of 18 or 24 items, each rated on a 7-8 point scale. It covers a broad range of areas, including thought disturbance, emotional withdrawal and retardation, anxiety and depression, and hostility and suspiciousness. It has been used extensively for decades as an outcome measure in treatment studies of schizophrenia. Reliability of the BPRS is good to excellent when raters are experienced, but this is difficult to achieve without substantial training.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. pp. 34-38.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 97.

The correct answer is: Brief Psychiatric Rating Scale

Question **28**

Not answered

Marked out of 1.00

Flag question

Which of the following interview schedules have replaced the Present Status Examination in contemporary epidemiological studies?

Select one:

- SCAN
- PANSS
- SSPI
- CASH
- SANS

Explanation:

The Schedules for Clinical Assessment in Neuropsychiatry (SCAN) has evolved from the older Present State Examination (PSE). SCAN is a set of instruments aimed at assessing and classifying psychopathology in adults. Instruments include the PSE-10 (the tenth edition of the Present State Examination), the SCAN glossary, the Item Group Checklist, and the Clinical History Schedule. The interview is semi-structured, the aim being to encapsulate the clinical interview while minimising its vagaries. Where there is doubt, the interviewer can proceed to a freestyle interview to clarify further. It is designed for use by psychiatrists or clinical psychologists, thereby utilising clinical interviewing skills in evaluating each symptom. The use of mental health professionals in interviewing with SCAN makes this an expensive method but has the advantage of approximating the ‘gold standard’ diagnosis achieved by clinical interview.

Ref: Casey P, Kelly B. Fish’s clinical pathology: Signs and symptoms in psychiatry, 4th Edition.

2019.p. 9.

The correct answer is: SCAN

Question **29**

Not answered

Marked out of 1.00

Flag question

Which one of the following questionnaires can be used to identify psychiatric 'caseness' in the general population?

Select one:

- BDI - Beck depression inventory
- HAM-D - Hamilton Depression Scale
- Structured clinical interview DSM-5
- GHQ - General Health Questionnaire
- SCAN - Schedules for Clinical Assessment in Neuropsychiatry

Explanation:

The General Health Questionnaire (GHQ) is a self-rated questionnaire used as a screening instrument for the presence of psychiatric illness. The patient is asked to report the presence of a list of symptoms in the preceding weeks. Four versions are available, using 12, 28 (max score = 84), 30, and 60 items.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 96.

The correct answer is: GHQ - General Health Questionnaire

Question **30**

Not answered

Marked out of 1.00

Flag question

Which of the following is not a rating scale for mood disorders?

Select one:

- Beck inventory
- Hamilton Rating scale
- Montgomery-Asberg rating scale
- Simpson-Angus scale
- Young rating scale

Explanation:

The Simpson-Angus Scale is a clinical tool commonly used to detect and monitor extrapyramidal symptoms. It rates 10 items from 0 to 4 points, yielding a total body score divided by 10, and considered normal up to 0.3. It has been criticized for its bias in the rigidity domain, accounting for 6 of 10 items.

Ref: Blanchet PJ, Rompré PH. Clinimetric evaluation of the Simpson-Angus Scale in older adults with schizophrenia. Journal of clinical psychopharmacology. 2014 Feb 1;34(1):36-9.

The correct answer is: Simpson-Angus scale

Question **31**

You have designed a study questionnaire and would like to measure the construct validity. Which of

Not answered

Marked out of 1.00

Flag question

the following will you be testing?

Select one:

- The questionnaire predicts what will happen in the future
- Constituent parts of the questionnaire are suitable and related to the phenomenon being tested
- The questionnaire covers most symptoms related to the phenomenon being measured
- The questionnaire gives same results when tested on the same patient twice by a single rater
- The questionnaire gives same results when used by two different raters on the same patient twice

Explanation:

In psychometrics, construct validity is the extent to which a test measures a specified construct or hypothetical construct, determined by interpreting the psychological meaning of the test scores and testing implications of this interpretation.

Ref: Colman AM. Oxford Dictionary of Psychology, 4 th Edition. 2015.

The correct answer is: Constituent parts of the questionnaire are suitable and related to the phenomenon being tested

Question **32**

Not answered

Marked out of 1.00

Flag question

Which one among the following rating scales is a self-report measure?

Select one:

- Abbreviated Mental Test
- Glasgow Coma Scale
- Hamilton Depression Scale
- Beck Depression Inventory
- Clinical Global Improvement scale

Explanation:

The Beck Depression Inventory (BDI) was developed in the early 1960s to rate depression severity, with a focus on behavioural and cognitive dimensions of depression. It is a self-rated questionnaire containing 21 statements, with four possible responses for each. The current version has added more coverage of somatic symptoms and covers the most recent 2 weeks. Because it is a self-report measure, it can be used to screen for major depression (in individuals aged 13 and older). A score of >17 indicates moderate and >30 indicates severe depression.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 39.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 97.

The correct answer is: Beck Depression Inventory

Question **33**

Not answered

Marked out of 1.00

Which of the following is correct concerning the SCID?

Select one:

- Based on a dimensional diagnostic scheme

Flag question

- Used in conjunction with ICD-10 classification
- Provides the Axis V diagnosis in DSM-IV
- Useful for untrained lay interviewers
- Primarily used in research

Structured Clinical Interview for DSM (SCID) is a clinician-administered semi-structured clinical interview for use with patients in whom a psychiatric diagnosis is suspected. It is primarily used in research with trained interviewers to inform the operationalized diagnosis psychiatric disorders.

The correct answer is: Primarily used in research

Question **34**

Not answered

Marked out of 1.00

Flag question

Mr. Y is a 30-year-old man who presented to the A and E after an overdose. He believes that there are cameras spying on his movements. He has stopped going out and no longer socialises with friends. He has been hearing voices discussing his sexuality in an unpleasant way. Which of the following tools would be helpful to assess and monitor his psychopathology?

Select one:

- MADRS
- PANSS
- MMSE
- HAS
- GHQ

Explanation:

The Positive and Negative Symptom Scale (PANSS) is a clinician-administered rating scale for the assessment of severity and monitoring of change of symptoms in patients with a diagnosis of schizophrenia. It contains items that cover positive symptoms, negative symptoms, and general psychopathology.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 97.

The correct answer is: PANSS

Question **35**

Not answered

Marked out of 1.00

Flag question

Which one of the following scales is used for monitoring clinical recovery?

Select one:

- SCAN
- CAGE
- MMPI
- AUDIT
- HoNOS

Explanation:

The Health of the Nation Outcomes Scales (HoNOS) is a 12-scale clinician-rated measure developed by the Royal College of Psychiatrists to guide everyday clinical practice and measure health and social care outcomes in secondary mental health service for adults age 18-65 years. It was designed to be sensitive to improvement, deterioration, or lack of change over time.

Ref: James M et al. A review and update of the Health of the Nation Outcome Scales (HoNOS). BJPsych bulletin. 2018 Apr;42(2):63-8.

The correct answer is: HoNOS

Question **36**

Not answered

Marked out of 1.00

Flag question

Which of the following scales can be used by primary care physicians, general medical professionals and also in community studies?

Select one:

- Signs and Symptoms in Psychotic Illness
- Scale of Assessment of Positive Symptoms
- General Health Questionnaire
- Simpson-Angus Scale
- Brief Psychiatric Rating Scale

Explanation:

The General Health Questionnaire (GHQ) is a self-rated questionnaire used as a screening instrument for the presence of psychiatric illness. The patient is asked to report the presence of a list of symptoms in the preceding weeks. Four versions are available, using 12, 28 (max score = 84), 30, and 60 items.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 96.

The correct answer is: General Health Questionnaire

Question **37**

Not answered

Marked out of 1.00

Flag question

Which of the following is true concerning Brief Psychiatric Rating Scale (BPRS)?

Select one:

- Useful for global psychiatric symptom ratings
- Cannot be used to detect treatment response
- It requires no clinical training to administer
- It has 20 items
- It has 8 items

Explanation:

The Brief Psychiatric Rating Scale (BPRS) was developed in the late 1960s as a short clinician-rated scale (based on observation) for measuring the severity of psychiatric symptomatology. It consists of 18 or 24 items, each rated on a 7-8 point scale. It covers a broad range of areas, including thought disturbance, emotional withdrawal and retardation, anxiety and depression, and hostility and suspiciousness. It has been used extensively for decades as an outcome measure in treatment studies of

schizophrenia. Reliability of the BPRS is good to excellent when raters are experienced, but this is difficult to achieve without substantial training.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. pp. 34-38.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 97.

The correct answer is: Useful for global psychiatric symptom ratings

Question **38**

Not answered

Marked out of 1.00

Flag question

Concurrent validity is also known as

Select one:

- Face validity
- Construct validity
- Incremental validity
- Criterion validity
- Predictive validity

In criterion-related validity, we examine whether the operationalised criteria used in a scale is aligned to the theory behind a construct that is being tested.

The correct answer is: Criterion validity

Question **39**

Not answered

Marked out of 1.00

Flag question

Brief Psychiatric Rating Scale is a clinician-rated scale used to measure the severity of clinical symptoms based on observation. It is most commonly used in patients with a diagnosis of

Select one:

- Alcohol use disorder
- Personality disorder
- Schizophrenia
- Bipolar disorder
- Somatisation disorder

Explanation:

The Brief Psychiatric Rating Scale (BPRS) was developed in the late 1960s as a short clinician-rated scale (based on observation) for measuring the severity of psychiatric symptomatology. It consists of 18 or 24 items, each rated on a 7-8 point scale. It covers a broad range of areas, including thought disturbance, emotional withdrawal and retardation, anxiety and depression, and hostility and suspiciousness. It has been used extensively for decades as an outcome measure in treatment studies of schizophrenia. Reliability of the BPRS is good to excellent when raters are experienced, but this is difficult to achieve without substantial training.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. pp. 34-38.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 97.

The correct answer is: Schizophrenia

Question **40**

Not answered

Marked out of 1.00

Flag question

You have designed a study questionnaire and would like to measure the inter-rater reliability. Which of the following will you be testing?

Select one:

- The questionnaire is valid when it gives same results if assessed on two different occasions by same rater
- The scores given by one rater are well concealed and cannot be copied by another rater
- The questionnaire gives same results when used by two different raters on the same patient
- Self-rating and clinician rating gives the same score from the questionnaire
- The questionnaire is valid when it gives same results if assessed by two different assessors on two different patients

Explanation:

Inter-rater reliability is a measure of the consistency of ratings, equal to the correlation between ratings given by different raters of the same people or stimuli.

Ref: Colman AM. Oxford Dictionary of Psychology, 4 th Edition. 2015.

The correct answer is: The questionnaire gives same results when used by two different raters on the same patient

Question **41**

Not answered

Marked out of 1.00

Flag question

In psychometry, internal consistency refers to

Select one:

- Degree of correlation between one test item with other items
- Degree of agreement between two raters
- Degree of agreement between results obtained in two different occasions
- Degree of correlation between two different versions of the scale
- Degree of agreement between two similar scales

Explanation:

In psychometrics, internal consistency is an aspect of reliability associated with the degree to which the items of a test measure the same construct or attribute. It is indexed by Cronbach's alpha, the Kuder-Richardson coefficient, or the split-half reliability.

Ref: Colman AM. Oxford Dictionary of Psychology, 4 th Edition. 2015.

The correct answer is: Degree of correlation between one test item with other items

Question **42**

Not answered

The assessment tool useful in measuring the severity and change in psychotic symptoms is

Marked out of 1.00

Flag question

Select one:

- Neuropsychiatric inventory
- CATEGO
- Young mania rating scale
- Brief psychiatric rating scale
- SCAN

Explanation:

The Brief Psychiatric Rating Scale (BPRS) was developed in the late 1960s as a short clinician-rated scale (based on observation) for measuring the severity of psychiatric symptomatology. It consists of 18 or 24 items, each rated on a 7-8 point scale. It covers a broad range of areas, including thought disturbance, emotional withdrawal and retardation, anxiety and depression, and hostility and suspiciousness. It has been used extensively for decades as an outcome measure in treatment studies of schizophrenia. Reliability of the BPRS is good to excellent when raters are experienced, but this is difficult to achieve without substantial training.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. pp. 34-38.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 97.

The correct answer is: Brief psychiatric rating scale

Question **43**

Not answered

Marked out of 1.00

Flag question

Which of the following scales is NOT used for measuring drug-induced movement abnormalities?

Select one:

- LUNSERS scale
- Lane scale
- Barnes Akathisia scale
- Simpson Angus scale
- AIMS scale

Lane scale is used for minor physical anomalies.

The correct answer is: Lane scale

Finish review

PREVIOUS ACTIVITY
Dynamic Psychopathology

NEXT ACTIVITY
EMIs Assessment Part 1

Jump to...

Drop here!

© 2022 SPMM Course Limited, UK.