Mock Paper A(1)

Started on	Monday, 28 May 2018, 12:17 AM
State	Finished
Completed on	Thursday, 7 June 2018, 1:03 PM
Time taken	10 days 12 hours
Overdue	10 days 11 hours
Grade	**0.00** out of 100.00

Question 1

Not answered

Marked out of 1.00

Flag question

MoK Classification 001
How many episodes of mania and/or depression should be seen in 12 months time to diagnose rapid cycling bipolar disorder?

Select one:

- 3
- 4
- 2
- 6
- 8

Rapid cycling bipolar disorder is defined in the DSM-5 as a type of bipolar illness in which the patient experiences four or more episodes of mania and/or major depression per year (Rapid cycling bipolar disorder, http://www.ncbi.nlm.nih.gov/pubmed/11825328).

The correct answer is: 4

Question 2

Not answered

Marked out of 1.00

Flag question

MoK Classification 002
Axis V of DSM-IV recorded Global Assessment of Functioning. What is the maximum score in the GAF scale?

Select one:

- 7
- 100
- 10
- 60
- 80

The GAF is a scale used by clinicians to describe a patient's level of functioning in three areas: social (i.e., marriage, friends), occupational, and psychological. The GAF scale ranges from 100 (highest functioning) to 0. The GAF is recorded on Axis V.

The correct answer is: 100

Question 3

MoK Classification 003

Not answered

Marked out of 1.00

Flag question

The term simple schizophrenia was added as to the categories of existing schizophrenias by

Select one:

- Bleuler
- Hubbard
- Hecker
- Angst
- Liddle

The diagnostic category called simple schizophrenia has a long history; it was one of the traditional schizophrenic subtypes identified by Bleuler and was later accepted by Kraepelin. The term has been removed from the official American nosology but has been retained in ICD system.

The correct answer is: Bleuler

Question 4

Not answered

Marked out of 1.00

Flag question

MoK Classification 004
Which of the following is correct with regard to a diagnosis of post schizophrenic depression?

Select one:

- The patient must not have a previous history of depression
- The patient must not be on any antipsychotic medication
- Some positive symptoms must be present along with depression
- It can be diagnosed only 12 months after the last psychotic episode
- It is classified along with psychotic disorders

Post-schizophrenic depression forms F20.4 - it is classified as a type of schizophrenia along with other psychotic disorders in ICD10. While the general criteria for schizophrenia must have been met within the past twelve months, they must not be met in full at present. But one of either positive/negative symptoms/formal thought disorder/catatonia symptoms must be present. The depressive symptoms must be sufficiently prolonged, severe and extensive to meet criteria for at least a mild depressive episode (F32.0). The patient may or may not be on treatment with antipsychotics or antidepressants. Previous history of depression does not preclude a diagnosis

The correct answer is: It is classified along with psychotic disorders

Question 5

Not answered

Marked out of 1.00

Flag question

MoK Classification 006
Which of the following features seen in a person suffering from grief reaction 1 year after the loss suggests depression?

Select one:

- Mummification
- Guilt regarding omission of care
- Physical health problems
- Preoccupation with the relative
- Anniversary reaction

Signs of difficulty in assimilating the loss vary from the slight (e.g., expecting the person to return home from work and setting two places for dinner) to the extreme of mummification (e.g., retaining a child's room exactly as it was on the day of death for years). These signs may be harbingers of depression if they persist

for more than one year. Mummification was first described by Gorer in 1965.

The correct answer is: Mummification

Question 6

Not answered

Marked out of 1.00

Flag question

MoK Classification 008

In order to make a ICD-10 diagnosis of delusional disorder, the delusions must be present for at least

Select one:

- 3 days
- 1 month
- 6 months
- 2 weeks
- 3 months

Persistent Delusional Disorder includes a variety of disorders in which long-standing (more than three months) delusions constitute the only, or the most conspicuous, clinical characteristic. Delusional disorders that have lasted for less than three months should be classified, at least temporarily, under schizophreniform illnesses.

The correct answer is: 3 months

Question 7

Not answered

Marked out of 1.00

Flag question

MoK Classification 010

The most common co-morbid psychiatric condition associated with panic disorder is

Select one:

- Depression
- Schizophrenia
- Agoraphobia
- Borderline personality disorder
- Social phobia

Agoraphobia is the most frequent comorbid diagnosis, with 30 - 60% of panic disorder patients satisfying the diagnostic criteria for agoraphobia.

The correct answer is: Agoraphobia

Question 8

Not answered

Marked out of 1.00

Flag question

MoK Classification 011

A person drinks alcohol as soon as he wakes up in the morning. During working hours, he again drinks in the lunch time and after getting home he needs to have at least 6 units of alcohol. His behaviour is best described as

Select one:

- Compulsive behaviour
- Alcohol dependence
- Rumination
- Impulsivity
- Obsessions

Compulsive behaviours seen in addictions are associated with 'pleasure seeking' i.e. seeking the drug to obtain 'chemical' pleasure. This is typical of drug dependence states.

The correct answer is: Alcohol dependence

Question 9

Not answered

Marked out of 1.00

Flag question

MoK Classification 013
What is least likely with regard to night terrors?

Select one:

- Associated with organised automatic behaviours
- Amnesia of the episode
- Occurs in deep slow wave sleep
- Often seen during early nocturnal sleep
- Increased autonomic arousal is noted during the episode

It is rarely associated with complex motor activities.

The correct answer is: Associated with organised automatic behaviours

Question 10

Not answered

Marked out of 1.00

Flag question

MoK Classification 014
Which of the following clinical features suggests a diagnosis of social phobia rather than agoraphobia?

Select one:

- Being house bound
- Having a fear of getting into situations of no escape
- Having a fear of being under continuous scrutiny from others
- Avoiding the use of public transport
- Avoiding crowded places

Both social phobia and agoraphobia can present with house-bound behaviour and avoiding public places. But in patients with social phobia the underlying fear is that of embarrassing oneself by 'doing something stupid' while others are watching them.

The correct answer is: Having a fear of being under continuous scrutiny from others

Question 11

Not answered

Marked out of 1.00

Flag question

MoK Classification 015
Marie presents with a first episode of depression. She is started on an SSRI. Seven days later, she is found to be overactive, easily distractible, irritable and sleepless, which lasts for about 5 days. The symptoms resolve when the SSRI was stopped. Most appropriate diagnosis among the following list is

Select one:

- Unipolar mania
- Bipolar type 2 disorder
- Mixed affective disorder
- Bipolar type 1 disorder
- Substance induced mood disorder

SSRI-induced mood switch or destabilisation is not a recognised classification in ICD or DSM. But as this lady has had a period of hypomania and past history of depression, she can be diagnosed with bipolar disorder type 2.

The correct answer is: Bipolar type 2 disorder

Question 12

Not answered

Marked out of 1.00

Flag question

MoK Classification 016
A 23-year-old woman is referred by her GP. She states that her eyes are set wide apart and people call her 'fish-face'. She spends hours in front of the mirror examining her eyes. She has seen surgeons twice to get plastic surgery done to her eye folds. However there is no obvious deformity. The most appropriate diagnosis is

Select one:

- Body dysmorphic disorder
- Social phobia
- Panic disorder
- Panic disorder
- Generalized anxiety disorder

In body dysmorphic disorder, there is an excessive concern (overvalued idea) about trivial or non-existent physical abnormalities, which are perceived to be deformities

The correct answer is: Body dysmorphic disorder

Question 13

Not answered

Marked out of 1.00

Flag question

MoK Classification 017
A 36-year-old woman complains of severe tiredness not relieved by rest for over 6 months and is unable to work. There is no identifiable organic pathology and she denies any pain in her body. The most appropriate diagnosis is

Select one:

- Fibromyalgia
- Generalized anxiety disorder
- Chronic Fatigue Syndrome
- Social phobia
- Depression

Chronic fatigue syndrome is characterized by persistent or relapsing unexplained chronic fatigue for at least 6 months, is of new or definite onset, is not the result of an organic disease or of continuing exertion and is not alleviated by rest. Fatigue results in a substantial reduction in previous occupational, educational, social, and personal activities. Four or more of the following symptoms, concurrently present for more than 6 months: impaired memory or concentration, sore throat, tender cervical or axillary lymph nodes, muscle pain, pain in several joints, new headaches, unrefreshing sleep, or malaise after exertion

The correct answer is: Chronic Fatigue Syndrome

Question 14

Not answered

Marked out of 1.00

Flag question

MoK Classification 018
Unstable affect and impulsive behaviour is characterised by which of the following personality disorders, according to DSM-5 Criteria?

Select one:

- Cluster E

- Cluster D
- Cluster C
- Cluster A
- Cluster B

There are four Cluster B personality disorders: antisocial, borderline, narcissistic, and histrionic. The DSM views these as a subset of personality disorders that are characterized by dramatic, emotional or erratic behavior.

The correct answer is: Cluster B

Question 15

Not answered

Marked out of 1.00

Flag question

MoK Classification 019
A 23-year-old lady has a history of olfactory hallucinations along with flushing. The most likely diagnosis is

Select one:

- Stuporous excitement
- Delirium
- Frontal partial seizures
- Complex partial seizures
- Occipital seizures

Wilder Penfield reported that fewer than 1% of patients with partial-onset seizures experienced olfactory symptoms. The majority of patients with epilepsy and olfactory auras will have clear complex partial or secondary generalized seizures in addition to their sensory symptoms although these episodes may occur at separate times.

The correct answer is: Complex partial seizures

Question 16

Not answered

Marked out of 1.00

Flag question

MoK Clinical Examination 001
Which of the following enzymes mediate alcohol related brain injury?

Select one:

- Alcohol dehydrogenase
- Lactate dehydrogenase
- Transketolase
- Aldehyde dehydrogenase
- Alpha ketoglutarate

Thiamine (vitamin B1) in its active form, thiamin pyrophosphate (TPP), is a co-enzyme for several enzymes, including transketolase. Transketolase is an important enzyme in the pentose phosphate pathway (PPP), essential for energy transduction and for generating ribose for nucleic acid synthesis. Transketolase also links the PPP to glycolysis, allowing a cell to adapt to a variety of energy needs, depending on its environment. Abnormal transketolase expression and/or activity have been implicated in a number of diseases where thiamin availability is low, including Wernicke-Korsakoff's Syndrome and other forms of alcohol-related brain injuries. Alexander-Kaufman, K. and Harper, C.(2009): The International Journal of Biochemistry and Cell Biology 41, 717-720.

The correct answer is: Transketolase

Question 17

Not answered

Marked out of 1.00

Flag question

MoK Clinical Examination 002
Length of time cannabis can be detected in the urine of a person who has used it only once is

Select one:

- 3 months
- 24 hours
- 1 month
- 7 days
- 3 days

With regular use, it may be present for weeks.

The correct answer is: 3 days

Question 18

Not answered

Marked out of 1.00

Flag question

MoK Clinical Examination 003
Which of the following reflexes may NOT be exaggerated in upper motor neuron lesions?

Select one:

- Abdominal reflex
- Ankle jerk reflex
- Biceps reflex
- Knee jerk reflex
- Jaw jerk reflex

Deep tendon reflexes are exaggerated. Abdominal reflex is not a deep tendon reflex; it is a superficial reflex.

The correct answer is: Abdominal reflex

Question 19

Not answered

Marked out of 1.00

Flag question

MoK Clinical Examination 004
Which of the following is the most important discriminating feature between mild cognitive impairment and dementia?

Select one:

- Educational attainment
- Duration of symptoms of memory disturbances
- Family history of Alzheimer's
- Activities of daily living
- MMSE score

Mild Cognitive Impairment (MCI) is a borderline state between age-associated cognitive decline and mild dementia. MCI is separated from mild dementia by an absence of global intellectual deterioration and the preservation of activities of daily living (ADL).

The correct answer is: Activities of daily living

Question 20

MoK Clinical Examination 005

Not answered

Marked out of 1.00

Flag question

Which one of the following is best suited for an assessment of global cognitive functioning?

Select one:

- Mini-COG
- Clock drawing test
- MMSE
- Montreal cognitive assessment guide
- Addenbrookes cognitive examination

Addenbrookes cognitive examination is a cognitive screening instrument used in the assessment of global cognitive functioning. It tests various cognitive domains such as Orientation, Attention, Memory, Verbal fluency, and Language and Visuospatial ability. It incorporates the questions on the MMSE and expands on the domains of memory, language and visuospatial concepts and adds tests of verbal fluency.

The correct answer is: Addenbrookes cognitive examination

Question 21

Not answered

Marked out of 1.00

Flag question

MoK Clinical Examination 006
A woman presents to A&E with the following features: weight loss, fatigue, hyperkalaemia and hyponatraemia. What is the most likely diagnosis?

Select one:

- Addison's disease
- Hypothyroidism
- Cushing's disease
- Diabetes insipidus
- Diabetes mellitus

Hyperkalaemia and hyponatraemia are seen in people with Addison's disease. The most common symptoms are fatigue, muscle weakness, fever, weight loss, darkening (hyperpigmentation) of the skin, including areas not exposed to the sun and orthostatic hypotension.

The correct answer is: Addison's disease

Question 22

Not answered

Marked out of 1.00

Flag question

MoK Clinical Examination 007
Mr. Y has a history of alcohol dependence. He is admitted to a medical ward as he developed memory difficulties following Wernicke's encephalopathy. The bedside test most likely to be impaired is

Select one:

- Digit span
- Address test
- Months of the year backwards
- Spelling world backwards
- Serial sevens

Korsakoff syndrome is a preventable memory disorder that usually emerges (although not always) in the aftermath of an episode of Wernicke's encephalopathy. Episodic memory is severely affected in the Korsakoff syndrome, and the learning of new semantic memories is variably affected. 'Implicit' aspects of memory are preserved.

The correct answer is: Address test

Question 23

Not answered

Marked out of 1.00

Flag question

MoK Clinical Examination 008
On ocular examination of a 74-year-old patient, constriction of pupils is noted on looking at near objects but no reaction to light. This is called

Select one:

- Senile pupil
- Adie's pupil
- Anisocoria
- Argyll Robertson Pupil
- Marcus Gunn pupil

Argyll Robertson Pupil (ARP): The pupils are bilaterally small and reduce in size when the patient focuses on a near object (i.e. Accommodation Reflexes Present - ARP again!) but do not constrict when exposed to bright light (i.e. pupillary reflexes absent). They are seen in patients with neurosyphilis (hence the name "Prostitute's Pupil") and diabetic neuropathy.

The correct answer is: Argyll Robertson Pupil

Question 24

Not answered

Marked out of 1.00

Flag question

MoK Descriptive Psychopathology 001
Which of the following syndromes is named after the person who initially described them?

Select one:

- Othello syndrome
- Fregoli syndrome
- Munchaussen syndrome
- Rabbit syndrome
- Cotard's syndrome

Cotard syndrome is named after Jules Cotard (1840-1889), a French neurologist who first described the condition, which he called le delire de negation.

The correct answer is: Cotard's syndrome

Question 25

Not answered

Marked out of 1.00

Flag question

MoK Descriptive Psychopathology 002
You are interviewing a newly admitted patient in your ward. When asked what he had for breakfast, he says 'cereals'. Then when you asked 'what is your name?' he says 'cereals'. When you asked 'what is his date of birth?' he continues to say 'cereals'. The most likely symptom exhibited here is

Select one:

- Mannerism
- Automatism
- Perseveration
- Stuttering
- Stereotypy

Perseveration is the uncontrolled repetition or continuation of a response (e.g., behavior, word, thought, activity, strategy, or emotion) in the absence of an ongoing occasion or rationale for that behavior or emotion (e.g., the topic or task requirements have changed).

The correct answer is: Perseveration

Question 26

Not answered

Marked out of 1.00

Flag question

MoK Descriptive Psychopathology 003
Which of the following is not a feature of Kubler-Ross model of grief?

Select one:

- Agitation
- Bargaining
- Denial
- Depression
- Acceptance

The Kubler-Ross model, commonly referred to as the "five stages of grief", is an explanatory model for grief reaction introduced by Elisabeth Kubler-Ross. When a person is faced with the reality of impending death or other extreme, awful fate, he or she will experience a series of emotional stages. This sequence can be remembered using the acronym DABDA, which include: denial, anger, bargaining, depression, and acceptance.

The correct answer is: Agitation

Question 27

Not answered

Marked out of 1.00

Flag question

MoK Descriptive Psychopathology 004
Inability to verbalize one's strong emotions is termed as

Select one:

- Anhedonia
- Avolition
- Apathy
- Ambivalence
- Alexithymia

Alexithymia refers to difficulties in emotionalizing, fantasizing, verbalizing, identifying, and analyzing emotions.

The correct answer is: Alexithymia

Question 28

Not answered

Marked out of 1.00

Flag question

MoK Descriptive Psychopathology 005
Which of the following may indicate poor response to psychotherapeutic interventions?

Select one:

- alexithymia
- insomnia
- apraxia
- anhedonia
- amnesia

Alexithymia is associated with poor outcome in both traditional psychodynamic psychotherapy and supportive therapy. This negative effect is found in individual and group psychotherapies. In the context of group therapy, higher levels of alexithymic features elicit negative reactions from one's therapist, which partially contribute to the poor outcome experienced by such patients. This negative reaction of therapists towards alexithymic patients appears to be in response to the lack of positive emotional expression. Ref: http://www.ncbi.nlm.nih.gov/pubmed/20471096

The correct answer is: alexithymia

Question 29

Not answered

Marked out of 1.00

Flag question

MoK Descriptive Psychopathology 007
A patient with schizophrenia protrudes his tongue repeatedly in and out when asked to do it once. He is exhibiting which of the following symptoms?

Select one:

- Chorea
- Stereotypy
- Ambivalence
- Hemiballism
- Ambitendence

Ambitendency can be regarded as a mild variety of negativism. The patient appears to be in conflict regarding moving their body. The presence of opposing tendencies to action is described as 'jack in the box' feature of catatonia.

The correct answer is: Ambitendence

Question 30

Not answered

Marked out of 1.00

Flag question

MoK Descriptive Psychopathology 009
Ben is worried that he has to shout 'Willy Wonka' whenever he sees an attractive woman, despite his best efforts to keep calm. This is best termed as

Select one:

- A first rank symptom of schizophrenia
- A mental compulsion
- An obsessional impulse
- Somatic passivity
- A made act

Obsessions can present as thoughts, images, impulses or 'sticky words'. In the above scenario, obsessions are presenting as impulses i.e. the rush to do something.

The correct answer is: An obsessional impulse

Question 31

Not answered

Marked out of 1.00

Flag question

MoK Descriptive Psychopathology 010
Which one of the following terms describes fantasy thinking?

Select one:

- Autistic thinking
- Conceptualism
- Rationalism
- Concrete thinking

- Magical thinking

Normal thinking is of three functional types: a) Fantasy/dereistic thinking or autistic thinking: There is no goal direction, unrealistic like daydreaming. Predominant in cluster A personality, dissociation and pseudologia fantastica. b). Imaginative thinking: Again fantasy elements but admixed with memory, involving abstract concepts but goal-directed and does not cross boundaries of possibility and realism. Determining tendency of thoughts is preserved e.g. creative lateral thinking. c). Rational or conceptual thinking: based on material reality and uses logic.

The correct answer is: Autistic thinking

Question 32

Not answered

Marked out of 1.00

Flag question

MoK Descriptive Psychopathology 011
Mr Y has a cardiac arrest. He is in the resuscitation room and experiences seeing himself outside of his body. This experience is called

Select one:

- Somatic hallucination
- Lilliputian hallucination
- Reflex hallucination
- Autoscopic hallucination
- Extracampine hallucination

Autoscopic hallucination is the visual experience of seeing oneself projected external to the body space. Males predominate in a 2:1 ratio; impaired consciousness is a common accompaniment and depression is the commonest psychiatric cause. They are also called phantom mirror images and may take the form of pseudohallucinations. Schizophrenia, TLE, parietal lesions are also implicated. In negative autoscopy, one looks into a mirror and sees no image at all.

The correct answer is: Autoscopic hallucination

Question 33

Not answered

Marked out of 1.00

Flag question

MoK Descriptive Psychopathology 012
You note tonic spasm in a woman who cannot resist closing her right eye while talking to you. She is able to carry out most of her daily activities including watching television normally. This is best described as

Select one:

- Tonic seizures
- Blepharospasm
- Normal phenomenon
- Catatonic posturing
- Multiple sclerosi

Blepharospasm is often idiopathic and benign.

The correct answer is: Blepharospasm

Question 34

Not answered

Marked out of 1.00

Flag question

MoK Descriptive Psychopathology 013
On listening to a joke shared by a friend, a woman loses her muscle tone and collapses suddenly. This is described as

Select one:

- Drop attacks
- Cataplexy
- Syncopy
- Catatonia
- Catalepsy

In a state of cataplexy, an intense emotional state triggers objective transient muscle weakness (often with areflexia). The clinical manifestations are varied, ranging from involuntary eye closure and neck weakness to a subtle buckling of the knees to generalized muscle weakness that causes the patient to collapse. Consciousness and awareness of the environment are preserved throughout the episode.

The correct answer is: Cataplexy

Question 35

Not answered

Marked out of 1.00

Flag question

MoK Descriptive Psychopathology 014
A patient with schizophrenia is observed in the waiting room with various semi-purposive movements, with an inability to change the movements at will or make them goal-directed. This is best described as

Select one:

- Mannerism
- Agitation
- Restlenessness
- Catatonic impulsiveness
- Akathisia

Catatonia often presents with immobility, stupor, posturing, rigidity, staring, grimacing, and withdrawal, The alternative presentation of catatonia (described here) is an excited state (impulsivity and associated agitation), along with combativeness, and autonomic instability.

The correct answer is: Catatonic impulsiveness

Question 36

Not answered

Marked out of 1.00

Flag question

MoK Descriptive Psychopathology 015
The following is the most common obsession in OCD:

Select one:

- Rumination
- Fear of contamination
- Symmetry
- Checking
- Magical thinking

The most common obsession is fear of contamination, followed by pathological doubt, a need for symmetry, and aggressive obsessions. The most common compulsion is checking, which is followed by washing, symmetry, the need to ask or confess, and counting. Swedo SE, et al. Obsessive-Compulsive Disorder in children and adolescents: Clinical phenomenology of 70 consecutive cases. Arch Gen Psychiatry 1989;46:335-41

The correct answer is: Fear of contamination

Question 37

Not answered

Marked out of 1.00

Flag question

MoK Descriptive Psychopathology 016
When asked what he did for a living, a patient stated that he worked at a shop selling clothes. He went on to say "I don't like the messages on t-shirts. I wish I was a sailor". He is exhibiting

Select one:

- Circumstantial speech
- Tangential speech
- Stilted speech
- Pressured speech
- Clang associations

The presence of excessive unwanted details suggests circumstantial speech, though the patient responds to the question appropriately.

The correct answer is: Circumstantial speech

Question 38

Not answered

Marked out of 1.00

Flag question

MoK Basic Psychology 001
With regard to the psychology of attention, all of the following are correct except

Select one:

- Focused attention is modality specific
- Sustained attention refers to vigilance
- Divided attention is the ability to attend to multiple task demands
- Alternating attention is the highest form of attention
- Shadowing is a test for selective attention

Sohlberg and Mateer proposed a hierarchic model based on the recovering of attention processes of brain-damaged patients when coming out of a coma. 1. Focused attention: The ability to respond discretely to specific visual, auditory or tactile stimuli. This is the most primitive form of attention 2. Sustained attention (vigilance): The ability to maintain a consistent behavioral response during continuous and repetitive activity. 3. Selective attention: The ability to maintain a behavioral or cognitive set in the face of distracting or competing stimuli. Shadowing is a test for selective attention in which simultaneously two different inputs are provided, one to each ear, and the subject is required to 'shadow' or follow one of the inputs. 4. Alternating attention: The ability of mental flexibility that allows individuals to shift their focus of attention and move between tasks having different cognitive requirements. 5. Divided attention: This is the highest level of attention, and it refers to the ability to respond simultaneously to multiple tasks or multiple task demands. In dual-task technique which tests divided attention, the subject is asked to attend to both inputs in an attempt to divide attention. Thus the ability to attend to multiple task demands is tested.

The correct answer is: Alternating attention is the highest form of attention

Question 39

Not answered

Marked out of 1.00

Flag question

MoK Basic Psychology 002
Cognitive theory suggests the following as the underlying factor for anxiety disorders:

Select one:

- Somatisation
- Catastrophisation
- Nihilism
- Depersonalisation
- Personalisation

The term catastrophizing was used by Albert Ellis, the founder of rational-emotional therapy, almost four decades ago. Catastrophizing is related to anxiety (e.g., anxiety is associated with the tendency to overemphasize the probability of a catastrophic outcome and the possible consequences of such an outcome). During the past decade, cognitive models of panic disorder have emphasized the role of catastrophizing in panic attacks (Retrieved from http://journals.lww.com/clinicalpain/Fulltext/2001/03000)

The correct answer is: Catastrophisation

Question 40

Not answered

Marked out of 1.00

Flag question

MoK Basic Psychology 003
Visual or iconic stimuli in immediate memory lasts for

Select one:

- 20-30 minutes
- 0.5 seconds
- 2-3 seconds
- 20-30 seconds
- 0.05 seconds

Visual or iconic- 0.5 seconds; Auditory or echoic -2-3 seconds.

The correct answer is: 0.5 seconds

Question 41

Not answered

Marked out of 1.00

Flag question

MoK Basic Psychology 006
Which one of the following is an example of non homeostatic motives?

Select one:

- All of the above
- Temperature control
- Achievement
- Hunger
- Thirst

There are two types of motivating forces; homeostatic and non homeostatic. Homeostatic motives arise from physiological needs such as thirst, hunger and temperature control and are concerned with basic survival. Non homeostatic motives refer to psychological drives such as curiosity, adventure, achievement, etc.

The correct answer is: Achievement

Question 42

Not answered

Marked out of 1.00

Flag question

MoK Basic Psychology 007
Perceptual constancy can be demonstrated for

Select one:

- All of the above
- Size
- Brightness constancies
- Colour
- Location

Perceptual constancy: It refers to the ability to perceive the appearance of objects as constant, even though, their retinal images keep changing. It includes several components: size, shape, colour, location and brightness constancies.

The correct answer is: All of the above

Question 43

Not answered

Marked out of 1.00

Flag question

MoK Basic Psychology 012

One of the local newspaper reports that drug addicts carry out muggings. An 83-year-old lady was mugged and the public concluded the attacker was a drug addict. Select the cognitive heuristic in this case.

Select one:

- Framing
- Availability
- Simulation
- Representation
- Anchoring

Availability- a heuristic for judging frequency and probability. Higgins and King (1981) described this phenomenon as a readiness with which a particular schema is used in information processing.

The correct answer is: Availability

Question 44

Not answered

Marked out of 1.00

Flag question

MoK Basic Psychology 013

In a program aimed at cessation of smoking, a 45-year-old woman is required to smoke many cigarettes in a small booth and over a short period of time. What is the process that is employed in this approach?

Select one:

- Aversive conditioning
- Stimulus control
- Systematic desensitization
- Variable ratio reinforcement
- Habituation

The smoker is being exposed to an aversive stimulus. The physiologic and psychological trauma is so severe and so aversive that the patient may choose to avoid smoking in the future.

The correct answer is: Aversive conditioning

Question 45

Not answered

Marked out of 1.00

Flag question

MoK Basic Psychology 014

A rat in the cage initially got electric shocks whenever goes to a particular area of the cage. Now it stays in electric shock free area. What is the learning process involved?

Select one:

- Avoidance conditioning
- Classical conditioning
- Reciprocal inhibition
- Shaping
- Higher order conditioning

Avoidance learning is a type of learning where an organism learns to avoid certain responses or situations. In avoidance conditioning, the response or act prevents the aversive stimulus from occurring. For example, people with posttraumatic stress after a road traffic accident may take a different road to work that avoids cues associated with the accident and, therefore, the anxiety that may be associated with the exposure. Avoidance is a powerful reinforcer that produces a strong form of learning that is difficult to extinguish.

The correct answer is: Avoidance conditioning

Question 46

Not answered

Marked out of 1.00

Flag question

MoK Basic Psychology 019
A forensic investigator tries to reconstruct the crime scene and encourages visits to scene of crime by witnesses. Which recall phenomenon is replicated in this study?

Select one:

- Blocking
- Failure of prospective memory
- Encoding failure
- State dependent memory
- Context dependent memory

Context-dependent memory refers to the situation or context in which you find yourself is a determinant of the ability to retrieve memories (Godden and Baddeley 1975). It suggests that contextual and environmental cues (aspects of the environment) that are effective in activating recall of information. This phenomenon has been useful particularly in crime detection. Reconstructing the crime scene and visits to scenes of crime has been found to be effective in triggering recall of details of the crime. Similarly, returning to a childhood home village/ town or old school/college brings long forgotten memories flooding back. It is sometimes not even necessary to visit a place to evoke memories but simply imagining the place or situation may be sufficient to retrieve a flood of memories.

The correct answer is: Context dependent memory

Question 47

Not answered

Marked out of 1.00

Flag question

MoK Basic Psychology 020
Which of the following is a test that primarily measures sustained attention?

Select one:

- Tower of London test
- Wisconsin card sorting test
- Rey verbal learning test
- Continuous performance test
- Trail making test B

A Continuous Performance Task/Test, or CPT, is a psychological test which measures a person's sustained and selective attention and impulsivity. Sustained attention is the ability to maintain a consistent focus on some continuous activity or stimuli and is associated with impulsivity. Selective attention is the ability to focus on relevant stimuli and ignore competing stimuli. This skill is associated with distractibility.

The correct answer is: Continuous performance test

Question 48

Not answered

Marked out of 1.00

Flag question

MoK Basic Psychology 021
A person with a fear of heights takes to parachute jumping for a hobby. This is best described as

Select one:

- Counterphobic behaviour

- Reciprocal inhibition
- Desensitisation
- reaction formation
- Projection

Counterphobic behaviour is any behavior that seeks out experiences that are consciously or unconsciously feared in an attempt to master those fears. An example would be parachute jumping for a person with a fear of heights.

The correct answer is: Counterphobic behaviour

Question 49

Not answered

Marked out of 1.00

Flag question

MoK Social Psychology 001
Fundamental attribution error refers to

Select one:

- Attributing personal factors for one's own behaviours
- Attributing external factors for others behaviours
- Failure to make attributions for observed behaviours
- Attributing personal factors for others' behaviours
- Attributing external factors for one's own behaviours

Often when we try to understand and explain what happens in social settings, we tend to explain behaviour of others in terms of internal disposition, such as personality traits, abilities, motives, etc. as opposed to external situational factors. This bias is more pronounce for external behaviours of others rather than self.

The correct answer is: Attributing personal factors for others' behaviours

Question 50

Not answered

Marked out of 1.00

Flag question

MoK Social Psychology 002
According to Allport, the denial of opportunity to the minority group due to the majority group practicing their prejudice is termed as

Select one:

- Avoidance
- Discrimination
- Antilocution
- Physical attack
- Extermination

Allport identified the following processes operating during conflicts between the minority and the majority: 1. Antilocution 2. Avoidance 3. Discrimination: Minority group is discriminated against by denying them opportunity and equality. 4. Physical attack 5. Extermination.

The correct answer is: Discrimination

Question 51

Not answered

Marked out of 1.00

Flag question

MoK Sociocultural Psychiatry 001
Which of the following is NOT true with regard to research ethics?

Select one:

- Written approval must be obtained from ethics committees
- Advertisements can be used to recruit subjects
- Subjects can be recruited by contacting medical colleagues
- A subject is ethically bound to complete a study after giving written informed consent
- Financial incentives can be offered for subjects to participate in research

Subjects, both healthy controls and patients, are reimbursed for their participation in research. But signing a consent form does not mean that they cannot discontinue the study during its course.

The correct answer is: A subject is ethically bound to complete a study after giving written informed consent

Question 52

Not answered

Marked out of 1.00

Flag question

MoK Sociocultural Psychiatry 002
Which of the following is true with regard to aetiological theories of suicide?

Select one:

- Sensational media reports of suicide reduce acute risk of suicide
- Serotonin receptor binding is decreased in suicidal patients
- The stressors in the stress-diathesis model can include genetic factors, physical illness or childhood experiences
- Serotonin transporter binding is increased in suicidal patients
- Durkheim proposed social models of suicide

According to stress-diathesis model, a typical stressor includes the acute worsening of a psychiatric disorder or an acute psychosocial crisis. Pessimism and aggression/impulsivity are components of the diathesis for suicidal behaviour; sex, religion, familial/genetic factors, childhood experiences and various other factors influence this diathesis. In the prefrontal cortex of people who committed suicide, serotonin transporter binding was decreased but serotonin receptor binding was increased. Mann (2003). Nature Reviews Neuroscience 4, 819-828.

The correct answer is: Durkheim proposed social models of suicide

Question 53

Not answered

Marked out of 1.00

Flag question

MoK Sociocultural Psychiatry 004
A patient is willing to stop alcohol use and is accepting the need for a positive change, though she has not taken any steps yet. According to Prochaska and DiClemente she in the stage of

Select one:

- Precontemplation
- Ready for Action
- Preparation
- Maintenance
- Acting out

The Stages of Change model shows that, for most persons, a change in behaviour occurs gradually, with the patient moving from being uninterested, unaware or unwilling to make a change (precontemplation), to considering a change (contemplation), to deciding and preparing to make a change.During the preparation stage, patients prepare to make a specific change. They may experiment with small changes as their determination to change increases. Genuine, determined action is then taken and over time, attempts to maintain the new behaviour occur. Relapses are almost inevitable in the process of working towards a life-long change.

The correct answer is: Ready for Action

Question 54
Not answered
Marked out of 1.00
Flag question

MoK Sociocultural Psychiatry 005
What is the number of women that have been reported being amputated in the genital area?

Select one:
- 1 million
- 10 millions
- 100 millions
- 100, 000
- 10000

WHO has estimated that 100 to 140 million girls and women worldwide are currently living with the consequences of FGM.

The correct answer is: 100 millions

Question 55
Not answered
Marked out of 1.00
Flag question

MoK Sociocultural Psychiatry 007
Schizophrenic patients were more likely to relapse if discharged to families exhibiting 'high expressed' emotion for more than:

Select one:
- 55 hours per week
- 33 hours per week
- 15 hours per week
- 24 hours per week
- 45 hours per week

Brown et al in 1962 found that schizophrenic patients were more likely to relapse if discharged to families exhibiting high expressed emotion (hostility, critical comments and emotional over-involvement) especially if they were exposed to these families for more than 35 hours per week. Relapse was less likely in low expressed emotion families that expressed warmth and positive remarks. Leff et al in 1985 demonstrated the interaction between expressed emotion and antipsychotic medication in a schizophrenic relapse and in a controlled trial they also showed the effectiveness of family intervention therapy consisting of education about schizophrenia and its treatment, communication training and support for relatives, in alleviating the effects of high expressed emotion. Ref; Core Psychiatry; pg 259-293.

The correct answer is: 33 hours per week

Question 56
Not answered
Marked out of 1.00
Flag question

MoK Sociocultural Psychiatry 008
All of the following are potential boundary violations between a therapist and a patient except

Select one:
- A private therapist expecting payment for each session of therapy
- A therapist colludes with a patient against a third party
- A clinician accepts gifts from one patient but not others
- Influencing patient for political causes
- A clinician tries to be impressive often by disclosing personal information

If the therapist expects all patients to pay for each session, then he may not be violating any boundaries by asking a patient do so.

The correct answer is: A private therapist expecting payment for each session of therapy

Question 57

Not answered

Marked out of 1.00

Flag question

MoK Sociocultural Psychiatry 009
Which of the following conditions is NOT strongly associated with social disadvantage?

Select one:

- Depression
- Alzheimer's disease
- Schizophrenia
- Bipolar disorder
- Alcohol use disorders

Early life socioeconomic level is related to level of cognition in late life but not to rate of cognitive decline or risk of AD.Neuroepidemiology. 2005;25(1):8-14. Epub 2005 Apr 25.

The correct answer is: Alzheimer's disease

Question 58

Not answered

Marked out of 1.00

Flag question

MoK Sociocultural Psychiatry 011
A Malaysian man shows sudden excitement with marked violence after a period of withdrawal. He returns to premorbid state after a violent incident. This is called as

Select one:

- Latah
- Cathard
- Windigo
- Amok
- Piblokto

Amok is a culture-bound syndrome seen in Malaysia and South East Asia. It is characterised by initial sullen period, followed by an outburst of violent, sometimes homicidal behaviour and a return to premorbid state after the episode. Though mostly dissociative, rarely amok may occur during a brief psychotic episode or constitute the onset or an exacerbation of a chronic psychotic process.

The correct answer is: Amok

Question 59

Not answered

Marked out of 1.00

Flag question

MoK Sociocultural Psychiatry 016
Which of the following is most associated with an increased risk of developing schizophrenia?

Select one:

- Being frequently placed in a 'double-blind situation' as a child
- Being born in an urban area
- Maternal cannabis use during pregnancy
- Experiencing high expressed emotion in the family home
- Maternal alcohol use during pregnancy

There is robust and consistent evidence from epidemiological studies showing that urban birth is associated with an increased risk of developing schizophrenia. Evidence suggests that this exposure may be associated with a sizeable proportion of cases. (McGrath 2006)

The correct answer is: Being born in an urban area

Question 60

Not answered

Marked out of 1.00

Flag question

MoK HumanDevelopment 001
Who proposed theories on cognitive development?

Select one:

- Sullivan
- Ainsworth
- Munroe
- Piaget
- Fromm

Swiss cognitive psychologist Jean Piaget contributed immensely to our understanding of the development of cognitive faculties in children. His key concepts assimilation, accommodation, equilibration, and schemas are applicable to learning at any age.

The correct answer is: Piaget

Question 61

Not answered

Marked out of 1.00

Flag question

MoK HumanDevelopment 002
Which of the following is a normal stage of grief?

Select one:

- Mourning
- Protest
- Acting Out
- Ambivalence
- Bargaining

The classic work on stages of grief came from Erich Lindemann at Massachusetts General Hospital, who studied 101 bereaved people. He wrote an article, published in 1944, titled 'Symptomology and Management of Acute Grief.' In this article he described a set pattern: After an unexpected death, there is an initial shock that lasts 10-14 days. After the initial shock comes a period of intense sadness, and the grieving person may withdraw from social contact. Next comes anger, as the grieving person seems to 'protest' the unexpected death. Finally, within a year or so, the grief is resolved, and the person returns to normal.

The correct answer is: Protest

Question 62

Not answered

Marked out of 1.00

Flag question

MoK HumanDevelopment 003
Which period refers to the autistic phase of Mahler's stages of early development?

Select one:

- 6-36 months
- 0-2 months
- 2-6 Months
- 6-12 months

- 12-24 months

According to Mahler, the first few weeks of infancy is termed as the 'Normal Autistic Phase' (she later discarded this designation). A newborn infant is blissfully unaware of anything but its own needs, and the mother needs to be available to meet those needs in this stage.

The correct answer is: 0-2 months

Question 63

Not answered

Marked out of 1.00

Flag question

MoK HumanDevelopment 004
Average age of puberty in boys is

Select one:

- 11-12 years
- 10-11 years
- 8-9 years
- 9-10 years
- 14-15 years

In the UK the average age of onset of puberty in males is 11.2 years; for females it is 11 years. Menarche on average is at 12.5 years for females.

The correct answer is: 11-12 years

Question 64

Not answered

Marked out of 1.00

Flag question

MoK HumanDevelopment 005
The strange-situation study of attachment was carried out on

Select one:

- Children aged 6-9 years
- Infants between 1-6 months
- Newborn babies
- Toddlers
- Children aged 12-24 months

The strange-situation study of attachment was carried out on children between 12-24 Months. It is a procedure designed to test the quality of the attachment bond between mother and their infants. It has three phases with the infant's reactions being noted in each of the phases; 1. Reaction to a stranger in the presence of the mother 2. Reaction when left alone with a stranger 3. Reaction when re united with the mother

The correct answer is: Children aged 12-24 months

Question 65

Not answered

Marked out of 1.00

Flag question

MoK HumanDevelopment 007
At what age can a child build a tower of 6 cubes?

Select one:

- 2 years
- 6 -10 months
- 5 years

- 12-18 months
- 4 years

At age 2, a child can run and build a tower of 6 cubes. At age 3, a child can make a tower of 9-10 cubes.

The correct answer is: 2 years

Question 66

Not answered

Marked out of 1.00

Flag question

MoK HumanDevelopment 011
Mastering intimate relationships despite the demands of work is a successful developmental task for ages between

Select one:

- 60 to 80
- 40 to 60
- 20 to 40
- 10 to 15
- 80 to 90

The Intimacy vs. Isolation conflict is emphasized around the ages of 30. At the start of this stage, identity vs. role confusion is coming to an end, though it still lingers at the foundation of the stage (Erikson, 1950).

The correct answer is: 20 to 40

Question 67

Not answered

Marked out of 1.00

Flag question

MoK HumanDevelopment 014
Which of the following is correct with regard to day-care for children?

Select one:

- 4 months day care before age 1 for more than 20hours/week may be detrimental
- Day care reduces attachment behaviour
- Day care induces neurotic clinging
- Regular day care affects cognitive development
- Day care increases maternal bonding

A series of studies by Belsky and Rovine (1988) claim that insecure attachments were more likely to develop if the child had been receiving care of at least 20 hours per week for 4 months or more before its first birthday.

The correct answer is: 4 months day care before age 1 for more than 20hours/week may be detrimental

Question 68

Not answered

Marked out of 1.00

Flag question

MoK Sociocultural Psychiatry 017
When a psychiatric patient is unwell, family and friends report their actions to mental health professionals in an attempt to get them admitted to an institution. What was the term used by Goffman to describe this?

Select one:

- Institutional neurosis
- Role stripping
- Mortification
- Batch living

Betrayal funnel

Goffman depicted the sequence of events through which the understandings of a 'pre-patient' would be betrayed using the process of referral and admission. Goffman described this as a "betrayal funnel" wherein during each stage of the admission process the family and friends might try to give patients the impression that they would not lose any more of the rights that they enjoyed in the community. (Goffman 1961, p. 140).

The correct answer is: Betrayal funnel

Question 69

Not answered

Marked out of 3.00

Flag question

MoK Assessment EMI001

Features of defence mechanisms

For each of the characteristic features described below chose the most appropriate defence mechanism suitable from the above list

Ascribing unacceptable motivations to others — Choose...

Resentment by noncooperation — Choose...

Failure to accept or face conflicts — Choose...

Explanation: In **projection**, intolerable feelings, impulses or thoughts are falsely attributed to other people. **Passive aggression** can manifest itself as repeated postponements, rigidity in task performance, resentment, sullenness, or repeated failure to take up the responsibility to complete tasks. **Denial** is characterised by a total failure to accept a problem even when faced with the facts contrary to the negation.Passive aggression

The correct answer is: Ascribing unacceptable motivations to others – Projection, Resentment by noncooperation – Passive aggression, Failure to accept or face conflicts – Denial

Question 70

Not answered

Marked out of 3.00

Flag question

MoK Assessment EMI002

Supportive techniques

For each of the following examples, identify the type of supportive techniques aimed at eliciting information

Sometimes when people are upset and low in their mood, they think of hurting themselves. Has this been true for you? — Choose...

I think it is best for you to consider psychotherapy at this time. If I am you, I will give this a serious thought — Choose...

This depression may be very difficult for you. But I think it is very likely with the proper treatment you can get back to your job. — Choose...

Explanation:

Validation/normalization: It helps to decrease a patient's sense of embarrassment about a feeling or behaviour. It can be done by quoting how it is normal for people to have different emotions/ reactions/ behaviours, etc.

Advice: Many patients seek help directly; it is acceptable to provide advice but based on sound understanding of the context. Premature advice can be obstructive than facilitative.

Ref: Adapted from Kay J & Tasman A. Essentials of Psychiatry, 2nd edition, 2006. John Wiley & Sons, Ltd.

The correct answer is: Sometimes when people are upset and low in their mood, they think of hurting themselves. Has this been true for you? – Validation, I think it is best for you to consider psychotherapy at this time. If I am you, I will give this a serious thought – Advice, This depression may be very difficult for you. But I think it is very likely with the proper treatment you can get back to your job. – Reassurance

Question 71

Not answered

Marked out of 4.00

Flag question

MoK Assessment EMI004

Rating scales and their uses

For each description given below, select the relevant rating scale from the list above;

This scale is useful for rating severity in patients with a pre-existing diagnosis of OCD	Choose...
Commonly used screening tool used in primary care and general population studies	Choose...
This scale measures overall psychosocial functioning	Choose...
7-point scale useful to assess global observation of severity of psychiatric illness	Choose...

Explanation:

Yale-Brown obsessive-compulsive scale (YBOCS) is a clinician-administered semi-structured interview, allowing rating of severity in patients with a pre-existing diagnosis of OCD.

General Health Questionnaire (GHQ) is a self-rated screening instrument for the presence of psychiatric illness. It is a commonly used screening tool used in primary care and general population studies.

GAF of Global assessment of functioning scale is a 100-item, self-report rating scale, measuring overall psychosocial functioning.

CGI or Clinical global impression scale is a 7-point scale useful for clinicians to assess the global observed severity of psychiatric illness.

The correct answer is: This scale is useful for rating severity in patients with a pre-existing diagnosis of OCD – Yale-Brown obsessive compulsive scales (YBOCS), Commonly used screening tool used in primary care and general population studies – General health questionnaire (GHQ), This scale measures overall psychosocial functioning – Global assessment of functioning scale (GAF), 7-point scale useful to assess global observation of severity of psychiatric illness – Clinical Global impressions scale (CGI)

Question 72

Not answered

Marked out of 3.00

Flag question

MoK Assessment EMI005

Dangerous complications of eating disorders

Identify the metabolic abnormalities commonly seen in the following clinical situations

A 16-year-old girl with history of significant weight loss, body image distortions and morbid fear of fatness had shown some endocrine abnormalities in her blood	Choose...
A 23-year-old woman with history of bulimia nervosa and is determined to reduce weight by repetitive vomiting	Choose...
A 20-year-old woman uses regular laxatives for last two years in order to reduce weight. In addition to hypokalemia associated with acidosis, what other electrolyte abnormalities results from chronic laxative abuse?	Choose...

Explanation: The diagnosis is anorexia nervosa. The common endocrine changes in anorexia include low concentrations of luteinising hormone, follicle stimulating hormone and oestradiol, low T3 but T4 in low normal range, average concentrations of thyroid stimulating hormone (low T3 syndrome), mild increase in plasma cortisol and raised growth hormone level.

Electrolyte disturbances can be of variable nature and are present in those who frequently vomit or misuse large quantities of laxatives or diuretics. Vomiting results in metabolic alkalosis and hypokalemia. In repetitive vomiting, loss of hydrochloric acid from gastric juices lead to metabolic alkalosis (loss of acid - alkalosis). Laxative misuse results in metabolic acidosis, hyponatraemia, hypokalemia. During laxative induced diarrhoea, a considerable amount of bicarbonate may be lost in the stool. With healthy kidneys, the

lost bicarbonate is replaced adequately, and a serious base deficit does not develop. When there is poor renal blood flow due to hypovolaemia/starvation, base deficit and acidosis develop rapidly. Acidosis also results from excessive production of lactic acid when patients have severe diarrhoea.

The correct answer is: A 16-year-old girl with history of significant weight loss, body image distortions and morbid fear of fatness had shown some endocrine abnormalities in her blood – Low T3 and high growth hormone, A 23-year-old woman with history of bulimia nervosa and is determined to reduce weight by repetitive vomiting – Hypokalemia, A 20-year-old woman uses regular laxatives for last two years in order to reduce weight. In addition to hypokalemia associated with acidosis, what other electrolyte abnormalities results from chronic laxative abuse?
– Hyponatremia

Question 73

Not answered

Marked out of 4.00

Flag question

MoK Assessment EMI006
Clinically notable motor disorders
Choose one option for each of the following descriptions

Sustained abnormalities of posture occurring focally in writer's cramp or torticollis	Choose...
Brief jerky downward movements of the wrist, when the patient extend both arms with the wrists dorsiflexed, palms facing forward, and eyes closed	Choose...
Quasi-purposeful movements affecting multiple joints with a distal preponderance	Choose...
Quivering of the muscle seen under the skin in neuromuscular disorder	Choose...

Explanation:

Writer's cramp is a form of focal dystonia, which is an involuntary, sustained muscle contraction causing twisting movements and abnormal postures. Writer's cramp is the most typical dystonia occurring in the setting of repetitive movement disorders.

The term chorea means dance. These are quasi-purposeful movements affecting multiple joints with a distal preponderance.

Asterixis represents the failure to maintain a position actively and is "caused by the abnormal function of diencephalic motor centres that regulate the tone of the agonist and antagonist muscles involved in maintaining posture" (Asterixis - NEJM - The New England Journal of Medicine.., http://www.nejm.org/doi/full/10.1056/NEJMicm0911157 (accessed April 21, 2015).

Fasciculations may be seen under the skin as quivering of the muscle. Although fasciculations are typically benign (particularly when they occur in the calf), if widespread, they can be associated with neuromuscular disease, including amyotrophic lateral sclerosis (ALS).
Neurological History and Physical Examination, http://emedicine.medscape.com/article/1147993-overview (accessed March 31, 2015).

The correct answer is: Sustained abnormalities of posture occurring focally in writer's cramp or torticollis – Dystonia, Brief jerky downward movements of the wrist, when the patient extend both arms with the wrists dorsiflexed, palms facing forward, and eyes closed – Asterixis, Quasi-purposeful movements affecting multiple joints with a distal preponderance – Chorea, Quivering of the muscle seen under the skin in neuromuscular disorder – Fasciculations

Question 74

Not answered

Marked out of 3.00

Flag question

MoK Assessment EMI007
Medical disorders & psychiatry
For each description below choose ONE option from the above list

A 44-year-old homeless man is brought into A&E with agitation and features of psychosis and depression. An EEG reveals diffuse attenuation in alpha activity, and his gait is abnormal. MRI shows atrophy of the striatum.	Choose...

A 25-year-old man recently returned from Kansas. He presents with unprovoked aggression. Of late he experienced recurrent seizures followed by periods of impaired mentation. His full blood profile and hepatic functions are normal, and a lumbar puncture is unremarkable.

Choose...

A 30-year-old woman has two-sided basal ganglia calcification. She is being reviewed for kidney stones and polyuria.

Choose...

Explanation: Case 1 is suggestive of Huntington's disease, which can present with nonspecific psychiatric disturbances.

Case 2 is descriptive of complex partial seizures. These begin as simple partial seizures and progress to impairment of consciousness. There may also be an impairment of consciousness at the onset. Aggressive behaviour is also a possibility in patients with complex partial seizures. The lead about America is a red herring.

Case 3 describes Fahr's disease, which is caused by idiopathic progressive calcium deposition in the basal ganglia. Early-onset type (between the ages of 20 and 40) is associated with schizophreniform psychoses and catatonic symptoms. Onset between the ages of 40 and 60 is associated with dementia and choreoathetosis. Depression is also very common, but mania much less so. 50% of patients have psychiatric symptoms, which correlate with the degree of calcification. The most frequent clinical features are Parkinsonism, dystonia, tremor, gait disturbance and dysarthria with some cases exhibiting chorea, seizures and myoclonus. The term Fahr's syndrome is preferred if specific causes of calcium deposition in the basal ganglia such as hypoparathyroidism are diagnosed. Radiological basal ganglia calcification on CT without clinical features may occur at a rate of about 0.9% MRI shows hypointensity of the striatum. EEG is non-specific - may show diffuse changes.
Neuropsychiatry of the basal ganglia - Re: Cognition Health, http://www.re-cognitionhealth.com/dynamicdata/MovementDisorders.pdf (accessed March 31, 2015)

The correct answer is: A 44-year-old homeless man is brought into A&E with agitation and features of psychosis and depression. An EEG reveals diffuse attenuation in alpha activity, and his gait is abnormal. MRI shows atrophy of the striatum. – Huntington's disease, A 25-year-old man recently returned from Kansas. He presents with unprovoked aggression. Of late he experienced recurrent seizures followed by periods of impaired mentation. His full blood profile and hepatic functions are normal, and a lumbar puncture is unremarkable. – Complex partial seizures, A 30-year-old woman has two-sided basal ganglia calcification. She is being reviewed for kidney stones and polyuria. – Fahr's syndrome

Question 75

Not answered

Marked out of 3.00

Flag question

MoK Psychology EMI001

Principles of Learning Theory

Which type of learning process is involved in each of the following situations?

The fear that a child with a spider phobia shows towards insects in general.

Choose...

Humans usually react with a fear response to snakes

Choose...

Cars have the habit of flashing a seat belt warning sign until you tighten your seat belt.

Choose...

Stimulus preparedness (Seligman) explains why snake and spider phobia are commoner than shoe phobia or watch phobia. In evolutionary terms, the stimuli that were threatening to hunter-gatherer man has been hard-wired in our system to respond immediately - and phobia develops more readily for such 'prepared stimuli'.

Pavlovs' dog experiments were conducted using human subjects by Watson & Rayner and paved the way for founding the school of behaviourism. Watson produced 'phobia' in an infant called Little Albert. By exposing him to loud, frightening noise whenever he was shown a white rat, eventually Albert became fearful of the white rat, even when he heard no loud noise. A similar fear response was seen when any furry white object was shown to Albert. This 'spread' of associative learning from one stimulus to other is called stimulus generalisation.

Reinforcement can be positive or negative according to the way reinforcers are handled i.e. if something is given or taken away. If a particular behaviour increases by providing a reinforcer, then this is positive reinforcement. If a particular behaviour increases by removal of an aversive stimulus then this is negative reinforcement. Similarly punishment can also be positive (smack) or negative (take the chocolate away) though it is often used in positive sense.

The correct answer is: The fear that a child with a spider phobia shows towards insects in general. – Stimulus generalisation, Humans usually react with a fear response to snakes – Stimulus preparedness, Cars have the habit of flashing a seat belt warning sign until you tighten your seat belt. – Negative reinforcement

Question 76

Not answered

Marked out of 3.00

Flag question

MoK Assessment EMI010

Eponymous syndromes

Find one most appropriate diagnosis/eponym each for the syndromes described below:

Mr. Yardley is a prisoner awaiting trial and when he was asked 'What is the capital of England, he answered 'Paris'. Choose...

A 34-year-old gentleman is convinced that his wife is not his real wife and has been replaced by an imposter who looks just like her. Choose...

Progressive global cognitive impairment with a clinical course characterised by gradual intellectual decline, generalised slowing, and motor problems. MRI scan shows multiple microvascular infarcts of perforating vessels and white matter leukoaraiosis Choose...

Explanation:

Ganser symptom refers to the production of approximate solutions as seen in case 1. Here the patient gives repeated wrong answers to questions, which are nonetheless approximately right.

Capgras syndrome (case 2) is a delusional misidentification resulting in the belief that a person known to a patient has been replaced by a 'double' who appears identical though not the real person. Capgras syndrome is sometimes referred to as the illusion of doubles though it is a delusion. This syndrome was originally reported by Kahlbaum (1866) but most subsequent research is attributed to Capgras and colleagues (1923, 1924).

Case 3 is Binswanger's disease (progressive small vessel disease) - a subcortical dementia with a clinical course characterised by slow intellectual decline and generalised slowing. The clinical picture ma be dominated by the dementia, and there may be concomitant physical problems such as gait disorders or dysarthria. MRI scan shows small distinct infarcts (Lacunae) or more generalised white matter changes (Leukoariasis).

The correct answer is: Mr. Yardley is a prisoner awaiting trial and when he was asked 'What is the capital of England, he answered 'Paris'. – Ganser's syndrome, A 34-year-old gentleman is convinced that his wife is not his real wife and has been replaced by an imposter who looks just like her. – Capgras syndrome, Progressive global cognitive impairment with a clinical course characterised by gradual intellectual decline, generalised slowing, and motor problems. MRI scan shows multiple microvascular infarcts of perforating vessels and white matter leukoaraiosis – Binswanger's disease

Question 77

Not answered

Marked out of 3.00

Flag question

MoK Assessment EMI011

Descriptive features in psychiatry

Identify the terms used to describe each of the following:

A 70-year-old man lives in a supported accommodation. He goes out for shopping on his own but forgets a few things on his shopping list. When confronted he says that he has already got enough of these items in stock at his flat. Choose...

A 72-year-old male admitted to A&E, when asked what the month was, said 'April' and when asked what the year was, said 'April.' Choose...

A patient with a moderate learning disability is unable to express her emotional problems verbally. When asked if she feels depressed, she says 'yes'. When a picture book with various emotional states is shown, she admits having whatever emotion is shown first in the booklet at different times.

Choose...

Explanation:

Confabulation may be spontaneous where often elements of fantasy are found. It can also occur in an attempt to avoid embarrassment as in this case.

Perseveration is repetitive goal-directed verbal or motor behaviour where the response set does not change despite changes in the stimuli.

Case 3: This patient is highly suggestible as she answers yes to closed questions and admits to the emotions in any random picture shown to her.

The correct answer is: A 70-year-old man lives in a supported accommodation. He goes out for shopping on his own but forgets a few things on his shopping list. When confronted he says that he has already got enough of these items in stock at his flat. – Confabulation, A 72-year-old male admitted to A&E, when asked what the month was, said 'April' and when asked what the year was, said 'April.' – Perseveration, A patient with a moderate learning disability is unable to express her emotional problems verbally. When asked if she feels depressed, she says 'yes'. When a picture book with various emotional states is shown, she admits having whatever emotion is shown first in the booklet at different times. – Suggestibility

Question 78

Not answered

Marked out of 3.00

Flag question

MoK Assessment EMI016

Personality disorders - diagnosis

Choose one option each from above list:

A 35-year-old man refuses to provide answers to standard questions during an initial clerking and threatens to stop the interview if recording his telephone number is insisted. It seems that he has taken many hospitals to court for suspicions about how his personal data has been handled.

Choose...

A 33-year-old night security at a local hospital prefers to be alone whenever possible. He has no friends, and he does not socialise. He does not keep update with current affairs and has no sexual interests. He spends most time day dreaming.

Choose...

A 21-year-old lady spends excessive time seeking attention by making herself attractive. She wears the 'fashion of the week' all the time and show shallow but labile affect. She comes across as being very impressionistic.

Choose...

Explanation:

Paranoid personality disorder is "characterised by pervasive suspiciousness, mistrust, and hypersensitivity to criticism and hostility". As a rule, paranoid people are ready to counter-attack, provoking repeated confrontations and induce hostility and resentment in others.

Schizoid personality disorder is "characterised by a persistent pattern of social withdrawal, discomfort in social interactions and are introverted, isolated and lonely".

Histrionic personality disorder is "characterised by excessive emotionality and attention seeking, and by dramatic, colourful, and extroverted behaviour that is egocentric, dependent, and demanding in interpersonal relationships". (All descriptions from New Oxford Textbook of Psychiatry)

The correct answer is: A 35-year-old man refuses to provide answers to standard questions during an initial clerking and threatens to stop the interview if recording his telephone number is insisted. It seems that he has taken many hospitals to court for suspicions about how his personal data has been handled. – Paranoid personality disorder, A 33-year-old night security at a local hospital prefers to be alone whenever possible. He has no friends, and he does not socialise. He does not keep update with current affairs and

has no sexual interests. He spends most time day dreaming. – Schizoid personality disorder, A 21-year-old lady spends excessive time seeking attention by making herself attractive. She wears the 'fashion of the week' all the time and show shallow but labile affect. She comes across as being very impressionistic. – Histrionic personality disorder

Finish review