Mock Paper A(1)

Started on	Thursday, 21 June 2018, 8:15 AM
State	Finished
Completed on	Thursday, 21 June 2018, 8:15 AM
Time taken	12 secs
Grade	**0.00** out of 100.00
Feedback	You are on the right track. Could you spend some more time on your revision?

Question 1
Not answered
Marked out of 1.00
Flag question

MoK Classification 012
Which of the following is a cardinal feature of depression?

Select one:
- Anhedonia
- Pessimism
- Suicidality
- Somatisation
- Guilt

Your answer is incorrect.

The role of anhedonia in the diagnosis of melancholia of the past years was considered to be crucial. Klein regarded anhedonia to be a symptom of depressive illness and probably the best clinical marker predicting response to antidepressant drugs. In 1980, the revised DSM presented the concept of anhedonia as one of the two cardinal symptoms of major depression (Anhedonia: a neglected symptom of psychopathology, http://www.ncbi.nlm.nih.gov/pubmed/8134519).

The correct answer is: Anhedonia

Question 2
Not answered
Marked out of 1.00
Flag question

MoK Classification 022
Which of the following clinical features suggests a diagnosis of mania rather than hypomania?

Select one:
- Decreased need for sleep
- Psychomotor agitation
- Inflated self esteem
- Flight of ideas
- Delusions of grandiosity

Your answer is incorrect.

Psychotic symptoms are suggestive of mania rather than hypomania.

The correct answer is: Delusions of grandiosity

Question 3
Not answered
Marked out of 1.00
Flag question

MoK Classification 025
Mrs Pinto lost her husband 9 months ago. She had continued attending her bingo sessions and cathedral groups since his death, but recently she has had pervasive guilt and suicidal thoughts. What is the most appropriate diagnosis?

Select one:
- Depression
- Paraphrenia
- Delusional disorder
- Bereavement
- Dementia

Your answer is incorrect.

The presence of delusional guilt or intense suicidal ideas indicates a major depressive episode rather than bereavement, especially after nine months of loss.

The correct answer is: Depression

Question 4
Not answered
Marked out of 1.00
Flag question

MoK Classification 027
A 37-year-old woman was admitted to the medical ward with history of sudden onset aphonia. Physical examination and investigations were normal. Her boyfriend has recently invited his mother to stay with them permanently, as his mother is feeling lonely and isolated. What is the most likely diagnosis according to ICD-10 criteria?

Select one:
- Adjustment disorder
- Munchausen syndrome
- Hypochondriasis
- Somatisation disorder
- Dissociative motor disorder

Your answer is incorrect.

Functional weakness describes weakness which is inconsistent and incongruent with a neurological disease. It is also referred to as motor conversion disorder (DSM-IV), dissociative motor disorder (ICD-10) and 'psychogenic' paralysis. Studies of aetiology have focused on risk factors such as childhood adversity and life events.(Stone et al. 2011)

The correct answer is: Dissociative motor disorder

Question 5

Not answered

Marked out of 1.00

Flag question

MoK Classification 032

A soldier is referred with symptoms of PTSD following return from Iraq. Which of the following feature is not consistent with the diagnosis of PTSD?

Select one:

- He is not able to enjoy watching television
- He is not able to share his war experience with his close friends
- He sleeps in his day time clothes
- He is keen to go to Iraq again
- He has sustained only mild injuries physicall

Your answer is incorrect.

Avoidance is a part of PTSD - wanting to go back seems odd.

The correct answer is: He is keen to go to Iraq again

Question 6

Not answered

Marked out of 1.00

Flag question

MoK Classification 034

Mrs Broadbent has been diagnosed with compensation neurosis. Which of the following is true regarding this?

Select one:

- Severe difficulties with sleep are noted
- Most patients do not return to work even after settlement
- Higher the severity of injury, greater the symptoms of compensation neurosis
- Absence of frontal headaches
- Improvements tend to occur after financial settlement

Your answer is incorrect.

Compensation neurosis is not an ICD diagnosis; its validity is questionable. Patients show scant improvement even after financial settlement. A follow-up study of 760 litigants found that, of the 264 subjects who were not working at the time of conclusion of litigation and who could be traced, 198 (75%) were not working after an average of 23 months following the finalization of their cases. (Mendelson, 1995. Compensation neurosis revisited. Journal of Psychosomatic Research, 39, pp. 695-706)

The correct answer is: Most patients do not return to work even after settlement

Question 7

Not answered

Marked out of 1.00

Flag question

MoK Classification 037

An elderly man presents with fluctuation in consciousness and visual hallucinations. His wife reports that he is not able to turn in his bed properly. You suspect

Select one:

- Vascular dementia
- Huntington's disease
- Lewy body dementia
- Wilson disease
- Alzheimer's dementia

Your answer is incorrect.

Not able to turn in bed indicates axial rigidity that may be related to Parkinsonism. This patient has Lewy body dementia.

The correct answer is: Lewy body dementia

Question 8

Not answered

Marked out of 1.00

Flag question

MoK Classification 039

Which of the following is consistent with a diagnosis of bipolar affective disorder, currently hypomanic episode?

Select one:

- Mood incongruent delusions
- 2 weeks duration of symptoms
- Fleeting hallucinations
- Minimal disturbance of psychosocial function
- Mood congruent delusions

Your answer is incorrect.

The presence of delusions or hallucinations should prompt to make a diagnosis of mania, not hypomania. By definition, the impairment in psychosocial functioning is of higher degree in mania than hypomania.

The correct answer is: Minimal disturbance of psychosocial function

Question 9

MoK Classification 041

Not answered

Marked out of 1.00

Flag question

Which of the following features is included in ICD-10 diagnosis for generalised anxiety disorder?

Select one:

- Nightmares
- Panic attacks
- Depression
- Low self confidence
- Motor tension

Your answer is incorrect.

To diagnose GAD, the sufferer must have primary symptoms of anxiety most days for at least several weeks at a time, and usually for several months. These symptoms should usually involve elements of 1. apprehension (worries about future misfortunes, feeling 'on edge', difficulty in concentrating, etc.); 2. motor tension (restless fidgeting, tension headaches, trembling, inability to relax); and 3. autonomic overactivity (lightheadedness, sweating, tachycardia or tachypnoea, epigastric discomfort, dizziness, dry mouth, etc.).

The correct answer is: Motor tension

Question 10

Not answered

Marked out of 1.00

Flag question

MoK Classification 042

A 33-year-old lady presents with excessive day time sleepiness and auditory hallucinations when waking from sleep. She also recalls episodes of feeling 'pinned down' to her bed even after waking up. The likely diagnosis is

Select one:

- Schizophrenia
- REM behavioural disorder
- Narcolepsy
- Sleep terror
- Primary hypersomnia

Your answer is incorrect.

The symptoms described here are hypnopompic hallucinations and sleep paralysis. These are characteristic of narcolepsy.

The correct answer is: Narcolepsy

Question 11

Not answered

Marked out of 1.00

Flag question

MoK Classification 043

Which one of the following is a feature of atypical Anorexia Nervosa?

Select one:

- Depressed mood
- Normal menstruation
- Binge eating
- Purging behaviour
- Restricting behaviour

Your answer is incorrect.

Atypical anorexia refers to an absence of one or more core symptoms needed to diagnose anorexia nervosa i.e. BMI criteria, body image criteria or amenorrhea.

The correct answer is: Normal menstruation

Question 12

Not answered

Marked out of 1.00

Flag question

MoK Classification 044

Bleuler discarded the term dementia praecox primarily because

Select one:

- Mania was commonly coexistent with dementia praecox
- Not all cases began during adolescence
- Praecox affect was rarely seen in the early cases
- No treatment was available for dementia praecox
- No cognitive impairment was noted in the early cases

Your answer is incorrect.

Kraepelin's original formulation of 'dementia praecox' was rejected by Bleuler, who favoured 'the group of schizophrenias' to describe the psychotic condition. Bleuler argued that the term dementia praecox was based on the notions of the age of onset and the illness course, for which convincing evidence was lacking. Bleuler emphasized the multiplicity of symptoms and considered the most important symptom to be fragmentation in the formulation and expression of thought, referred to as 'loosening of associations.'

The correct answer is: Not all cases began during adolescence

Question 13

Not answered

Marked out of 1.00

Flag question

MoK Classification 045

Mrs. Z arrives at work at least an hour earlier than anyone else to plan the day. She becomes upset if the cleaners have moved anything on her desk and in her office, which is very tidy. Her home and car are spotlessly clean. This is best described as;

Select one:

- Obsessional personality trait
- Normal phenomenon
- Hoarding behaviour

- Obsessive compulsive disorder
- Compulsive behaviour

Your answer is incorrect.

Obsessive-compulsive personality disorder (OCPD) is a personality disorder characterized by excessive perfectionism and a need for control over one's environment. People with OCPD tend to be reliable and orderly, but also inflexible and unable to surrender control. This makes it hard for people with this condition to express their feelings, have close relationships and enjoy their successes. This lady is exhibiting some of the above traits.

The correct answer is: Obsessional personality trait

Question 14
Not answered
Marked out of 1.00
Flag question

MoK Classification 046
Which of the following is true with respect to adjustment disorder?

Select one:
- usually requires prolonged treatment
- correlates with the severity of the stressor
- occurs more often in males than in females
- occurs in all age groups
- it is a type of bereavement

Your answer is incorrect.

Although an adjustment disorder can occur at any age, it is more common at times in life when major transitions occur, such as adolescence, mid-life, and late-life (Mental Health: Adjustment Disorder - WebMD, http://www.webmd.com/mental-health/mental-health-adjustment-disorder).

The correct answer is: occurs in all age groups

Question 15
Not answered
Marked out of 1.00
Flag question

MoK Classification 047
With respect to models of diagnostic classification, which of the following is correct?

Select one:
- Ontological theories suggest that mental state does not exist by virtue of its abstract nature
- Fulford claims that psychiatric diagnoses are not value laden
- ICD-10 is entirely atheoretical
- Pheneticists stress on the natural existence of hierarchical structure among disease entities
- Categorical entities are not used in modern classification

Your answer is incorrect.

In contrast to the essentialist strategy that treats diseases as entities that exist in nature and form natural groups in essence, phenetic approach, (also called numerical taxonomy) shifts the emphasis to the systematic description of the observed disease presentations to enable manual classification. This approach treats all characters and attributes as having equal weight. While ICD and DSM are atheoretical in terms of assuming aetiology and causal models, they are not entirely atheoretical in descriptions used (e.g. dissociative disorder, conversion, etc. employ psychodynamic concepts).

The correct answer is: Ontological theories suggest that mental state does not exist by virtue of its abstract nature

Question 16
Not answered
Marked out of 1.00
Flag question

MoK Clinical Examination 016
Which of the following assessment is not a part of MMSE?

Select one:
- Repeating phrase
- Clock Drawing
- Motor sequence
- Testing attention
- Copying polygons

Your answer is incorrect.

Clock drawing provides complementary information, but not a part of MMSE.

The correct answer is: Clock Drawing

Question 17
Not answered
Marked out of 1.00
Flag question

MoK Clinical Examination 018
Which of the following is a frontal lobe test?

Select one:
- Repertory Grid
- Line bisection test
- Star cancellation test
- Cognitive Estimates Test
- Bar chart test

Your answer is incorrect.

Cognitive estimates test is a part of the frontal assessment battery. It includes questions such as 'What is the average height of a British woman?'

The correct answer is: Cognitive Estimates Test

Question 18
Not answered
Marked out of 1.00
Flag question

MoK Clinical Examination 020
Subtracting 7 out of 100 repeatedly (serial sevens test) is most useful to test one's

Select one:
- Attention
- Alertness
- Concentration
- Working memory
- Motivation

Your answer is incorrect.
Concentration is also known as sustained attention.
The correct answer is: Concentration

Question 19
Not answered
Marked out of 1.00
Flag question

MoK Clinical Examination 023
A 56-year-old woman is brought to the hospital with a stroke. A CT scan shows evidence of left occipital lobe damage. Which of the following is most likely to be seen?

Select one:
- Quadrantanopsia
- Cortical blindness
- Anosmia
- Bitemporal hemianopsia
- Apraxia

Your answer is incorrect.
Cortical blindness is the total or partial loss of vision in a normal-appearing eye caused by damage to the occipital cortex of the brain. Cortical blindness can be acquired or congenital, and may also be transient in certain instances. Acquired cortical blindness is most often caused by loss of blood flow to the occipital cortex from either unilateral or bilateral posterior cerebral artery blockage due to ischemic stroke. In most cases, the patient may recover some of their vision
The correct answer is: Cortical blindness

Question 20
Not answered
Marked out of 1.00
Flag question

MoK Clinical Examination 024
A boxer is knocked out by a blow to his lateral skull. Despite recovering after a few minutes, and being asymptomatic for the next 12 hours, he develops severe headache, vomiting and reduced consciousness. Which of the following is the most indicated investigation?

Select one:
- X ray skull
- SPECT scan
- EEG
- MRI brain
- CT scan brain

Your answer is incorrect.
In cases of suspected bony fracture and fresh hemorrhage, a CT scan is more useful than plain X-ray, which cannot pick up the bleed or MRI brain which is slower to obtain and may miss fresh bleeding.
The correct answer is: CT scan brain

Question 21
Not answered
Marked out of 1.00
Flag question

MoK Clinical Examination 025
Hyperhomocysteinemia is associated with increased risk for which of the following conditions?

Select one:
- Alcoholism
- Borderline personality disorder
- Panic disorder
- Self mutilation
- Dementia

Your answer is incorrect.
An increased plasma homocysteine level is a strong, independent risk factor for the development of dementia and Alzheimer's disease. N Engl J Med 2002; 346:476-483
The correct answer is: Dementia

Question 22
Not answered
Marked out of 1.00
Flag question

MoK Clinical Examination 026
Which of following is not commonly seen in nicotinic acid deficiency?

Select one:
- Insomnia

- Diarrhoea
- Glossitis
- Dermatitis
- Dementia

Your answer is incorrect.

Dementia occurs only in an extreme form of nicotinic acid deficiency. Commonly, acute deficiency causes pellagra characterized by 3 Ds - diarrhoea, dermatitis and disturbed mental functions.

The correct answer is: Dementia

Question 23

Not answered

Marked out of 1.00

Flag question

MoK Clinical Examination 027
Which of the following best describes ideomotor apraxia?

Select one:

- Inability to coordinate and carry out facial and lip movements
- Inability to construct elements to a meaningful whole
- Difficulty in learning new information
- Inability to perform a multiple-step task.
- Difficulty in copying skilled movements

Your answer is incorrect.

Ideomotor apraxia is a disorder of goal-directed movement. The patient knows what to do but not how to do it. There is an inability to mimic a learned motor task or behavior, such as combing one's hair. Ideomotor apraxia results from damage to the dominant parietal lobe. Abnormalities include (i) abnormal amplitude of motor movements (ii) body-part-as-object substitution, e.g. the patient uses his own finger to represent a toothbrush when asked to brush his teeth; (iii) abnormal orientation of body parts when attempting to perform an action.

The correct answer is: Difficulty in copying skilled movements

Question 24

Not answered

Marked out of 1.00

Flag question

MoK Clinical Examination 028
While screening for alcohol use problems, which of the following questions is the best one to be asked first?

Select one:

- Have you ever felt bad or guilty about your drinking?
- Have people annoyed you by getting on your case about your drinking?
- Do you enjoy a drink now and then?
- Have you ever needed to take a drink first thing in the morning?
- Have you felt you should cut down on your drinking?

Your answer is incorrect.

This question is not about CAGE. It is about sensitive interview techniques. A sensitive topic must be gradually introduced with non-judgemental tone.

The correct answer is: Do you enjoy a drink now and then?

Question 25

Not answered

Marked out of 1.00

Flag question

MoK Clinical Examination 029
A patient with schizoaffective disorder is on both lithium and risperidone. He develops tremors as a side effect of his treatment. Which of the following will help detecting the offending agent?

Select one:

- Finger nose test
- Peripheral visual field
- Examination of tone of joints
- Frontal release sign
- Physical examination for tendon reflexes

Your answer is incorrect.

EPSE with risperidone can cause tremors (Parkinsonian) and rigidity, unlike Lithium tremor where one would not expect rigidity.

The correct answer is: Examination of tone of joints

Question 26

Not answered

Marked out of 1.00

Flag question

MoK Descriptive Psychopathology 018
When asked where do he lived when he was a child, a patient replied "Winchester. Manchester City was my favourite football team." The type of speech abnormality exhibited is

Select one:

- Clang association
- Pressured speech
- Tangential
- Coprolalia
- Circumstantial

Your answer is incorrect.

This is an example of clang association, which is an abnormality of speech where the connection between words is their sound rather than their meaning. It may often occur in patients with mania.

The correct answer is: Clang association

Question 27
Not answered
Marked out of 1.00
Flag question

MoK Descriptive Psychopathology 025
Capgras syndrome is:

Select one:
- A form of negative autoscopy
- A form of autoscopy
- Also called the phantom mirror image
- A perceptual abnormality
- Delusional misidentification

Your answer is incorrect.

Capgras syndrome is included in delusional misidentification syndromes along with reduplicative paramnesia, Fregoli's syndrome and intermetamorphosis. It is not a perceptual error. Phantom mirror image refers to autoscopy which is a perceptual phenomenon.

The correct answer is: Delusional misidentification

Question 28
Not answered
Marked out of 1.00
Flag question

MoK Descriptive Psychopathology 027
A 20-year-old man is unable to express his problem coherently but appears very confused and feels that 'something is happening but not sure what it is'. This sense of perplexity and uncertainty that exists during a prodrome of psychosis may be linked most appropriately to which of the following experiences?

Select one:
- Negative symptoms
- Depression
- Delusional memory
- Delusional mood
- Autochthonous delusions

Your answer is incorrect.

Delusional mood refers to a state of perplexity in which the patient has some sense of some inexplicable change in his environment. He senses 'something going on' which he cannot identify, but which has a peculiar significance for him.

The correct answer is: Delusional mood

Question 29
Not answered
Marked out of 1.00
Flag question

MoK Descriptive Psychopathology 030
A 23-year-old man notices the inscribed image of the Queen on a coin, and says "I know now, my mother is an alien". This is an example of

Select one:
- Ideas of reference
- Delusional mood
- Delusional memory
- Delusional misidentification
- Delusional perception

Your answer is incorrect.

This is delusional perception that is a primary delusion, which is recalled as having arisen as a result of a perception. The percept is a real external object, not a hallucinatory experience.

The correct answer is: Delusional perception

Question 30
Not answered
Marked out of 1.00
Flag question

MoK Descriptive Psychopathology 033
A 43-year-old woman, who was involved in a serious flight accident, felt at the time of the accident as if the world had ceased to exist. For some time she lost her sense of time perception and felt detached from the rescue proceedings around her. This is known as:

Select one:
- Nihilistic delusion
- Derealisation
- Suicidal thought
- Depersonalisation
- None of the above

Your answer is incorrect.

In depersonalisation, an unpleasant sense of detachment can occur. Disturbed time perception and feelings of 'unreality' are also noted. In isolated derealisation (which is rare in the absence of depersonalisation), one may not have a personal sense of detachment but may feel that the world around them is unreal, sometimes 'robotic', made of cardboards!

The correct answer is: Depersonalisation

Question 31
Not answered
Marked out of 1.00
Flag question

MoK Descriptive Psychopathology 034
One among the following statements is NOT true about depersonalisation?

Select one:
- Depersonalisation is easily distinguished from derealizatio
- Depersonalisation is always subjective
- Depersonalisation is frequently accompanied by the symptom of derealization
- Depersonalisation is a frequent experience of normal healthy people
- Affect is invariably involved in Depersonalisation

Your answer is incorrect.
It is hard to accurately distinguish the two clinically.
The correct answer is: Depersonalisation is easily distinguished from derealizatio

Question 32
Not answered
Marked out of 1.00
Flag question

MoK Descriptive Psychopathology 035
Of various thought disorders classified by Andreasen, which of the following are more common in schizophrenia than mania?

Select one:
- Derailment
- Circumstantiality
- Neologisms
- Clanging
- Flight of ideas

Your answer is incorrect.
Of various thought disorders classified by Andreasen, clanging and flight are more common in mania while derailment (loosening) and thought blocking and to some extent tangentiality, poverty of content of speech are often seen in schizophrenia - other items were largely non-specific
The correct answer is: Derailment

Question 33
Not answered
Marked out of 1.00
Flag question

MoK Descriptive Psychopathology 038
The phenomenon in which a subject perceives other people's heads to be as big as their actual body size can be termed as

Select one:
- Pareidolia
- Dysmegalopsia
- Lilliputian Hallucinations
- Micropsia
- Macropsia

Your answer is incorrect.
Dysmegalopsia is characterised by alterations in the symmetry/size of perceived objects.
The correct answer is: Dysmegalopsia

Question 34
Not answered
Marked out of 1.00
Flag question

MoK Descriptive Psychopathology 039
A patient repeats whatever the examiner says. He is exhibiting

Select one:
- Perseveration
- Logoclonia
- Parapraxis
- Stereotypy
- Echolalia

Your answer is incorrect.
Echolalia is an echophenomenon seen in catatonia. Other echophenomena are echopraxia (repeating what you do) and mirror movements (doing as you do at same time)
The correct answer is: Echolalia

Question 35
Not answered
Marked out of 1.00
Flag question

MoK Descriptive Psychopathology 040
The nature of speech produced in Broca's aphasia is

Select one:
- Fluent and intelligible
- Impaired speech comprehension
- Fluent but unintelligible
- Effortful but agrammatism is seen
- Effortful but with grammatical use of language

Your answer is incorrect.

Broca's aphasia is characterised by retained use of nouns and verbs, but the patient may have lost all forms of pronouns, articles, and conjunctions and hence may struggle to speak more than one word at a time even with enormous effort.

The correct answer is: Effortful but agrammatism is seen

Question 36
Not answered
Marked out of 1.00
Flag question

MoK Descriptive Psychopathology 041
Mr X was admitted to the psychiatric unit with a history of elated mood, overactivity, restlessness, grandiose delusions, disinhibited behaviour and was diagnosed with mania. A student nurse observed that he also experienced third person auditory hallucinations and wondered whether first rank symptoms could occur in manic patients. He wanted to clarify this with the treating psychiatric doctor in the team. Choose the most appropriate statement from the following:

Select one:

- First rank symptoms never occur in manic patients
- First rank symptoms may present in 10-25% of manic patients
- First rank symptoms are very rarely seen in manic patients
- First rank symptoms are present in all cases of mania with psychosis
- First rank symptoms are present in nearly half of all patients with mania

Your answer is incorrect.

First rank symptoms are not pathognomonic of schizophrenia. They are observed to occur on bipolar mania, epilepsy (TLE) and also in organic psychotic states.

The correct answer is: First rank symptoms may present in 10-25% of manic patients

Question 37
Not answered
Marked out of 1.00
Flag question

MoK Dynamic Psychopathology 001
Which of the following defence mechanisms is most prominent in subjects with narcissism?

Select one:

- Idealisation
- Discrimination
- Acting out
- Anticipation
- Projection

Your answer is incorrect.

Projection is commonly seen in narcissism. It involves taking one's own unacceptable qualities or feelings and ascribing them to other people. For example, a narcissist with a strong dislike for someone, might instead believe that the other person does not like him.

The correct answer is: Projection

Question 38
Not answered
Marked out of 1.00
Flag question

MoK Dynamic Psychopathology 002
Avoidance is a defence mechanism most commonly used in

Select one:

- OCD
- Agoraphobia
- Hypochondriasis
- Somatoform disorder
- Depression

Your answer is incorrect.

Avoidance is a powerful cognitive and behavioural strategy that serves as a negative reinforcer to reduce anxiety associated with agoraphobia. (Most questions on defence mechanism/cognitive styles are based on psychotherapeutic theoretical grounding; not on quantitative evidence regarding frequency and prevalence.)

The correct answer is: Agoraphobia

Question 39
Not answered
Marked out of 1.00
Flag question

MoK Dynamic Psychopathology 003
You are a CT1 psychiatric trainee. One of your patients on the ward who is a 62-year-old man has an acute onset chest pain and sweating. Accompanying ECG showed ST elevation in leads V1-4. What is the abnormality?

Select one:

- Sick sinus syndrome
- Anterior MI
- First degree block
- Inferior MI
- Prolonged QTc

Your answer is incorrect.

ST elevation and/or Q waves in V1-4/V5 is suggestive of anterior wall MI

The correct answer is: Anterior MI

Question 40
Not answered
Marked out of 1.00
Flag question

MoK Dynamic Psychopathology 004
Which of the following defence mechanisms involves realistic planning for future discomfort?

Select one:
- Depreciation
- Anticipation
- Altruism
- Reaction formation
- Denial

Your answer is incorrect.
Anticipation is a mature defence mechanism
The correct answer is: Anticipation

Question 41
Not answered
Marked out of 1.00
Flag question

MoK Dynamic Psychopathology 005
The expression of ideas, thoughts, and suppressed material that is accompanied by an emotional response producing a state of relief in the patient is called as

Select one:
- Democratisation
- Modelling
- Mirroring
- Rapprochement
- Catharsis

Your answer is incorrect.
Catharsis is the expression of ideas, thoughts and suppressed material that is accompanied by an emotional response that produces a state of relief in the patient
The correct answer is: Catharsis

Question 42
Not answered
Marked out of 1.00
Flag question

MoK Dynamic Psychopathology 008
Which of the following ego defence mechanism has been put forth as an explanation for phobia?

Select one:
- Denial
- Intellectualization
- Reaction formation
- Splitting
- Displacement

Your answer is incorrect.
Freud invoked the mechanism of displacement to explain the fear of horses in Little Hans. The fear of his father was displaced on to horses.
The correct answer is: Displacement

Question 43
Not answered
Marked out of 1.00
Flag question

MoK Basic Psychology 005
The span of immediate memory is estimated to be around

Select one:
- 8±2
- 9±2
- 5±2
- 3±2
- 7±2

Your answer is incorrect.
The 'magical number seven' (G. A. Miller; PA, Vol 31:2914,) demonstrated the concept of limited channel capacity and specified 7 (+/-2) as the immediate memory span. Though the span is limited, the immediate memory capacity is relatively insensitive to the amount of information per item. Miller, and later Baddeley emphasized the importance of the recoding of information (the concept of chunking).
The correct answer is: 7±2

Question 44
Not answered
Marked out of 1.00
Flag question

MoK Basic Psychology 009
Which one of the following theories is not associated with encoding failure of forgetting?

Select one:
- State dependent memory
- Blocking
- Failure of prospective memory
- Context dependent memory

- All of the above

Your answer is incorrect.

All the above theories are associated with retrieval failure of forgetting and not due to encoding failure. Encoding failure: Forgetting over time may occur due to defective or impaired initial encoding of information into memory. It may also be due to shallow processing of information. Retrieval failure; This is the commonest cause of failure to access long term memory. Although the memory trace is there, the route to retrieving is blocked, which can be facilitated by cues.

The correct answer is: All of the above

Question 45
Not answered
Marked out of 1.00
Flag question

MoK Basic Psychology 011
Abnormalities in Stroop test suggests

Select one:

- Cortical blindness
- Lesion of optic chiasma
- Attentional bias
- Short term memory disturbance
- Damage to occipital cortex

Your answer is incorrect.

Abnormal Stroop effect with the loss of appropriate inhibitory control is seen in frontal lobe lesions. Abnormalities in modified emotional stroop interference effects (e.g. on threat words in anxious subjects on the emotional Stroop task) have generally been interpreted as evidence for mood-congruent attentional bias in various mental states.

The correct answer is: Attentional bias

Question 46
Not answered
Marked out of 1.00
Flag question

MoK Basic Psychology 017
Tom is a 4-year-old boy. He has a rabbit at home. When he reached out to stroke the rabbit, a hammer is struck against an iron rod causing a loud noise, which frightens him. This is repeated 5 times. As a result of this, the sight of the rabbit on its own now frightens Tom. This type of conditioning is called

Select one:

- Classical conditioning
- Operant conditioning
- Observational learning
- Stimulus discrimination
- Trace conditioning

Your answer is incorrect.

Classical conditioning is a process of behaviour modification by which a subject comes to respond in a desired manner to a previously neutral stimulus that has been repeatedly presented along with an unconditioned stimulus that elicits the desired response

The correct answer is: Classical conditioning

Question 47
Not answered
Marked out of 1.00
Flag question

MoK Basic Psychology 018
The process which leads to the transfer of information from short-term memory to long-term memory is called:

Select one:

- Rehearsal
- Encoding
- Consolidation
- Displacement
- Interference

Your answer is incorrect.

The transfer of information from short to long term, more permanent storage, is called memory consolidation. This can be facilitated by motivation (conscious effort) or association or elaboration of meaning. Rehearsal or repetition also facilitates the transfer of items in short term memory to long term storage.

The correct answer is: Consolidation

Question 48
Not answered
Marked out of 1.00
Flag question

MoK Basic Psychology 023
All of the following are tests of recent memory except

Select one:

- Name and address recall task
- Rey-osterrieth complex figure test
- Three word learning task
- Wechsler memory scale
- Digit span

Your answer is incorrect.

Digit span is the commonest test of auditory, verbal working memory. Both forward and backward digit span are tested in routine clinical practice. The normal range of digit forwards is 6+/-1 and for reverse digit span is 5+/-1 Three word learning task (E.g. apple, table, penny) is a test of Anterograde memory and learning. Name and address recall task (7 items) is the commonest test of recent memory. Here the subject is asked to recall as many items, without prompts, in five or ten minutes.

The correct answer is: Digit span

Question 49
Not answered
Marked out of 1.00
Flag question

MoK Basic Psychology 028
A middle-aged woman treated for agoraphobia has improved with repeated exposure to crowds. Which psychological process is involved here?

Select one:
- Instrumental conditioning
- Massed practices
- Cognitive restructuring
- Counter-conditioning
- Extinction

Your answer is incorrect.

Extinction is the conditioning phenomenon in which a previously learned response to a cue is reduced when the cue is presented in the absence of the previously paired stimulus

The correct answer is: Extinction

Question 50
Not answered
Marked out of 1.00
Flag question

MoK Basic Psychology 029
If a parent intentionally ignores a child's misbehaviour, this is an example of which operant conditioning principle?

Select one:
- Positive reinforcement
- Punishment
- Extinction.
- Negative reinforcement
- Shaping

Your answer is incorrect.

The attention given to a misbehaving child might positively reinforce the unwanted behaviour. By ignoring the misbehaving child, this positive reinforcement is withdrawn or extinguished (extinction). Negative reinforcement, on the other hand, is the removal of an undesirable effect that increases the probability of a target behaviour occurring in the future (e.g. flashing red lights are activated when a person starts a car without wearing seat belts. This is an undesirable effect. By buckling the seat belt, one can remove these aversive antecedents. Thus the target behaviour of 'buckling up' is reinforced by the 'removal' of the unwanted flashes of red light).

The correct answer is: Extinction.

Question 51
Not answered
Marked out of 1.00
Flag question

MoK Basic Psychology 043
Whose theory of emotions proposed that emotion is secondary to physiological changes and that there is a distinct pattern of changes for different emotions?

Select one:
- Cannon baird theory
- Cognitive appraisal theory
- James Lange theory
- Schema theory
- None of the above

Your answer is incorrect.

According to James-Lange theory emotion, is secondary to physiological changes and that there is a distinct pattern of changes for different emotions. The Cannon-Baird theory of emotion states that the subjective experience of emotions is independent of physiological changes. Any stimulus is processed by the thalamus that sends impulses to both the cortex and the hypothalamus. The cortex is responsible for emotions while the hypothalamus is responsible for physiological changes. The cognitive appraisal conscious or unconscious always precedes emotional experience. Lazarus proposed this theory.

The correct answer is: James Lange theory

Question 52
Not answered
Marked out of 1.00
Flag question

MoK Basic Psychology 044
Who postulated that thinking occurs peripherally in the speech apparatus?

Select one:
- Sternberg
- John Watson
- Whorf
- Spearman
- Maslow

Your answer is incorrect.

The Linguistic relativity hypothesis also called as the Whorfian hypothesis, places language as primary and thoughts as secondary. But in peripheralism, John Watson postulated that thinking occurs peripherally in the speech apparatus.

The correct answer is: John Watson

Question 53
Not answered
Marked out of 1.00
Flag question

MoK Basic Psychology 045
Which of the following are correctly paired with the concepts they introduced?

Select one:
- Freud : negative and positive symptoms
- Jung : introversion
- Eugen Bleuler : dementia praecox
- Manfred Bleuler : dementia praecox
- Charcot : dissociation

Your answer is incorrect.

Jung developed a personality typology that distinguished between introversion and extroversion. Introverts are people who prefer their internal world of thoughts, feelings, fantasies, dreams, and so on, while extroverts prefer the external world of things and people and activities.

The correct answer is: Jung : introversion

Question 54
Not answered
Marked out of 1.00
Flag question

MoK Social Psychology 004
Which one among the following is not one of the Gestalt principles of perception?

Select one:
- Proximity
- Similarity
- Closure
- Continuity
- Familiarity

Your answer is incorrect.

Gestalt law of perceptual organization includes Proximity: Objects close to each other are perceived as one figure, Closure: Incompletely closed figures are perceived as fully closed, Continuity: Continuous items are perceived together, Similarity: Similar items are grouped together based on colour or shape, etc. and Common fate: Things moving together are perceived together.

The correct answer is: Familiarity

Question 55
Not answered
Marked out of 1.00
Flag question

MoK Social Psychology 006
Beautiful people are generally viewed as more kind, convincing and effective. This is called

Select one:
- Barnum effect
- Fundamental attribution error
- Halo effect
- Pygmalion effect
- Hawthorne effect

Your answer is incorrect.

Halo effect refers to our tendency to allow one specific trait or our overall impression of a person to influence our judgment of their other traits. This is also called the physical attractiveness stereotype or "what is beautiful is good" principle.

The correct answer is: Halo effect

Question 56
Not answered
Marked out of 1.00
Flag question

MoK Sociocultural Psychiatry 014
Which of the following ethical principles does the issue of capacity to make treatment decisions uphold?

Select one:
- Non maleficience
- Justice
- Confidentiality
- Autonomy
- Beneficience

Your answer is incorrect.

Individual freedom is based on the principle of autonomy in modern medical ethics. It is the principle that a person should be free to make his or her own decisions.

The correct answer is: Autonomy

Question 57
Not answered
Marked out of 1.00
Flag question

MoK Sociocultural Psychiatry 019
Under the principle of necessity which of the following can be done without patient's consent?

Select one:
- Covert administration of medication in a competent non consenting adult
- A psychiatric inpatient who needs overnight detention as he attempts to leave
- Sterilisation of a patient with repeated postpartum mania

- Liver transplantation to a patient who consumed overdose of paracetamol
- Blood sample drawn to confirm suspected myocardial infarction in a schizophrenia patient

Your answer is incorrect.

Emergency interventions for life-saving measures could be carried out under common law or necessity principle. Irreversible or long term procedures cannot be carried out without formalised legal procedures.

The correct answer is: Blood sample drawn to confirm suspected myocardial infarction in a schizophrenia patient

Question 58
Not answered
Marked out of 1.00
Flag question

MoK Sociocultural Psychiatry 022
Tarasoff case is related to which of the following ethical principles?

Select one:
- Autonomy
- Beneficience
- Confidentiality
- Justice
- Capacity

Your answer is incorrect.

Tarasoff case refers to US federal law and highlights the duty to warn third parties who may be directly harmed by a patient, in addition to a doctor's duty of care towards patients. Tatiana Tarasoff was killed by Prasanjit Poddar, who divulged his intentions to his therapist before the homicide. Though the therapist arranged to admit him, no one directly warned Tatiana Tarasoff. Following his discharge from a brief hospitalization, Poddar killed Tarasoff. Turner M, Kennedy M. Tarasoff and the duty to warn third parties. Psychiatric Bull 1997; 21: 465-466;

The correct answer is: Confidentiality

Question 59
Not answered
Marked out of 1.00
Flag question

MoK Sociocultural Psychiatry 024
According to the Holmes and Rahe Social readjustment scale, which of the following has the highest 'Life Change' value?

Select one:
- Pregnancy
- Retirement
- Death of a spouse
- Marriage
- Minor violation of law

Your answer is incorrect.

The Holmes and Rahe stress scale is a list of 43 stressful life events that can contribute to mental/physical illness. When using Holmes and Rahe Stress Scale, the number of "Life Change Units" for relevant events in the past year of one's life are summed up to provide a final score of total stress exposure. The top one on the list is the death of a spouse followed by divorce.

The correct answer is: Death of a spouse

Question 60
Not answered
Marked out of 1.00
Flag question

MoK Sociocultural Psychiatry 027
Which of the following ethical theories concerns itself with the actions of an individual, rather than the thinking that lead up to it or the consequences?

Select one:
- Deontology
- Teleology
- Consequentialism
- Utilitarianism
- Virtue theory

Your answer is incorrect.

Deontological ethics derives the rightness or wrongness of one's conduct from the character of the behaviour itself rather than the outcomes of the conduct.

The correct answer is: Deontology

Question 61
Not answered
Marked out of 1.00
Flag question

MoK Sociocultural Psychiatry 031
A patient demands to have a CT scan of her head to evaluate her migrainous headaches. Your clinical judgment suggests that the pain is due to a migraine and there is no necessity for a CT scan.She does not agree and repeatedly asks for the scan. The best option in this situation is

Select one:
- Do not order the CT scan and ask her to seek medical care elsewhere
- Do not order the CT scan, but discharge her from your care
- Order the CT scan so that the patient will not lodge a complaint with hospital managers.
- Do not order the CT scan, but offer to refer her for a second opinion if she wishes to consider this.
- Ask her to arrange for a privately paid CT scan if she is willing to pay for this.

Your answer is incorrect.

Ordering unnecessary tests is both inappropriate utilization of health care resources and needlessly expensive. The CT scan should not be ordered for this patient if it is not indicated, regardless of their request.Provided the professional adheres to an existing level of standard of care to this patient, then a malpractice or negligence should not be of concern. There is not a clear reason to discharge the patient from the caseload completely and to do so may be construed as patient abandonment. The best choice, in this case, is to offer the patient an opportunity to get a second opinion on her symptoms.

The correct answer is: Do not order the CT scan, but offer to refer her for a second opinion if she wishes to consider this.

Question 62
Not answered
Marked out of 1.00
Flag question

MoK Sociocultural Psychiatry 032
In some families, contradictory parental demands on children lead to confusion and frustration, increasing the risk of emotional difficulties. Which of the following term best captures this situation?

Select one:
- Learned Helplessness
- Rapproachment
- Marital Schism
- Double bind
- Projective identification

Your answer is incorrect.

The essential hypothesis of the double bind theory is that the 'victim'-the person who becomes psychotically unwell-finds him or herself in a communicational matrix, in which messages contradict each other. These are described as injunctions as below: Primary Negative Injunction: 'I'll kill you if you stay.' i.e. 'Do not stay.' Secondary Negative Injunction: 'I am used to people leaving me.' i.e. 'Do not abandon me.' Tertiary Negative Injunction: 'That's right. It is so typical of you to irritate me and then leave me. No guts to face your problems. I'm prepared to stick here'. i.e. Don't leave.

The correct answer is: Double bind

Question 63
Not answered
Marked out of 1.00
Flag question

MoK Sociocultural Psychiatry 033
Which of the following is not a risk factor for ADHD?

Select one:
- Large size of the family
- Family history of ADHD
- Low birth weight
- Low educational attainment in parents
- Food additives

Your answer is incorrect.

There is no proven link between food additives and ADHD. ADHD and its common complications are associated with low social class, large family size, paternal antisocial personality, and maternal mental disorder.

The correct answer is: Food additives

Question 64
Not answered
Marked out of 1.00
Flag question

MoK Sociocultural Psychiatry 037
All of the following factors commonly influence the basic features of health explanatory belief models except

Select one:
- history of good physical health
- age
- time-point and context of assessment
- educational status
- gender

Your answer is incorrect.

Kleinman's work in the development of a health explanatory belief model has led to an increased appreciation of the role of a patients' understanding of the causes of an illness when attempting to treat the illness. Age, gender, cultural background, time and context of assessment and educational status can influence one's explanatory models. Ref: Kramer et al., West J Med. 2002 Sep; 176(4): 227-231.

The correct answer is: history of good physical health

Question 65
Not answered
Marked out of 1.00
Flag question

MoK HumanDevelopment 012
The average age of the earliest retrieved memory is

Select one:
- 2.5 years
- 3.5 years
- 5 years
- 4.5 years
- 2 years

Your answer is incorrect.

The average age of the earliest retrieved memory is 3.5 years. There is a total lack of memories for events occurring during the first few years of life, and there is amnesia for events that occurred in the first 2 to 5 years. This is also termed as infantile amnesia.

The correct answer is: 3.5 years

Question 66
Not answered
Marked out of 1.00
Flag question

MoK HumanDevelopment 013
Infants exhibit the capacity for colour vision and accommodation at the age of

Select one:
- 4 months
- 2 months
- 1 month
- 6 months
- Birth

Your answer is incorrect.

At birth, infants are able to track and scan objects, can discriminate levels of brightness, able to fix objects, fixed focus at about 20 cm, figure-ground discrimination and can have slightly impaired visual acuity. At 1month, they can differentiate faces with preference shown for complex stimuli. At 2 months- possesses depth perception, prefers 3 dimensional rather than 2 dimensional representations of a face. At 4 months, colour vision and accommodation develops. At 6 months accurate acuity is achieved(6:6).

The correct answer is: 4 months

Question 67
Not answered
Marked out of 1.00
Flag question

MoK HumanDevelopment 019
In a Strange Situation experiment, a 2-year-old girl continues to play without getting disturbed when her mother leaves the room and does not show any response when she comes back. What type of attachment is seen here?

Select one:
- Absence of attachment
- Disorganised attachment
- Resistant attachment
- Secure attachment
- Anxious avoidant attachment

Your answer is incorrect.

Anxious Avoidant attachment is characterized by anxiety and fright within the child and signs of not trusting the mother resulting in an indifferent attitude towards mother leaving the room or entering the room. Distress, when left alone, can be comforted by a stranger rather easily.

The correct answer is: Anxious avoidant attachment

Question 68
Not answered
Marked out of 1.00
Flag question

MoK HumanDevelopment 022
A boy can identify that two balls are of the same size even though they appear to be of different sizes due to a distance illusion. This corresponds to which stages of Paget's cognitive development?

Select one:
- Formal operational
- Operational stage
- Sensorimotor
- Pre-operational
- Concrete operational

Your answer is incorrect.

In concrete operational stage children start achieving 'conservation'. They are successful in conservation tasks involving numbers, quantity and volume by 7-8 years. The child realizes that certain properties like volume, number, length and mass of objects remain unchanged even when the objects appearances are altered.

The correct answer is: Concrete operational

Question 69
Not answered
Marked out of 5.00
Flag question

MoK Assessment EMI008
Interview techniques
For each of the following statements given below, identify the interview technique used.

You seem very upset when talking about your girlfriend. Are you having any problems with your relationship?	Choose...
The psychiatrist uses approval nods, leaning forwards slightly to express interest, using responses such as 'Yes. And then?' 'Yeah, go on..', 'Uh-huh', etc.	Choose...
You mentioned having taken overdoses since your last admission. How many times have you taken overdoses in total? Seven...? Eight...?	Choose...
What sort of recreational drugs do you usually use when you are drinking?	Choose...
The patient said 'It is not good if one's parents are divorced even before one goes to school.' The doctor replied 'I'd like to hear more about your parents, but first let me get a picture of what's happening to you of late.'	Choose...

Your answer is incorrect.

Explanation: Interpretation is clarifying certain associations or relationships that the patient may not see. It is a sophisticated technique and should be used only after the doctor has established some rapport. Should be stated as a hypothesis after sufficient collection of evidence from the interview.

Facilitation is helping patients continue the interview by providing both verbal and nonverbal encouragement. It helps patient to feel that the doctor is listening actively. Encourages flow of information.

Symptom exaggeration: When deception or minimisation is expected, overstating a guessed frequency in order to elicit a true answer. It is also helpful in reducing guilt to a certain extent as the patient feels that the doctor has expected a higher amount of problem that what she/he has brought.

Symptom expectation: Without a formal admission from the patient, asking about details of problem behaviour. The doctor assumes (rightly) that the patient is involved in the act.

Redirection is gently reorienting patient towards the topic of discussion. The motive is to keep the patient back on track.

Ref: Adapted from Kay J & Tasman A. Essentials of Psychiatry, 2nd edition, 2006. John Wiley & Sons, Ltd.

The correct answer is: You seem very upset when talking about your girlfriend. Are you having any problems with your relationship? – Interpretation, The psychiatrist uses approval nods, leaning forwards slightly to express interest, using responses such as 'Yes. And then?' 'Yeah, go on..', 'Uh-huh', etc. – Facilitation, You mentioned having taken overdoses since your last admission. How many times have you taken overdoses in total? Seven...? Eight...? – Symptom exaggeration, What sort of recreational drugs do you usually use when you are drinking? – Symptom expectation, The patient said 'It is not good if one's parents are divorced even before one goes to school.' The doctor replied 'I'd like to hear more about your parents, but first let me get a picture of what's happening to you of late.' – Redirection

Question 70

Not answered

Marked out of 3.00

Flag question

MoK Assessment EMI013

Maladaptive defences

For each of the disorders below choose the most prominent defence mechanisms used.

Fugue reactions (one) Choose...

Depression (one) Choose...

Pseudoseizures (one) Choose...

Your answer is incorrect.

Explanation: **Dissociation** is the predominant defence seen in dissociative fugue and amnesia. This involves temporarily but drastically modifying one's sense of personal identity or mental functions to avoid emotional distress.Moving back into childish or earlier developmental phase to avoid confronting a conflict is called as **regression**. Regression may explain the withdrawal and tearfulness seen in depression.**Conversion** refers to the switching of mental conflicts into somatic symptoms.

The correct answer is: Fugue reactions (one) – Dissociation, Depression (one) – Regression, Pseudoseizures (one) – Conversion

Question 71

Not answered

Marked out of 4.00

Flag question

MoK Assessment CXEMI014

Rating scales in clinical use

Identify the relevant rating scales for each of the following conditions

1. Choose TWO observer-rated tests to monitor depression in a woman recently diagnosed with postnatal depression (Choose Two)

2. Choose the most appropriate tool to screen for depression in mothers following childbirth

3. Choose one scale to assess EPSEs due to antipsychotic use.

Explanation:

Hamilton and Montgomery Asberg depression rating scales are observer-rated tests used to monitor depression.

The Edinburgh Postnatal Depression Scale (EPDS) have been used to screen for depression in women during the antepartum and postpartum periods. Edinburgh Postnatal Depression Scale was specially developed for assessing postpartum depression and relies much less on somatic questions. The questions on the Edinburgh scale (10 items, can be self or clinician-rated) are framed within the "past seven days", and the response format is frequency-based. Routine use of EPDS during the postpartum period has been shown to increase the detection of postnatal depression compared with usual care, but it is not sensitive to detect a change in the severity.

The Simpson-Angus Scale is helpful to assess EPSEs due to the use of antipsychotic medications. It is less sensitive than AIMS (Abnormal Involuntary Movement Scale) scale to detect tardive dyskinesia.

Screening for Depression Across the Lifespan: A Review .., http://www.aafp.org/afp/2002/0915/p1001.html (accessed March 31, 2015).

The correct answer is:
Choose TWO observer-rated tests to monitor depression in a woman recently diagnosed with postnatal depression (Choose Two) - Hamilton Depression scale, Montgomery Asberg Depression Rating Scale,

Choose the most appropriate tool to screen for depression in mothers following childbirth - Edinburgh postnatal depression scale,

Choose one scale to assess EPSEs due to antipsychotic use - Simpson-Angus Scale

Question 72

Not answered

Marked out of 4.00

Flag question

MoK Assessment EMI018

Varieties of Delusions

For each of the descriptions below choose the most closely associated psychopathological terms from the given list.

A 32-year-old psychotic patient with persistent auditory hallucinations is now developing a belief that a transmitter has been placed in his ear. Choose...

A 21-year-old psychotic patient remembers his parents taking him for an operation as a child becoming convinced that he had been implanted with monitoring devices, which have become active in his life. Choose...

A 29-year-old psychotic patient with an odd belief that certain individuals are not who they externally appear to be. Choose...

A 33-year-old psychotic patient is convinced that the Mafia is after him and are conspiring to ruin his life Choose...

Your answer is incorrect.

Explanation:

Secondary delusions that arise when a patient attempts to find an explanation for primary psychopathological processes (in case 1, a hallucination) are called delusional elaborations. In the above scenario, a patient with auditory hallucinations is now developing a belief that a transmitter has been placed in his ear.

Delusional memory is a primary delusion, which is recalled as arising as a result of a memory. In the above scenario, a psychotic patient remembers his parents taking him for an operation as a child (memory). However, in a delusional manner he is now convinced that control and monitoring devices were implanted. Delusional memory can also be a retrospective delusion where something that never happened and so false, irrational or bizarre is reported as if it has occurred in the past.

Delusional misidentification is a delusional belief that certain individuals are not who they externally appear to be. It includes Capgras and Fregoli syndrome).

A persecutory delusion is a belief that one's life is being interfered with in a harmful way.

(Ref: Oxford Handbook of Psychiatry 1e-Pg 84-87)

The correct answer is: A 32-year-old psychotic patient with persistent auditory hallucinations is now developing a belief that a transmitter has been placed in his ear. – Delusional elaboration, A 21-year-old psychotic patient remembers his parents taking him for an operation as a child becoming convinced that he had been implanted with monitoring devices, which have become active in his life. – Delusional memory, A 29-year-old psychotic patient with an odd belief that certain individuals are not who they externally appear to be. – Delusional misidentification, A 33-year-old psychotic patient is convinced that the Mafia is after him and are conspiring to ruin his life – Persecutory delusions

Question 73
Not answered
Marked out of 5.00
Flag question

MoK Assessment EMI021
Disorders of perception
Which of the listed descriptive psychopathological terms refers to the following symptoms?

A 34-year-old gentleman in the psychiatric outpatient clinic mentioned 'I heard a voice calling my name which then startled me back to wakefulness to find no one there.'	Choose...
A 20-year-old university student mentioned that while she was walking across a lonely park at night; she became anxious on mistaking bushes rustling in the wind as a hidden attacker but later calmed down	Choose...
A woman with bilateral cataracts says that she sees a number of her dead relatives walking in the house	Choose...
A 46-year-old woman on the psychiatric ward says 'All my blood has dropped into my legs, and I can feel a water level in my chest.'	Choose...
A 37-year-old gentleman on the psychiatric ward described 'I thought my life was outside my feet and made them vibrate.'	Choose...

Your answer is incorrect.

Explanation:

Hypnagogic hallucinations: A transient false perception experienced while on the verge of falling asleep. (Go for gogic) It is usually auditory. The same phenomenon if experienced when waking up is called hypnopompic hallucinations. Both can occur in healthy individuals.

In affect illusion, there is a mixture of heightened emotions and misperception.

Charles Bonnet Syndrome: Elderly patients, with normal consciousness and no brain pathology, with reduced visual acuity due to ocular problems, experience vivid, distinct, usually coloured hallucinations. The images are mostly humans; at times animated cartoons are reported. The vividness of this experience is a striking contrast to the real sensory percepts which appear blurred due to the visual impairment. Insight about unreality is usually preserved - though they may evoke emotions including fear and joy.

Case 4 is a description of a hygric hallucination (fluid-wetness), which is a type of superficial somatic hallucination.

Kinaesthetic or proprioceptive hallucinations is also a form of somatic hallucination. It refers to joint or muscle sense, often linked to bizarre somatic delusions. They are also seen in benzodiazepine withdrawal and alcohol intoxication

The correct answer is: A 34-year-old gentleman in the psychiatric outpatient clinic mentioned 'I heard a voice calling my name which then startled me back to wakefulness to find no one there.' – Hypnagogic hallucination, A 20-year-old university student mentioned that while she was walking across a lonely park at night; she became anxious on mistaking bushes rustling in the wind as a hidden attacker but later calmed down – Affect illusions, A woman with bilateral cataracts says that she sees a number of her dead relatives walking in the house – Charles Bonnet syndrome, A 46-year-old woman on the psychiatric ward says 'All my blood has dropped into my legs, and I can feel a water level in my chest.' – Hygric hallucinations, A 37-year-old gentleman on the psychiatric ward described 'I thought my life was outside my feet and made them vibrate.' – Kinaesthetic hallucination

Question 74
Not answered
Marked out of 5.00
Flag question

MoK Assessment EMI023
Speech disorder
Identify the correct terms that represent each of the following speech disorders

Decrease in spontaneous talking occurring in depression	Choose...
Increased quantity of speech seen in early manic states	Choose...
The patient is mute but remain aware of the environment though cannot move or respond	Choose...
Difficulty in finding the correct name for an object	Choose...
Often seen in children who refuse to speak to certain people	Choose...

Your answer is incorrect.

Explanation:

Alogia is a term used to denote poverty of speech and a decrease in spontaneous talking; it occurs in depression and schizophrenia.

Logorrhoea is increased quantity of speech, without pressure of speech or formal thought disturbances and seen especially in early manic states.

Nominal aphasia is also called as amnestic aphasia as it manifests as a difficulty in finding the correct name for an object.

Akinetic mutism is associated with lesions of the upper midbrain or posterior diencephalon. Here the patient is mute but remains aware of the environment though cannot move or respond to commands.

Elective mutism: Mostly seen in children who refuse to speak to certain people; for example, the child may not speak at school but may speak at home.

The correct answer is: Decrease in spontaneous talking occurring in depression – Alogia, Increased quantity of speech seen in early manic states – Logorrhoea, The patient is mute but remain aware of the environment though cannot move or respond – Akinetic mutism, Difficulty in finding the correct name for an object – Nominal aphasia, Often seen in children who refuse to speak to certain people – Elective mutism

Question 75
Not answered
Marked out of 6.00
Flag question

MoK Psychology CXEMI 005
Cloninger's Theory of Personality
Identify the correct behavioural traits to match the dimensions of temperament described by Cloninger.

1. Harm avoidance traits (Choose TWO)		
2. Novelty seeking traits (Choose TWO)		
3. Reward dependence traits (Choose TWO)		

Explanation: Cloninger's psychobiological model of personality includes four dimensions of temperament (each 50 to 60 % heritable), which manifest early in life and 3 components of character, which are shaped by environment. The temperamental dimensions include

- (1) Novelty-seeking (includes frustration avoidance, impulsive decision-making)
- (2) Harm-avoidance (pessimistic worry about the future, passive avoidant behaviour, fear of uncertainty);
- (3) Reward-dependence (sentimentality, social attachment, and dependence on praise and approval)
- (4) Persistence (high perseverance and tolerance of frustration). The character dimensions are self-directedness, cooperativeness, and self-transcendence.

The correct answer is:
Harm avoidance traits (Choose TWO)- Fear of uncertainty, Pessimism,

Novelty seeking traits (Choose TWO)- Frustration Avoidance, Impulsivity,

Reward dependence traits (Choose TWO)- Praise dependence, Sentimentality

Finish review