What style of parenting is characterised by children following rules in response to their parents' arbitrary use of hostile control or harsh punishment?

Select one:

- Neglectful/uninvolved
- Permissive/indulgent
- Authoritarian ✔
- Authoritative
- Overbearing

Your answer is correct.

The authoritarian parenting style is characterised as low in responsiveness but high in demandingness. Parents of this style tend to use hostile control or harsh punishment in an arbitrary way to gain compliance, but they seldom provide explanation or allow verbal give-and-take.

Ref: Bi X et al. Parenting styles and parent–adolescent relationships: The mediating roles of behavioral autonomy and parental authority. Frontiers in psychology. 2018 Nov 13;9:2187.

The correct answer is: Authoritarian

Question 2

Correct

Mark 1.00 out of 1.00

Flag question

In attachment theory (according to Bowlby), what is the sequence of the stages of separation anxiety?

Select one:

- Despair, protest, detachment
- Despair, detachment, protest
- Protest, detachment, despair
- Protest, despair, detachment ✔
- Detachment, protest, despair

Your answer is correct.

Underlying attachment theory were numerous observations of separations and reunions between young parents and children. The process has been described as occurring in phases; called protest, despair, and denial by Robertson; and protest, despair, and detachment by Bowlby. Bowlby, among others, preferred the term detachment because it was a natural counterpart to attachment. When separation anxiety is activated, the most prominent sign of the first phase – protest – is the child's acute distress. If the parent fails to return for long, the behaviour will be replaced by expressions of the next phase – despair. If the separation continues, the despair is gradually replaced by detachment.

Ref: Klette T, Killén K. Painful transitions: a study of 1-year-old toddlers' reactions to separation and reunion with their mothers after 1 month in childcare. Early Child Development and Care. 2018 Jan 11.

The correct answer is: Protest, despair, detachment

Question 3

Correct

Mark 1.00 out of 1.00

Flag question

According to Freud, when conflict is not worked through to a realistic solution, drives or wishes are expelled from consciousness through which of the following?

Select one:

- Depression
- Repression ✔
- Altruism
- Sublimation
- Regression

Your answer is correct.

The classic Freudian view of the genesis of neuroses regards conflict as essential. The conflict can arise between instinctual drives and external reality or internal agencies, such as the id and the superego or the id and the ego. Moreover, because the conflict has not been worked through to a realistic solution, the drives or wishes that seek discharge are expelled from consciousness through repression. Repression consists of the expelling and withholding from consciousness awareness of an idea or feeling.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. pp. 1067-1068..

The correct answer is: Repression

Question **4**

Incorrect

Mark 0.00 out of 1.00

Flag question

Which recombination fraction value denotes that two genes are likely to be independently assorted?

Select one:

- 0
- 0.25
- 1 ✖
- 0.75
- 0.5

Your answer is incorrect.

The evidence for linkage between two loci depends on the recombination frequency between them. Recombination frequency is measured by the recombination fraction, equal to the genetic distance between the two loci. 1% recombination equals 1 centimorgan (cM) in the genetic distance and, on average, covers a physical distance of about 1 megabase of DNA. Genes are said to independently assort, when they are apart at a distance of 50 cM or more. Thus, a recombination fraction of 0.5 or 50% indicates that two loci are not linked but rather that they are segregating independently.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 988.

The correct answer is: 0.5

Question 5

Incorrect

Mark 0.00 out of 1.00

Flag question


Deep brain stimulation in OCD targets which of the following structures?

Select one:

- Anterior limb of internal capsule
- Dorsolateral prefrontal cortex
- Subgenual cingulate gyrus ✖
- Orbitofrontal cortex
- Cerebellopontine angle

Your answer is incorrect.


Deep brain stimulation is best regarded as an experimental treatment for OCD and depression. DBS involves implantation of bilateral electrodes under stereotactic guidance and MRI confirmation. Targets for DBS in OCD include the anterior limb of the internal capsule and, for depression, the subgenual cingulate gyrus.



Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 313.

The correct answer is: Anterior limb of internal capsule

Which pattern of inheritance is shown in this pedigree?



Select one:

- Autosomal recessive ✖
- Spontaneous (no inheritance)
- X-linked recessive
- X-linked dominant
- Autosomal dominant

Your answer is incorrect.

Autosomal dominant (AD) disorders are those in which a patient manifests clinical symptoms when only a single copy of the mutant gene is present. Inheritance of AD disorders follows several general principles: 1) Each affected person has an affected parent; 2) Affected persons, on average, have equal numbers of affected and unaffected children; 3) Normal children of affected parents only have unaffected children; 4) Males and females are affected in equal proportions; 5) Each sex is equally likely to transmit the disorder to male and female children; and 6) Vertical transmission of the disorder occurs through successive generations.

Ref: Levine F. 1 - Basic Genetic Principles. Editor(s): Polin, RA et al. Fetal and Neonatal Physiology (5th Edition), 2017. p. 1-13.

The correct answer is: Autosomal dominant

Question **7**

Incorrect

Mark 0.00 out of 1.00

Flag question

Which ion channel is most implicated in schizophrenia?

Select one:

- Sodium ✖
- Potassium
- Chloride
- Calcium
- Chlorine

Your answer is incorrect.

Genetic association studies of schizophrenia have consistently demonstrated a convergence of genetic risk upon sets of genes with synaptic functions, including discrete signal transduction pathways such as voltage-gated calcium channel complexes and glutamate receptor complexes. The relationship between the expression of genes involved in voltage-gated cation channel activity and association with schizophrenia has been shown to be strongest for the subset of genes related to calcium channel activity.

Ref: Clifton NE et al. Developmental profile of psychiatric risk associated with voltage-gated cation channel activity. Biological psychiatry. 2021 Sep 15;90(6):399-408.

The correct answer is: Calcium

Question 8

Correct

Mark 1.00 out of 1.00

Flag question

Southern blots are used to identify which of the following?

Select one:

- Post translational modification
- DNA fragment sequence ✔
- DNA binding protein
- Protein sequence
- RNA sequence

Your answer is correct.

Developed by and named after Edward Southern, a Southern blot is used to identify if a known DNA sequence is present in a sample by using a specific probe to identify the sequence in a large mixture of other DNA fragments. Southern blots were traditionally used to validate the successful introduction or removal of DNA sequences in the genome of genetically modified animals. Now, modern PCR techniques are more sensitive and a faster alternative to identifying a DNA fragment.

Ref: Carter M, Essner R, Goldstein N, Iyer M. Guide to Research Techniques in Neuroscience, 3rd Edition. 2022. p. 239.

The correct answer is: DNA fragment sequence

Question 9

Incorrect

Mark 0.00 out of 1.00

Flag question

Adverse Lyonization is seen in which of the following conditions?

Select one:

- Fragile X syndrome
- Huntington disease
- Wilson disease
- Klinefelter syndrome
- Down syndrome ✖

Your answer is incorrect.

Fragile X syndrome (FXS)-affected individuals display a spectrum of neurologic, psychiatric and developmental issues which often manifest with less severity in full-mutation (FM) females as a result of X-chromosome inactivation or skewed or adverse Lyonization. Random inactivation (or Lyonisation) of either the paternal or the maternal X chromosomes occurs in each cell in female embryos, resulting in a mosaic expression of X-linked genes in females with 50:50 expression of maternal and paternal alleles. However, in the presence of a structural abnormality such as fragile X chromosome, a deviation from the 50:50 ratio, known as skewed or adverse Lyonization, may occur wherein the inactivation of the unaffected X chromosome is favored. This leads to the clinical expression of FXS in some women (30-50% of the Fragile X chromosome carriers). In general, the X-chromosome inactivation ratio is considered skewed if it is ≥65:35.

Ref: Brand BA, Blesson AE, Smith-Hicks CL. The impact of X-chromosome inactivation on phenotypic expression of X-linked neurodevelopmental disorders. Brain Sciences. 2021 Jul 9;11(7):904.

The correct answer is: Fragile X syndrome

Question **10**

Correct

Mark 1.00 out of 1.00

Flag question

The dentate gyrus is a part of which structure?

Select one:

- Pons
- Cerebellum
- Hippocampus ✔
- Midbrain
- Medulla

Your answer is correct.

The dentate gyrus is a subregion of the hippocampus. Encoding spatial information as well as pattern separation are seen as its major functions, among many other roles relevant to human memory.

Ref: Borzello M et al. Assessments of dentate gyrus function: discoveries and debates. Nature Reviews Neuroscience. 2023 Jun 14:1-6.

The correct answer is: Hippocampus

Question **11**

Incorrect

Mark 0.00 out of 1.00

Flag question

Which of the following inflammatory biomarkers is commonly elevated in depression?

Select one:

- IL-13
- IL-4 ✖
- IL-22
- IL-6
- IL-11

Your answer is incorrect.

Biomarkers of inflammation in patients with depression include elevated blood levels of Il-1beta, IL-6, TNF, and CRP. Of these, prospective associations of depression and inflammatory biomarkers are observed in both directions, particularly for IL-6.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 257.
Giollabhui MN, Ng TH, Ellman LM, Alloy LB. The longitudinal associations of inflammatory biomarkers and depression revisited: systematic review, meta-analysis, and meta-regression. Molecular Psychiatry. 2021 Jul;26(7):3302-14.

The correct answer is: IL-6

Question 12

Correct

Mark 1.00 out of 1.00

Flag question

The locus coeruleus is involved in which of the following pathways?

Select one:

- Serotonergic
- Dopaminergic
- Noradrenergic ✔
- Cholinergic
- GABA-ergic

Your answer is correct.

CNS norepinephrine (noradrenaline) synthesis takes place primarily in the locus coeruleus, which is located in the dorsal portion of the pons. While the locus coeruleus gives rise to norepinephrine tracts, the raphe nucleus gives rise to serotonin tracts.

Ref: Kaufman DM, Geyer HL, Milstein MJ, Rosengard JL. Kaufman's Clinical Neurology for Psychiatrists, 9th Edition. 2023. p. 526.

The correct answer is: Noradrenergic

Question 13

Incorrect

Mark 0.00 out of 1.00

Flag question

Fluorodeoxyglucose (FDG) is used in which of the following neuroimaging techniques?

Select one:

- PET
- CT
- MRI
- MRS ✖
- fMRI

Your answer is incorrect.

PET relies on positron-emitting, biologically active radioisotopes (radioligands) incorporated into organic molecules. The radioligands, inhaled or injected intravenously, undergo metabolism in the brain and emit positrons. The reaction between positrons and electrons yields photons that PET detects and transforms into images. Most PET studies measure the metabolism of a substitute for glucose, fluorine-18-labeled fluorodeoxyglucose (FDG).

Ref: Kaufman DM, Geyer HL, Milstein MJ, Rosengard JL. Kaufman's Clinical Neurology for Psychiatrists, 9th Edition. 2023. pp. 507.

The correct answer is: PET

Question **14**

Incorrect

Mark 0.00 out of 1.00

Flag question

In vivo measurement of neuroreceptors is possible by which of the following techniques?

Select one:

- fMRI
- PET
- MEG
- MRS ✖
- CT

Your answer is incorrect.

Dynamic positron emission tomography (PET) with appropriate neuroreceptor ligands permits the measurement of neurotransmitter release in the human brain with appropriate pharmacological or behavioural challenge. This imaging technique has been used for more than two decades to probe the neurochemical underpinnings of cognitive and motivational functions in normal and neuropathological conditions. PET is the only technology to date that has the capability to reveal the metabolism of molecules, receptors, and neurotransmitters in vivo with the appropriate radioligands and precursor.

Ref: Petibon Y et al. PET imaging of neurotransmission using direct parametric reconstruction. NeuroImage. 2020 Nov 1;221:117154.
Su Y et al. Real-time in vivo detection techniques for neurotransmitters: A review. Analyst. 2020;145(19):6193-210.

The correct answer is: PET

Question 15

Correct

Mark 1.00 out of 1.00

Flag question

In ICD-11, disorders of Intellectual development are characterised by intellectual functioning and adaptive behaviour that are at least how many standard deviations below the mean?

Select one:

- 3
- 1
- 2 ✔
- 4
- 5

Your answer is correct.

In ICD-11, disorders of Intellectual development are characterised by intellectual functioning and adaptive behaviour that are approximately two or more standard deviations below the mean (approximately less than the 2.3rd percentile), based on appropriately normed, individually administered standardised tests.

Ref: International Statistical Classification of Diseases and Related Health Problems (11th ed. ICD-11; World Health Organization, 2020)

The correct answer is: 2

Question **16**

Correct

Mark 1.00 out of 1.00

Flag question

At what stage in Conrad's model of the development of delusional psychosis does the individual start to find new meaning for psychological events?

Select one:

- Anastrophy
- Trema
- Apophany ✔
- Consolidation
- Residuum

Your answer is correct.

Conrad proposed five stages in the development of delusional psychosis:
1) Trema: delusional mood representing a total change in perception of the world
2) Apophany: a search for, and the finding of, new meaning for psychological events
3) Anastrophy: heightening of the psychosis
4) Consolidation: forming of a new world or psychological set based on new meanings<b5)
Residuum: eventual autistic state

Ref: Oyebode F. Sims' symptoms in the mind: Textbook of descriptive psychopathology. 7th edition, 2022. p. 119.

The correct answer is: Apophany

Question 17

Correct

Mark 1.00 out of 1.00

Flag question

A patient with mania demonstrates the following speech: "Proctors, doctors. Doctors are in the hospital. Apples keep doctors away. Newton had an apple. Newton discovered gravity. Graves are in the ground. Ground, found." What is this speech known as?

Select one:

- Perseveration
- Derailment
- Flight of ideas ✔
- Circumstantiality
- Vorbeireden

Your answer is correct.

In flight of ideas, thoughts follow each other rapidly; there is no general direction of the thinking; and the connections between successive thoughts appear to be due to chance factors which, however, can usually be understood. The absence of a determining tendency to thinking allows the associations of the train of thought to be determined by chance relationships, verbal associations of all kinds, clang associations, proverbs, maxims, and cliches. The chance linkage of thoughts in flight of ideas is demonstrated by the fact that one could completely reverse the sequence of the record of a flight of ideas, and the progression of thought would be understood just as well (or just as poorly).

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 44.

The correct answer is: Flight of ideas

Question **18**

Correct

Mark 1.00 out of 1.00

Flag question

A middle-aged man with OCD believes that his thoughts and actions can cause specific outcomes in ways that defy the normal laws of cause and effect. The most appropriate term for this defence mechanism is

Select one:

- Reaction formation
- Magical thinking ✔
- Displacement
- Rationalisation
- Acting out

Your answer is correct.

Magical thinking refers to a belief in personal power to control or cause external events in the real world beyond culturally and rationally accepted laws of causality. It is found when one attributes magical properties to thoughts or behaviours (e.g. 'if I throw this piece of paper in the bin five times in a row, then I'll pass my exams'). This defence is associated with OCD.

Ref: Fite RE, Adut SL, Magee JC. Do you believe in magical thinking? Examining magical thinking as a mediator between obsessive-compulsive belief domains and symptoms. Behavioural and Cognitive Psychotherapy. 2020 Jul;48(4):454-62. Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 893.

The correct answer is: Magical thinking

Question 19

Correct

Mark 1.00 out of 1.00

Flag question

Mild personality disorder in ICD-11 is equivalent to what level of personality functioning in DSM-5?

Select one:

- Extreme impairment
- Moderate impairment ✔
- No impairment
- Some impairment
- Severe impairment

Your answer is correct.

The threshold for a personality disorder diagnosis is at least Mild personality disorder (i.e., some impairment as per ICD-11) or Moderate impairment of personality functioning (DSM-5).

Ref: Bach B, First MB. Application of the ICD-11 classification of personality disorders. BMC psychiatry. 2018 Dec;18:1-4.

The correct answer is: Moderate impairment

Question **20**

Correct

Mark 1.00 out of 1.00

Flag question

In ICD-11, which of the following can differentiate a manic episode from a hypomanic episode?

Select one:

- Decreased need for sleep
- Psychotic symptoms ✔
- Reckless behaviour
- Racing thoughts
- Rapid speech

Your answer is correct.

In ICD-11, a hypomanic episode is accompanied by characteristic symptoms such as rapid speech, rapid/racing thoughts, increased self-esteem, an increase in sexual drive or sociability, decreased need for sleep, distractibility, and/or impulsive or reckless behaviour. In a hypomanic episode, there are no accompanying delusions or hallucinations.

Ref: International Statistical Classification of Diseases and Related Health Problems (11th ed. ICD-11; World Health Organization, 2020)

The correct answer is: Psychotic symptoms

Question 21

Incorrect

Mark 0.00 out of 1.00

Flag question

In ICD-11, which of the following requires the longest duration of symptoms for diagnosis?

Select one:

- Adjustment disorder
- Schizoaffective disorder
- Schizotypal disorder
- Schizophrenia
- Delusional disorder ✖

Your answer is incorrect.

For ICD-11 diagnosis of schizotypal disorder, symptoms should have been present, continuously or episodically, for at least 2 years. Symptoms of schizoaffective disorder must have persisted for at least one month. ICD-11 notes that the duration of adjustment disorder varies widely; symptoms that resolve within a few days do not typically warrant a diagnosis of adjustment disorder, but ICD-11 does not designate a specific duration of symptoms. Symptoms must have persisted for at least one month in order for a diagnosis of schizophrenia to be assigned. In ICD-11, delusional disorder is characterised by the development of a delusion or set of related delusions, typically persisting for at least 3 months and often much longer.

The correct answer is: Schizotypal disorder

Question 22

Incorrect

Mark 0.00 out of 1.00

Flag question

According to ICD-11, which of the following is most likely to differentiate intermittent explosive disorder from oppositional defiant disorder?

Select one:

- Defiant behaviours
- Outbursts out of proportion to the provocation ✖
- Vindictive behaviours
- Headstrong behaviours
- Significant physical aggression

Your answer is incorrect.

According to ICD-11, individuals with intermittent explosive disorder (IED) are more likely to exhibit significant physical aggression than those with oppositional defiant disorder (ODD). Severe temper outbursts that are grossly out of proportion to the provocation are core to IED, but also occur in the context of ODD with chronic irritability-anger. Individuals with ODD with chronic irritability-anger typically display other features of ODD - including defiant, headstrong, or vindictive behaviours - which are not characteristic of IED.

Ref: International Statistical Classification of Diseases and Related Health Problems (11th ed. ICD-11; World Health Organization, 2020)

The correct answer is: Significant physical aggression

Question 23

Correct

Mark 1.00 out of 1.00

Flag question

A woman sees colours when she hears her husband's voice or laughter. This is known as what?

Select one:

- Dysmegalopsia
- Pseudologia fantastica
- Synaesthesia ✔
- Autoscopy
- Vorbeireden

Your answer is correct.

Synaesthesia is the experience of a stimulus in one sense modality producing a sensory experience in another (e.g. 'tasting' sounds or 'hearing' colours).

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 36.
Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 118.

The correct answer is: Synaesthesia

Question **24**

Incorrect

Mark 0.00 out of 1.00

Flag question

Which of the following is true about stupor?

Select one:

- The patient appears awake and alert
- The patient is able to move
- It exists on a spectrum of wakefulness to coma
- Consciousness is severely impaired ✖
- It is the same as coma

Your answer is incorrect.

Note that in neurology the term stupor is used to refer to any state of reduced consciousness. In psychiatry, stupor is the absence of movement and communicative behaviour (akinesia + mutism) where there is no impairment of consciousness. It can be regarded as an extreme form of hypokinesia. Functional stupor occurs in a variety of psychiatric illnesses wherein patients are mute and paralyzed but generally appear alert with preserved intellectual function.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 118.
Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 98.

The correct answer is: The patient appears awake and alert

Question 25

Correct

Mark 1.00 out of 1.00

Flag question

Which of the following is true regarding alcohol use by young people in the UK?

Select one:

- Those who drink alcohol regularly from an early age are more likely to develop later alcohol misuse/abuse ✔
- Young people in the UK aged 15-17 years are unlikely to binge drink
- Alcohol seldom has negative impacts on young people's educational performance
- Binge drinking in young people has no association with self-harm and thoughts of suicide
- The Health Behaviour in School-Aged Children Survey (HSBC) provides data on cannabis, but not alcohol

Your answer is correct.

Young people between the ages of 15 and 17 years are more likely to binge drink, which is associated with other health risk behaviours such as unprotected or regretted sexual activity; antisocial and criminal behaviour; and self-harm and thoughts of suicide. Those who drink regularly from an early age are more likely to develop later alcohol misuse or abuse. Alcohol use can negatively impact upon young people's educational performance, their friendships, their relationships with their parents, and the dynamics of their peer group. The Health Behaviour in School-Aged Children Survey (HBSC), as used in the UK, provides information on alcohol and substance use, including cannabis.

Ref:
https://stateofchildhealth.rcpch.ac.uk/evidence/hea
behaviours/alcohol-drug-use-young-people/

The correct answer is: Those who drink alcohol regularly from an early age are more likely to develop later alcohol misuse/abuse

Question **26**

Incorrect

Mark 0.00 out of 1.00

Flag question

Which of the following is a first rank symptom?

Select one:

- Upon hearing a clock strike 12, a man believes that he is the Messiah
- A man believes that his neighbours are poisoning him
- After his wife dies, a man briefly thinks that he sees her in the shadows
- A man hears voices commanding him to hurt other people ✖
- A man feels unsettled and believes that something is about to happen

Your answer is incorrect.

A delusional perception is a primary delusion of any content that is reported by the patient as having arisen following the experience of a normal perception. If the patient is asked to recall the point when they became aware of the delusion and its significance to them, they will report that the delusional perception arose on seeing a normal percept (the percept is a real external object, not a hallucinatory experience). Delusional perception is a first-rank symptom of schizophrenia.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 78, 106-107.

The correct answer is: Upon hearing a clock strike 12, a man believes that he is the Messiah

Question **27**

Correct

Mark 1.00 out of 1.00

Flag question

The cocktail party phenomenon involves

Select one:

- Cognitive misattribution
- Involuntary attention
- Visual perception
- Gustatory perception
- Selective auditory attention ✔

Your answer is correct.

The cocktail party phenomenon involves selective attention in speech perception that enables a listener to attend to one among several equally loud conversations occurring simultaneously, factors such as voice quality and directional cues facilitating the task. Recordings of conversations under similar circumstance are very much harder to understand. The term was introduced in 1957 by the British telecommunications engineer Colin Cherry.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015. p. 140.

The correct answer is: Selective auditory attention

Question **28**

Correct

Mark 1.00 out of 1.00

Flag question

What is the name given to the phenomenon in which individuals are reluctant to help in an emergency when there are many people present?

Select one:

- Domino effect
- Bystander effect ✔
- Halo effect
- Hawthorne effect
- Pygmalion effect

Your answer is correct.

The 'bystander effect' (also known as the 'Genovese syndrome') is the reluctance of bystanders to intervene in an emergency, especially when a person appears to be in distress, or a crime is being committed. Scores of experiments have shown that people are much less likely to intervene in an emergency, and are generally slower to respond, when other people are present than when they are alone, and this phenomenon is sometimes called 'group inhibition of helping' or 'bystander apathy effect'.

Ref: Marsh HW, Reeve J, Guo J, Pekrun R, Parada RH, Parker PD, Basarkod G, Craven R, Jang HR, Dicke T, Ciarrochi J. Overcoming limitations in peer-victimization research that impede successful intervention: Challenges and new directions. Perspectives on psychological science. 2023 Jul;18(4):812-28.

The correct answer is: Bystander effect

Question **29**

Incorrect

Mark 0.00 out of 1.00

Flag question

Which of the following is true of bereavement and grief in children?

Select one:

- Grief in children closely resembles that in adults ✖
- 1% of school-aged children/young people in the UK experience family bereavement
- Grief reactions are influenced by developmental levels and concepts of death
- Grief in children is an illness
- A child's grief usually resolves completely within 3 months following bereavement

Your answer is incorrect.

Grief reactions in children are influenced by developmental levels and concepts of death and may not resemble adult reactions. Family bereavement has continuous, cumulative effects on children's emotional and social well-being, long after bereavement happens. Grief is not an illness, but it does increase the risk of mental health difficulties, both in childhood and later life. Around 1/3 of bereaved children reach clinical levels of emotional/behavioural difficulties in the two years following a parent's death. Around 1 in 29 school-age children and young people (5-16) in the UK are bereaved at some point in their childhood of a parent or sibling.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 852.
https://committees.parliament.uk/writtenevidence/

The correct answer is: Grief reactions are influenced by developmental levels and concepts of death

Question **30**

Correct

Mark 1.00 out of 1.00

Flag question

Sherif and colleagues' (1954) field experiment observing group behaviours that involved two groups of eleven-year-old boys was focused on which of the following?

Select one:

- Effects of wakefulness and fatigue
- Non-verbal communication
- Group conflict and cooperation ✔
- Peer pressure and bullying
- Effects of persuasive speaking

Your answer is correct.

Realistic group conflict theory is the notion that group conflict and hostility are caused chiefly by competition for resources and power, as proposed by psychologist Muzafer Sherif and several colleagues in the Robbers Cave experiment. The field experiment involved two groups of white, middle-class eleven-year-old boys, and was named after the Robbers Cave State Park where the experiment took place.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015. pp. 638, 662.

The correct answer is: Group conflict and cooperation

Question **31**

Correct

Mark 1.00 out of 1.00

Flag question

A policeman comes to the scene of a crime, sees a well-dressed man, and decides he cannot be the culprit. What is the name given to this phenomenon?

Select one:

- Bystander effect
- Hawthorne effect
- Halo effect ✔
- Pygmalion effect
- Barnum effect

Your answer is correct.

The halo effect is a generalisation from the perception of one prominent or salient characteristic, trait, or personality attribute to an impression of the whole, leading to inflated correlations between rated items.

Ref: Batres C, Shiramizu V. Examining the "attractiveness halo effect" across cultures. Current Psychology. 2023 Oct;42(29):25515-9.

The correct answer is: Halo effect

Question **32**

Correct

Mark 1.00 out of 1.00

Flag question

Healthcare workers wash their hands more often following the implementation of a new protocol which involves the workers being observed. Which of the following explains their improvement in hand washing?

Select one:

- Pygmalion effect
- Bystander effect
- Barnum effect
- Hawthorne effect ✔
- Halo effect

Your answer is correct.

The Hawthorne effect refers to improvement in the performance or productivity of workers or students resulting from the introduction of new working methods or conditions, irrespective of the nature of the changes, the effect often being attributed to the feeling of being under concerned observation.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015. p. 332. Cohen AL. Hawthorne Effect. In Problems and Pitfalls in Medical Literature: A Practical Guide for Clinicians 2023 Sep 28 (pp. 33-35). Cham: Springer International Publishing.

The correct answer is: Hawthorne effect

Question 33

Correct

Mark 1.00 out of 1.00

Flag question

Which of the following is true of the concept of 'storm and stress' in adolescence?

Select one:

- Negative behaviours in adolescence are normative ✔
- Parent-adolescent relationships are unaffected by this phenomenon
- Percentages of adolescents displaying specific negative behaviours are specified
- Adolescents exhibit less negative externalising behaviours than children
- Positive behaviours are more common than negative behaviours in adolescence

Your answer is correct.

A primary aspect of the 'storm and stress' characterisation of adolescence is that adolescents exhibit more negative, or undesirable, characteristics in the domains of externalising, internalising, and parent-adolescent relationships than do younger and older individuals. The original characterisation by Hall (in 1904) who spoke of behaviour such as 'cruelty, laziness, lying, and thievery' as normative, is now seen as inaccurate and does not specify percentages of adolescents who are expected to display specific negative behaviours. Nevertheless, this characterisation has influenced the public opinion and is often referred to in the context of risk behaviour, mood disruption, and parent-child conflicts.

Ref: Buchanan CM et al. Adolescent Storm and Stress: A 21st Century Evaluation. Frontiers in Psychology. 2023 Aug 3;14:1257641.

The correct answer is: Negative behaviours in adolescence are normative

Question **34**

Incorrect

Mark 0.00 out of 1.00

Flag question

According to Bandura's classic experiments, which of the following is true about effectively reducing aggressive behaviour in children?

Select one:

- Observing an adult aggressor receive punishment reduces the probability of aggressive behaviour in children
- Not reinforcing an adult's aggressive behaviour leads to decreased aggression in children ✖
- Children who watch a film showing aggressive cartoons tend not to imitate this behaviour
- Positive reinforcement of an adult's aggressive behaviour has no effect on aggression in children
- Putting children into a new, different context eliminates the possibility of aggressive behaviour

Your answer is incorrect.

Key findings of Bandura's experiments on aggressive in children include the following: 1) Observation of an aggressive model is sufficient to elicit aggressive behaviour in the young child. 2) If the adult model is punished for their aggressive behaviour, the probability that the child will show aggressive behaviour is reduced. 3) Positive reinforcement or no reinforcement of the adult model's aggression leads to increased aggression on the part of the child (vicarious/indirect learning). 4) Children who watch a film showing aggressive people or cartoons tend to imitate this behaviour. 5) Children transfer, by means of generalisation, aggression into new, different contexts, even when the aggressive model is no longer present (delayed imitation).

Ref: Galanaki E, Malafantis KD. Albert Bandura's experiments on aggression modelling in children: A psychoanalytic critique. Frontiers in psychology. 2022 Nov 25;13:7360.

The correct answer is: Observing an adult aggressor receive punishment reduces the probability of aggressive behaviour in children

Question 35

Incorrect

Mark 0.00 out of 1.00

Flag question

Which of the following reflects the constructivist theory of perception?

Select one:

- Affordance is what the object offers the viewer in terms of potential action
- Perception does not require knowledge-based processes
- Depth and form are obtained directly from a visual image
- Perception involves making inferences about past experiences
- Gestalts include proximity features ✖

Your answer is incorrect.

Richard Gregory was a pioneer of cognitive psychology whose scientific work involved the development and interpretation of visual illusions, which he used as a tool to work out the perceptual mechanisms involved in the way that the visual world is normally perceived. Gregory's account of perception requires not only information from the retina but also knowledge-based processes required for top-down or indirect construction of perception: past experience, assumptions, and active processing. An alternative approach was advocated by the psychologist James Gibson who argued that depth and form are obtained directly from the visual image. Gibson also argued that an object's affordance is what it can offer the viewer in terms of potential action.

Ref: Land MF, Heard P. Richard Langton Gregory. 24 July 1923—17 May 2010. Biogr. Mems Fell. R. Soc. 64:163—182.
Gregory RL. The Medawar lecture 2001 knowledge for vision: Vision for knowledge. Philosophical Transactions of the Royal Society B: Biological Sciences. 2005 Jun 29;360(1458):1231-51.

The correct answer is: Perception involves making inferences about past experiences

Question **36**

Incorrect

Mark 0.00 out of 1.00

Flag question

Sherif and Hovland's assimilation-contrast theory involves which of the following?

Select one:

- Aerial perspective
- Individual assertiveness
- Attachment theory
- Group norms ✖
- Persuasive communication

Your answer is incorrect.

Assimilation-contrast theory is a theory of judgement and attitude change. A consequence of this theory is that an extreme and ego-involving judgement or attitude tends to be polarised (to become even more extreme) in response to most types of information or persuasive communication.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015. p. 58.

The correct answer is: Persuasive communication

Question **37**

Correct

Mark 1.00 out of 1.00

Flag question


Reciprocal inhibition is central to which of the following psychotherapeutic techniques?

Select one:

- Implosion
- Floodling
- Mindfulness
- Exposure and response prevention
- Systematic desensitisation ✔

Your answer is correct.

Systematic desensitisation (Wolpe) is frequently used in the treatment of phobic anxiety disorders. It is based on the behavioural principle of reciprocal inhibition (i.e. anxiety and relaxation cannot coexist).



Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 908.

The correct answer is: Systematic desensitisation

Question **38**

Incorrect

Mark 0.00 out of 1.00

Flag question

What is true about prelinguistic communication?

Select one:

- It includes eye gaze, but not gestures
- It is not used in adolescence or adulthood
- Skills learned in this period are lost over later life
- It has no bearing on communication as an adult ✖
- This period is extended in children with autism spectrum disorder

Your answer is incorrect.

The prelinguistic stage is viewed as the period between birth and when an individual begins to use words/signs meaningfully. It is a time when children typically increase their ability to communicate with others, first using eye gaze, attending, and social-emotional affect and later adding gestures and other nonverbal means to communicate. Individuals with significant intellectual disability, including autism spectrum disorder, can have substantially protracted prelinguistic periods of communication and language development. For some, a singular reliance on prelinguistic communication may continue into adolescence or adulthood. The skills typically learned during this stage are critical to effective communication throughout the lifespan.

Ref: Crais E, Ogletree BT. Prelinguistic communication development. Prelinguistic and minimally verbal communicators on the autism spectrum. 2016:9-32.

The correct answer is: This period is extended in children with autism spectrum disorder

Extrapyramidal symptoms (EPS) are produced at what percentage of D2 occupancy?

Select one:

- 0.8 ✔
- 0.6
- 0.2
- 0.4
- 0.5

Your answer is correct.

The incidence of EPS is significantly correlated with the D2 receptor occupancy in the substantia nigra and striatum areas. Approximately 75-80% blockade of D2 receptors by antipsychotics can lead to the induction of EPS. Notably, this does not apply to D2 partial agonists as they show high D2R occupancies but still lower odds of EPS.

Ref: Siafis S, Wu H, Wang D, Burschinski A, Nomura N, Takeuchi H, Schneider-Thoma J, Davis JM, Leucht S. Antipsychotic dose, dopamine D2 receptor occupancy and extrapyramidal side-effects: a systematic review and dose-response meta-analysis. Molecular Psychiatry. 2023 Aug 3:1-1.

The correct answer is: 0.8

Question 40

Correct

Mark 1.00 out of 1.00

Flag question

Which of the following is not seen in serotonin syndrome?

Select one:

- Tremors
- Nausea
- Diplopia ✔
- Shivering
- Mydriasis

Your answer is correct.

Serotonin syndrome is characterised by altered mental state, agitation, tremors, shivering, diarrhoea, hyperreflexia, myoclonus, ataxia, and hyperthermia. Autonomic signs and symptoms include hyperthermia (that may be secondary to prolonged seizure activity, rigidity, or muscular hyperactivity), GI upset (including nausea and diarrhoea), mydriasis, tachycardia, and hyper/hypo-tension.

Your answer is correct.

Serotonin syndrome is characterised by altered mental state, agitation, tremors, shivering, diarrhoea, hyperreflexia, myoclonus, ataxia, and hyperthermia. Autonomic signs and symptoms include hyperthermia (that may be secondary to prolonged seizure activity, rigidity, or muscular hyperactivity), GI upset (including nausea and diarrhoea), mydriasis, tachycardia, and hyper/hypo-tension.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 1022.
Francescangeli J. et al. The serotonin syndrome: from molecular mechanisms to clinical practice. International journal of molecular sciences. 2019 May 9;20(9):2288.

The correct answer is: Diplopia

Question **41**

Correct

Mark 1.00 out of 1.00

Flag question

Metabotropic receptors are

Select one:

- Ligand gated K channels
- Ligand gated Na channels
- G protein coupled ✔
- Ligand gated Ca channels
- Ligand gated Cl channels

Your answer is correct.

Metabotropic receptors are linked to G proteins. Metabotropic receptors get their name from the fact that the movement of ions through them depends on certain metabolic processes. While they do not have ion channels included in their own structure (they are monomeric proteins with an extracellular neurotransmitter binding domain), the activation of intermediate molecules called G-proteins (an intracellular domain) enable eventual ion movement. For a self-explanatory figure, see https://www.ncbi.nlm.nih.gov/books/NBK10855/fig

Ref: Stahl SM. Stahl's Essential Psychopharmacology: Neuroscientific Basis and Practical Applications, 5th Edition. 2021. p. 100.

The correct answer is: G protein coupled

Question **42**

Correct

Mark 1.00 out of 1.00

Flag question

Which of the following medications works by acetylcholinesterase inhibition?

Select one:

- Galantamine ✔
- Pimavanserin
- Brexpiprazole
- Memantine
- Bupropion

Your answer is correct.

Galantamine is an inhibitor of the enzyme acetylcholinesterase (AChE). Central cholinergic neurons are important for regulation of memory, and thus, in the central nervous system, the boost of acetylcholine caused by AChE blockade contributes to improved cognitive functioning.

Ref: Stahl SM. Stahl's Essential Psychopharmacology: Prescriber's Guide, 7th Edition. 2021. p. 517.

The correct answer is: Galantamine

Question **43**

Correct

Mark 1.00 out of 1.00

Flag question

Which of the following physiologic/pharmacokinetic changes is seen in pregnancy?

Select one:

- Increased glomerular filtration rate (GFR) ✔
- Increased binding of drugs to albumin
- Decreased volume of distribution of water-soluble drugs
- Decreased cardiac output
- Decreased renal blood flow

Your answer is correct.

Increased cardiac output of pregnancy is accompanied by an increase in renal blood flow, which, in conjunction with haemodilution and a reduction in oncotic pressure from reduced plasma proteins, increases the GFR. Albumin concentrations are 70-80% lower compared to non-pregnancy. Consequently, drugs that are highly protein bound will typically have greater free fractions in pregnancy. Fluid volume increases in pregnancy. Depending on a number of factors, the volume of distribution for hydrophilic medications may be as large as the volume of total body water. In pregnancy, the increase in total body water (and thus volume of distribution) leads to reduced plasma and tissue concentrations for the same initial dose compared to the nonpregnant state.

Ref: Pinheiro EA, Stika CS. Drugs in pregnancy: pharmacologic and physiologic changes that affect clinical care. In Seminars in perinatology 2020 Apr 1 (Vol. 44, No. 3, p. 151221). WB Saunders.

The correct answer is: Increased glomerular filtration rate (GFR)

Question **44**

Correct

Mark 1.00 out of 1.00

Flag question

What is the primary mechanism of action of memantine?

Select one:

- 5HT-2A antagonism
- NMDA agonism
- Cholinesterase inhibition
- Non-competitive NMDA antagonism ✔
- 5HT-1A agonism

Your answer is correct.

Memantine is a non-competitive, low-affinity NMDA receptor antagonist. It also has weak 5HT-3 antagonist properties.

Ref: Stahl SM. Stahl's Essential Psychopharmacology: Prescriber's Guide, 7th Edition. 2021. p. 522.

The correct answer is: Non-competitive NMDA antagonism

Question 45

Incorrect

Mark 0.00 out of 1.00

Flag question

Where are the muscarinic receptors of the peripheral nervous system mainly found?

Select one:

- Vascular endothelial cells
- Sympathetic ganglia
- Peripheral postsynaptic receptors
- Neuromuscular junctions ✖
- Parasympathetic ganglia

Your answer is incorrect.

Muscarinic acetylcholine receptors are G-protein coupled metabotropic receptors. Various subtypes are distributed predominantly in the CNS. Iin the peripheral nervous system, they are found primarily on autonomic effector cells innervated by postganglionic parasympathetic nerves.

Ref: Brown, Joan Heller, et al. "Muscarinic Receptor Agonists and Antagonists." Goodman & Gilman's: The Pharmacological Basis of Therapeutics, 13e Eds. Laurence L. Brunton, et al. McGraw Hill, 2017.

The correct answer is: Parasympathetic ganglia

Question **46**

Incorrect

Mark 0.00 out of 1.00

Flag question

What is the purpose of naloxone being mixed with buprenorphine in the treatment of opioid dependence?

Select one:

- Reduces the risk of relapse ✖
- Reduces the risk of misuse/diversion
- Increases the efficacy of treatment
- Reduces the cost of medication
- Reduces the duration of treatment

Your answer is incorrect.

A buprenorphine/naloxone preparation (suboxone) may reduce the risk of diversion. The different sublingual and parenteral absorption profiles of buprenorphine and naloxone are the key factors: if used sublingually, the naloxone will have negligible effects. However, if the combined preparation is injected, the naloxone will have a substantial effect and will attenuate the effects of buprenorphine in the short term and is also likely to precipitate withdrawal in opioid-dependent individuals on full opioid agonists.

Ref: Taylor DM, Barnes TRE, Young AH. The Maudsley Prescribing Guidelines in Psychiatry, 14th Edition. 2021. p. 490.

The correct answer is: Reduces the risk of misuse/diversion

Question **47**

Incorrect

Mark 0.00 out of 1.00

Flag question

What is the first line treatment of clozapine-induced constipation?

Select one:

- Methylcellulose
- Senna
- Lactulose ✖
- Macrogols
- Docusate

Your answer is incorrect.

Stimulant laxatives (senna, bisacodyl or sodium picosulfate) are the first-line treatments for clozapine-induced constipation, if the individual has not passed a bowel movement in two days, after excluding intestinal obstruction. If the laxative is not working after 24 hours, consider adding an osmotic laxative (macrogols or lactulose) or emollient stool softeners (docusate sodium). Regular lactulose and plenty of fluids is important part of management. Avoid bulk-forming laxatives because the cause of constipation in this case is gastric hypomotility. Faecal impaction may require treatment with high dose macrogol or repeated doses of suppositories.

Ref: Taylor DM, Barnes TRE, Young AH. The Maudsley Prescribing Guidelines in Psychiatry, 14th Edition. 2021. p. 212.

The correct answer is: Senna

Question **48**

Correct

Mark 1.00 out of 1.00

Flag question

When a child is obedient to rules and there is parental responsiveness without rigidity, what is the most likely style of parenting?

Select one:

- Neglectful/uninvolved
- Authoritative ✔
- Permissive/indulgent
- Authoritarian
- Overbearing

Your answer is correct.

The authoritative parenting style is characterised as high in responsiveness and demandingness. Authoritative parents provide not only support and warmth, but also clearly defined rules and consistent discipline.

Ref: Bi X et al. Parenting styles and parent–adolescent relationships: The mediating roles of behavioral autonomy and parental authority. Frontiers in psychology. 2018 Nov 13;9:2187.

The correct answer is: Authoritative

Question **49**

Partially correct

Mark 2.00 out of 3.00

Flag question

Family Systems in Development

Identify the correct term for each of the following descriptions.

1. Family members' interactions are influenced by each other, which in turn are influenced by others. Complementarity ✗

2. Over-involved relationships that develop from diffuse boundaries within family systems. Enmeshment ✓

3. Involvement of a child in a parental conflict to defuse tension within the interparental subsystem. Triangulation ✓

EXPLANATION: Circular causality focuses on the reciprocal relationship between two events. While linear causality implies a process in which one event causes the occurrence of another, circular causality implies that one has influence over the other. Circular questioning is used in family therapy to yield information about the dynamics and relationships in a family.

Salvador Minuchin used the term enmeshment to describe the overinvolved relationships that develop from diffuse boundaries within family systems and between family members and other systems.

In the family context, triangulation refers to the involvement of a child in a parental conflict to defuse tension within the interparental subsystem, and generally includes parents' pressure on children to become involved as well as children's self-directed attempts to intervene.

Ref: Mantova KS. THEORETIC OVERVIEW OF CIRCULAR QUESTIONS IN FAMILY AND SYSTEMIC THERAPY. KNOWLEDGE-International Journal. 2022;54(1):187-91.
D'Astice, T., Russell, W.P. Enmeshment in Couples and Families. In: Lebow, J.L., Chambers, A.L., Breunlin, D.C. (eds) Encyclopedia of Couple and Family Therapy.

CORRECT ANSWERS:
1) Family members' interactions are influenced by each other, which in turn are influenced by others. – Circular causality
2) Over-involved relationships that develop from diffuse boundaries within family systems. – Enmeshment
3) Involvement of a child in a parental conflict to defuse tension within the interparental subsystem. – Triangulation

EXPLANATION: P-glycoprotein, commonly referred to as P-gp, is a membrane transporter within the ATP-binding cassette (ABC) transporter family. It is encoded by the ABCB1 or MDR1 gene, located on chromosome 7. P-gp functions as an efflux pump, contributing to drug absorption, distribution, and elimination.

Alpha synuclein is a small protein that is highly charged and coded by the SNCA (synuclein) gene. Alpha synuclein is predominantly found at the presynaptic terminals, where it associates with synaptic vesicles. Alpha synuclein is involved in synucleinopathies like Parkinson disease, multiple systems atrophy, and dementia with Lewy bodies.

The serotonin transporter (SERT), encoded by the SLC6A4 gene located on the long arm of chromosome 17 (17q11-17q12), plays a critical role in regulating serotonin neurotransmission in the brain. SERT is responsible for serotonin reuptake from the synaptic cleft into presynaptic neurons, thereby regulating the amount of serotonin available to bind to serotonin receptors in the brain. The most investigated region of the SLC6A4 gene is the serotonin transporter gene promoter region (5-HTTLPR). Several classes of antidepressant drugs, including SSRIs, SNRIs, and TCAs, target SERT and inhibit its activity.

Question **50**

Partially correct

Mark 1.00 out of 3.00

Flag question

Gene Coding

Choose the gene that codes for each of the following.

1. P-glycoprotein CYP1A2 ✖

2. Alpha-synuclein SNCA ✔

3. Serotonin transporter SNCA ✖

EXPLANATION: P-glycoprotein, commonly referred to as P-gp, is a membrane transporter within the ATP-binding cassette (ABC) transporter family. It is encoded by the ABCB1 or MDR1 gene, located on chromosome 7. P-gp functions as an efflux pump, contributing to drug absorption, distribution, and elimination.

Alpha synuclein is a small protein that is highly charged and coded by the SNCA (synuclein) gene. Alpha synuclein is predominantly found at the presynaptic terminals, where it associates with synaptic vesicles. Alpha

Question **51**

Partially correct

Mark 2.00 out of 3.00

Flag question

Memory & forgetting

Choose the term that best applies to each of the following descriptions.

1. Partial or total inability to recall past experiences and events.

Amnesia ✔

2. Fluent plausible lying in those without organic brain pathology.

Pseudologia fantastica ✔

3. When new memories impede the retrieval of old material.

Proactive interference ✖

EXPLANATION: Amnesia is defined as partial or total inability to recall past experiences and events. Its origin may be organic or

Parkinson disease, multiple systems atrophy, and dementia with Lewy bodies.

The serotonin transporter (SERT), encoded by the SLC6A4 gene located on the long arm of chromosome 17 (17q11-17q12), plays a critical role in regulating serotonin neurotransmission in the brain. SERT is responsible for serotonin reuptake from the synaptic cleft into presynaptic neurons, thereby regulating the amount of serotonin available to bind to serotonin receptors in the brain. The most investigated region of the SLC6A4 gene is the serotonin transporter gene promoter region (5-HTTLPR). Several classes of antidepressant drugs, including SSRIs, SNRIs, and TCAs, target SERT and inhibit its activity.

Ref: Radosavljevic M et al. The Role of Pharmacogenetics in Personalizing the Antidepressant and Anxiolytic Therapy. Genes. 2023 May 16;14(5):1095.
Simuni T, et al. A biological definition of neuronal a-synuclein disease: towards an integrated staging system for research. The Lancet Neurology. 2024 Feb 1;23(2):178-90.

CORRECT ANSWERS:
1) P-glycoprotein – ABCB1/MDR1
2) Alpha synuclein – SNCA
3) Serotonin transporter – SLC6A4

EXPLANATION: Amnesia is defined as partial or total inability to recall past experiences and events. Its origin may be organic or psychogenic.

Pseudologia fantastica or fluent plausible lying (pathological lying) describes confabulation that can be seen in those without organic brain pathology, such as personality disorders/difficulties of the antisocial or dissocial type.

Ordinary memory failure can be caused by interference from related material. In proactive interference, old memories interfere with new learning and hence with recall, while in retroactive interference new memories interfere with the retrieval of old material.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. pp. 64-69.

CORRECT ANSWERS:
1) Partial or total inability to recall past experiences and events. – Amnesia
2) Fluent plausible lying in those without organic brain pathology. – Pseudologia fantastica
3) When new memories impede the retrieval of old material. – Retroactive interference