Started on	Saturday, 23 July 2022, 11:08 AM
State	Finished
Completed on	Saturday, 23 July 2022, 11:08 AM
Time taken	17 secs
Marks	0.00/63.00
Grade	**0.00** out of 10.00 (**0**%)

Question **1**

Not answered

Marked out of 1.00

Flag question

A psychology student is keen to know about the ICD-10 duration criteria for the diagnosis of mania. The episode should last at least:

Select one:

- 2 days
- 1 day
- 4 weeks
- 2 weeks
- 1 week

Your answer is incorrect.

Both ICD-10 and DSM-4 specify the manic episode should last at least 1 week and the disturbance is sufficiently severe to impair social and occupational functioning.

The correct answer is: 1 week

Question **2**

Not answered

Marked out of 1.00

Flag question

The symptoms of adjustment disorder resolve within:

Select one:

- 1 month of termination of the stressor

Your answer is incorrect.

Adjustment disorder requires a stressor and that the symptoms should resolve within 6 months of termination of the stressor or its consequences.

The correct answer is: 6 months of termination of the stressor

Question 3

Not answered

Marked out of 1.00

Flag question

What is the most common psychiatric diagnosis among those presenting to emergency departments following self-harm?

Select one:

- Post-traumatic stress disorder
- Depression
- Adjustment disorder
- Panic disorder
- Schizophrenia

Your answer is incorrect.

The most common psychiatric diagnosis among those presenting to emergency departments following self-harm is adjustment disorder.

(Ref: Taggart et al; Gen Hosp psychiatry 2006: 28: 330-5)

The correct answer is: Adjustment disorder

Question 4

Not answered

Marked out of 1.00

Flag question

According to the DSM-5, which of the following is true regarding changes to the diagnostic criteria of psychotic illnesses?

- Catatonia is strongly associated with schizophrenia

Your answer is incorrect.

The DSMV is the 2013 update to the American Psychiatric Association's (APA) classification and diagnostic tool. Changes were made in the diagnostic criteria and coding of Schizophrenia spectrum and other psychotic disorders. This included the removal of all subtypes of schizophrenia (paranoid, disorganized, catatonic, undifferentiated, and residual).

The correct answer is: All subtypes of schizophrenia were removed

Question 5

Not answered

Marked out of 1.00

Flag question

During Fresher's Week, Kaley, a 20-year-old law student, with no prior psychiatric history, presents to Accident and Emergency with delirium, labile blood pressure, dry mouth and dilated pupils. Which of the following could explain her presentation?

Select one:

- Neuroleptic malignant syndrome
- Paracetamol overdose
- Anti-muscarinic psychosis
- Cocaine intoxication
- Serotonin Syndrome

Your answer is incorrect.

Cocaine is a stimulant made from an alkaloid contained in the leaves of the coca bush, Erythroxylon coca. Crack or rock cocaine is a ready-to-smoke, potent, relatively inexpensive, and highly addictive freebase forms of cocaine. The patient in question is presenting with typical symptoms of cocaine intoxication. (Ref. University of Southern California).

The correct answer is: Cocaine intoxication

Question 6

Not answered

Marked out of 1.00

Flag question

- ICD 2
- ICD 1
- ICD 9

Your answer is incorrect.

The World Health Organisation's ICD-9, published in 1977, listed homosexuality as a mental illness. It was later removed from the ICD-10, a decision that was endorsed by the 43rd World Health Assessment on 17th May 1990.

The correct answer is: ICD 10

Question 7

Not answered

Marked out of 1.00

Flag question

Mrs. Dellmore, who is 83 years old, has been referred to a clinic for older adults following concerns that she has been forgetting the names of common objects, and losing her way on returning from the local shops. The psychologist uses a number of rating measures. Which one ofthe following is observer-rated?

Select one:

- Meyers Briggs
- NPI-C
- SADQ
- PHQ-9
- GAD-7

Your answer is incorrect.

The Neuropsychiatric Inventory- Clinician rating scale is observer-rated, focussing on neuropsychiatric symptoms in people with dementia. The other scales are self-rated (A and B for depression and anxiety, D for alcohol, and Meyers Briggs for personality types).

The correct answer is: NPI-C

Question 8

Not answered

Select one:

- Hypercholesterolemia
- Atrial fibrillation
- Obesity
- Diabetes
- Hypertension

Your answer is incorrect.

The Syst-Eur and PROGRESS studies showed small but significant reductions in dementia from treating hypertension (Forette 2002; Tzourio 2003). Hypertension is now considered as strongest treatable risk factor for vascular dementia.

The correct answer is: Hypertension

Question 9

Not answered

Marked out of 1.00

Flag question

'Driving a car' is an example of what type of memory?

Select one:

- Semantic
- Working
- Declarative
- Long term memory
- Episodic

Your answer is incorrect.

Driving a car is an example of procedural memory, which is classified under non-declarative memory. Non-declarative memory is a component of long-term memory.

The correct answer is: Long term memory

Question 10

Not answered

Marked out of 1.00

- Digital span
- Similarities
- Picture completion
- Digit symbol
- Block design

Your answer is incorrect.

Tests on WAIS are either ‘hold' or ‘no hold'. The so-called ?hold tests in WAIS are thought to reflect the use of old knowledge and are relatively resistant to the effects of brain damage and ageing. These include vocabulary, information, object assembly, and picture completion. ?No hold tests require speed of response, working memory, or the creation of new relations between unrelated items. These are more likely to show early decline with ageing and cognitive impairment. These include digit symbol, digit span, similarities, and block design. This is the reason why WAIS is not the best test to measure premorbid intelligence.

The correct answer is: Picture completion

Question 11

Not answered

Marked out of 1.00

Flag question

An ECG should be carried out on presentation in a patient with anorexia nervosa if BMI is less than;

Select one:

- 25
- 16
- 20
- 22
- 19

Your answer is incorrect.

In patients with anorexia, a screening assessment of risk should include a minimum of blood pressure, pulse, BMI, temperature and tests of proximal myopathy. An electrocardiogram (ECG) should be carried out if weight is low (BMI <16). Where a low BMI combines with abnormalities on ECG, rapid gradient of weight loss, abnormal electrolytes and muscle wasting, these can be considered ‘rough and ready' predictors of higher physical risk. ECG is recommended routinely if

Not answered

Marked out of 1.00

Flag question

Andrew is a 10-day-old genetic male who appears not to have developed the characteristic of male genitalia as expected. The cause is thought to be linked to excessive virilisation in female adults. Which condition does this relate to?

Select one:

- 3-beta-yydroxysteroid dehydrogenase
- 5-beta-reductase deficiency
- 17-beta-hydroxysteroid dehydrogenase deficiency
- 17-alpha-hydroxysteroid dehydrogenase deficiency
- 5-alpha-reductase deficiency

Your answer is incorrect.

17-beta-hydroxysteroid dehydrogenase deficiency is an autosomal recessive form of male pseudohermaphroditism due to impaired testicular conversion of androstenedione to testosterone. It can produce impaired virilization of genetically male infants & children and excessive virilization of female adults. It affects male sexual development and their bodies do not produce enough of testosterone.

The correct answer is: 17-beta-hydroxysteroid dehydrogenase deficiency

Question 13

Not answered

Marked out of 1.00

Flag question

Which of the following conditions is associated with polycystic ovarian syndrome?

Select one:

- Binge eating disorder
- Anorexia nervosa
- Muscle dysmorphia
- Body dysmorphic disorder
- Bulimia nervosa

Question 14

Not answered

Marked out of 1.00

Flag question

During a cognitive examination, Betty, an 85-year-old former shop assistant is asked to read a telephone number and then dial the number without looking at it. What type of memory is being tested here?

Select one:

- Iconic memory
- Recognition memory
- Episodic memory
- Semantic memory
- Haptic memory

Your answer is incorrect.

Iconic memory is the memory of visual stimuli. It is part of the visual memory system that includes visual short-term memory and long-term memory. It is a sensory memory that is thought to last milliseconds before disappearing.

The correct answer is: Iconic memory

Question 15

Not answered

Marked out of 1.00

Flag question

Which one among the following is an environmental factor associated with ADHD?

Select one:

- Essential fatty acid deficiencies
- Prematurity
- Food and additive allergies
- Low-protein

inconsistent parenting, high expressed emotion, family conflict, early institutional rearing) and Acquired neurobiological risks (e.g. head trauma) Ref: Asherson 2005. All others listed above are biochemical risk factors for ADHD.

The correct answer is: Prematurity

Question 16

Not answered

Marked out of 1.00

Flag question

Betty is then shown a list of words. After a few moments she is shown another list of words and asked to identify words that appear in both lists.

Select one:

- Semantic memory
- Iconic memory
- Episodic memory
- Recognition memory
- Haptic memory

Your answer is incorrect.

Recognition memory is the ability to recognize previously encountered events, objects or people.

The correct answer is: Recognition memory

Question 17

Not answered

Marked out of 1.00

Flag question

Your colleague Bill asks you to call a GP practice on his behalf. He reads out loud the number for you, and you manage to remember it long enough to dial it. What kind of memory does this illustrate?

Select one:

- Episodic memory
- Working memory

Your answer is incorrect.

Working memory is the ability to process and remember information at the same time, and usually hold small amounts of information (typically around 7 items). It is how we process information such as phone numbers, which we can remember sufficiently long to perform the task of dialling it.

The correct answer is: Working memory

Question 18

Not answered

Marked out of 1.00

Flag question

A child attains gender identity in which stage of Freud's psychosexual development?

Select one:

- Anal phase
- Genital phase
- Phallic phase
- Oral phase
- Latent phase

Your answer is incorrect.

Explanation:

The phallic stage begins sometime during year 3 and continues until approximately the end of year 5. The phallic stage lays the foundation for gender identity and serves to integrate residues from the previous stages into a predominantly genital-sexual orientation.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1062-1063.

The correct answer is: Phallic phase

Question 19

Not answered

Marked out of 1.00

Flag question

- Formal operational
- Concrete Operational
- Sensorimotor
- Pre-operational

Your answer is incorrect.

Explanation.

This child is displaying animism. In Piaget's pre-operational stage, children use animistic thinking, which is the tendency to endow physical events and objects with life-like psychological attributes, such as feelings and intentions.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1008.

Stage	Age	Piaget's Cognitive Developmental Characteristics
Sensorimotor	0-2 years	Divided into six phases, characterised by: 1. Inborn motor and sensory reflexes 2. Primary circular reaction - coordinates activities of own body and five senses 3. Secondary circular reaction – seeks out new stimuli 4. Use of familiar means to obtain ends; plays peek-a-boo 5. Tertiary circular reaction and discovery through active experimentation 6. Insight and object permanence
Preoperational	2-7 years	Deferred imitation, symbolic play, graphic imagery (drawing), mental imagery, and language Egocentrism, phenomenalistic causality, and animistic thinking
Concrete operations	7-11 years	Conservation of quantity, weight, volume, length, and time based on reversibility by inversion or reciprocity; operational thought; class inclusion and seriation Syllogistic reasoning – logical conclusion formed from two premises
Formal operations	11 years-end of adolescence	Combinatorial system, whereby variables are isolated and all possible combinations are examined; hypotheticodeductive reasoning Complex use of language and abstract thinking

Ref: Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition, 2022

SPMM COURSE

The correct answer is: Pre-operational

Question 20

Not answered

Marked out of 1.00

Flag question

Pauline is a first-time mother to Stephanie, her 18-month-old daughter. She is worried that Stephanie isn't talking as well as the other children at the nursery. If an 18-month-old child has

- 85-125
- 0-10

Your answer is incorrect.

Explanation:

By age 18 months, a child is generally using 20 single words. They will understand up to 150 words.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 889.

The correct answer is: 20-50

Question 21

Not answered

Marked out of 1.00

Flag question

In your CAMHS clinic, you observe Mollie, a five-year-old girl, as she plays with her doll. You notice that she is talking to her doll as though it is real. What is the name of this phenomenon?

Select one:

- Artificialism
- Animism
- Centration
- Symbolic play
- Egocentrism

Your answer is incorrect.

Explanation:

Animism is the belief that plants, inanimate objects and natural phenomena possess a living soul and have human-like feelings and intentions. In Piaget's pre-operational stage, children use such animistic thinking to endow physical events and objects with life-like psychological attributes, such as feelings and intentions.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1008.

The correct answer is: Animism

According to Kohlberg, which one among the following represent Conventional Stage?

Select one:

- Authority
- Obedience
- Social Contract
- Punishment
- Universal Ethics

Your answer is incorrect.

Explanation:

Lawrence Kohlberg integrated Jean Piaget's concepts and described multiple stages of moral development within three significant levels of morality. The first level is pre-conventional morality, in which punishment and obedience are the determining factors. The second level is morality of conventional role-conformity, in which children try to conform to gain approval and maintain good relationships with others and are also guided by rigid codes of 'law and order' (authority orientation). The third and highest level is the morality of self-accepted moral principles (post-conventional), in which children voluntarily comply with rules based on a concept of universal ethical principles and make exceptions to rules in certain circumstances.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 902.

Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: Authority

Question 23

Not answered

Marked out of 1.00

Flag question

Which of the following is seen in a normal child development?

Select one:

- Babbling with consonant sounds at 7 months
- Toilet training at 5 years
- Beginning to crawl at 15 months
- Social smile at 6 months
- Putting two words together at 4 years

Question 24

Not answered

Marked out of 1.00

Flag question

In a strange situation experiment, Bobby, a 2-year-old boy, is only comforted when his mother returns to the room and is not comforted by the stranger. What type of attachment is seen here?

Select one:

- Anxious Resistant
- Anxious Avoidant
- Secure attachment
- Disorganized
- Insecure

Your answer is incorrect.

Explanation:

Mary Ainsworth devised the Strange Situation procedure with 12- to 18-month infants, which categorised infant-parent relationships. The procedure involves separations and reunions with caregivers and observes response. In secure attachment, the child is distressed by separation but can quickly be soothed on reunion with the caregiver.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 658-659.

The correct answer is: Secure attachment

Question 25

Not answered

Marked out of 1.00

Flag question

Which of the following is NOT a risk factor for delayed speech development?

Select one:

- Smoking in pregnancy
- Small family size

Your answer is incorrect.

The important family risk factors associated with delayed language development in children include large family size, smoking in pregnancy, rapid successive pregnancies, absence of father, low levels of parental education, divorce, marital disharmony & violence.

The correct answer is: Small family size

Question 26

Not answered

Marked out of 1.00

Flag question

Which of the following has the strongest correlation with developmental language delay?

Select one:

- Low socio-economic status of mother
- High fibre diet
- Maternal smoking during pregnancy
- Immigration
- Maternal alcohol use during pregnancy

Your answer is incorrect.

Smoking during pregnancy and large family size are the strongest risk factors for developmental language delay

The correct answer is: Maternal smoking during pregnancy

Question 27

Not answered

Marked out of 1.00

Flag question

The preferential attachment of a baby to one adult female figure is called;

Select one:

- Universal attachment
- Proximity seeking

Your answer is incorrect.

Explanation:

Attachment theory originated in the work of John Bowlby. Being **monotropic**, infants tend to attach to one person, but they can form attachments to several people. A basic sense of security and safety is derived from a continuous and close relationship with a caregiver, usually the mother.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1011.

The correct answer is: Monotropy

Question **28**

Not answered

Marked out of 1.00

Flag question

Angela Stubbs has been participating in psychotherapy for the past seven months. She has the sense that the process has been helping her develop a more robust sense of self, a feeling of continuity, and of affiliation with others. What term best describes this process?

Select one:

- Symbiosis
- Deindividuation
- Identity diffusion
- Individuation
- Introjection

Your answer is incorrect.

Explanation:

Carl Jung described individuation as “becoming a single, homogeneous being, and, in so far as ‘individuality’ embraces our innermost, last, and incomparable uniqueness, it also implies becoming one’s own self. We could therefore translate individuation as ‘coming to selfhood’ or ‘self-actualisation’."

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: Individuation

Question **29**

Not answered

Marked out of 1.00

- Premack's principle
- Aversive conditioning
- Covert reinforcement
- Avoidance learning
- Covert sensitization

Your answer is incorrect.

Aversive conditioning: This is an operant conditioning where punishment is used to reduce the frequency of target behaviour e.g. the use of Disulfiram (noxious stimuli) to reduce the frequency of drinking alcohol.

The correct answer is: Aversive conditioning

Question 30

Not answered

Marked out of 1.00

Flag question

Timmy has a pet mouse which he spends time with. As part of a psychological experiment, his father creates a loud noise whenever Timmy is playing with his mouse, and the noise frightens Timmy. This is done a number of times, so that whenever Timmy sees his pet mouse, he feels frightened. Which of these is now classed as the conditioned stimulus?

Select one:

- The mouse + noise
- The fear + noise
- The mouse
- The noise
- The fear

Your answer is incorrect.

The mouse was previously a neutral stimulus; it has been associated with an unconditioned stimulus (the noise) that naturally triggers an automatic response in Timmy (fear). The mouse therefore becomes a conditioned stimulus, eliciting fear in Timmy.

The correct answer is: The mouse

A psychologist is involved in treating a patient with fear of heights. He requested the patient to imagine standing on top of the building without actual real exposure to them. The psychological principle employed here is:

Select one:

- Punishment
- Flooding
- Reciprocal inhibition
- Implosion
- Aversion

Your answer is incorrect.

Flooding is an operant conditioning technique where actual exposure to feared stimulus takes place for a substantial amount of time so the accompanying anxiety response fades away while the stimulus is continuously present. This will lead to extinction of fear. When a similar technique is attempted with imagined exposure then it is called **implosion**.

The correct answer is: Implosion

Question **32**

Not answered

Marked out of 1.00

Flag question

A teacher rewarded Jimmy with star sticker for his good behaviour. It represents which type of learning?

Select one:

- Social learning
- Cognitive learning
- Classical conditioning
- Operant conditioning
- Aversive learning

Your answer is incorrect.

Explanation:

behaviour is an example of positive reinforcement.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. pp. 907-908.

The correct answer is: Operant conditioning

Question 33

Not answered

Marked out of 1.00

Flag question

Mohammed is a 7-year-old boy. His mother told him that he could play outside once he had finished his homework. This is based on which of the following psychological mechanisms?

Select one:

- Operant conditioning
- Premack principle
- Classical conditioning
- Reciprocal inhibition
- Cognitive mapping

Your answer is incorrect.

Explanation:

The Premack principle asserts that access to a preferred action will reinforce any action that is less preferred. Any behaviour that is preferred to a second behaviour will theoretically reinforce the second behaviour. This is also known as ‘Grandma‘s rule' – ‘eat the carrot and get the dessert'.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1019.

The correct answer is: Premack principle

Question 34

Not answered

Marked out of 1.00

Flag question

A grandmother says to her grandchild "You must eat your greens before you go to play outside". Which psychological mechanism is used here?

- Reciprocal inhibition

Your answer is incorrect.

Explanation:

The Premack principle asserts that access to a preferred action will reinforce any action that is less preferred. Any behaviour that is preferred to a second behaviour will theoretically reinforce the second behaviour. The Premack principle is also called relative theory of reinforcement, often used with young children. It states that more probable behaviour (like play outside) will reinforce less probable behavior (like eating green leafy vegetables).

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1019.

The correct answer is: Premack principle

Question **35**

Not answered

Marked out of 1.00

Flag question

Sammy Taylor is coming to see you in clinic as an emergency book-in. He is 19 years old and has developed some voice-hearing experiences and bizarre thoughts regarding the MI5. Which of these might be a precipitating factor?

Select one:

- Having dropped out of school prematurely at the age of 15
- Having lost his adoptive father to a road traffic accident two weeks previously
- Having had a traumatic birth
- Having a brother who has a diagnosis of schizophrenia
- Having been adopted at the age of 7 months

Your answer is incorrect.

Explanation:

The others may be contributing factors, but most are predisposing rather than precipitating. A precipitating factor has a strong temporal link to the onset of the episode of mental illness (e.g. it occurs just before).

Four 'Ps' of case formulation:

Predisposing factors: those that put a person at risk of developing a problem; may include genetics, previous life events, or temperament

pediatric psychology. 2016 Mar 1;41(2):159-81.

The correct answer is: Having lost his adoptive father to a road traffic accident two weeks previously

Question 36

Not answered

Marked out of 1.00

Flag question

According to the World Health Organization, how many women have experienced intimate partner violence or non-partner sexual violence in their lifetime?

Select one:

- 1 in 20
- 1 in 10
- 1 in 3
- 1 in 5
- 1 in 100

Your answer is incorrect.

Explanation:

According to the World Health Organisation, almost 1 in 3 women have experienced one or both of these forms of violence (intimate partner violence or non-partner sexual violence) at least once in their lifetime. This recent data confirms that physical and sexual intimate partner violence and sexual violence more broadly remain pervasive in the lives of women and adolescent girls across the globe.

Ref: World Health Organization. Violence against women prevalence estimates, 2018: global, regional and national prevalence estimates for intimate partner violence against women and global and regional prevalence estimates for non-partner sexual violence against women. 2021.

The correct answer is: 1 in 3

Question 37

Not answered

Marked out of 1.00

Flag question

- Individualism
- Democracy
- Communism

Your answer is incorrect.

Group rights, also known as collective rights, are rights held by a group rather than by its individual members, the latter of which is termed individualism.

The correct answer is: Collectivism

Question 38

Not answered

Marked out of 1.00

Flag question

William and Diane Corndale are seeking help for infertility. They have been trying to conceive for 5 years, but have so far been unsuccessful. Which of the following is the most likely consequence of this stressor?

Select one:

- Feelings of guilt
- Negative impact on self-esteem
- Decrease in frequency of sexual intercourse
- Increase in number of disagreements
- All of the listed options

Your answer is incorrect.

Difficulty in conceiving may place significant stress on a couple, and there's the potential for a wide spectrum of psychological struggles to ensue.

The correct answer is: All of the listed options

Question 39

Not answered

Marked out of 1.00

Flag question

- Suicide rates are higher in women than in men
- Suicide was more common in civilians than in soldiers
- The risk of suicide is fixed

Your answer is incorrect.

Explanation:

Emile Durkheim was a French sociologist who challenged the assumption that social processes can be understood in terms of the actions of individuals and argued that even suicide, apparently the ultimately individualistic act, cannot be fully explained at the purely individual level. In his book *Suicide*, Durkheim used survey data purporting to show that egoistic suicide, resulting from feelings of self-reproach and sense of failure, arises from a lack of social cohesion; that altruistic suicide, carried out for the benefit of others, arises from a sense of failure to society; and that anomic suicide, arising from a sense that life is pointless, arises from an absence of social norms that he called anomie.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: Suicide could be attributed to a lack of social cohesion

Question 40

Not answered

Marked out of 1.00

Flag question

Mr Loughton has come to see you in clinic, following a referral from the police. You are aware that your colleague will want to know what his future risk of sexual violence will be. Which of the following is the strongest predictive factor for future sexual violence?

Select one:

- History of past sexual violence
- Having numerous siblings
- History of alcohol misuse
- History of bipolar affective disorder
- History of depression

Your answer is incorrect.

Explanation:

History of sexual violence perpetration is the strongest predictor of sexual violence perpetration.

Question 41

Not answered

Marked out of 1.00

Flag question

Mandip is a 33-year-old Sikh male from India. He is an IT contractor and speaks English at work. He regularly goes to the pub after work for drinks with his colleagues. When he goes home he speaks Punjabi with his parents and wears traditional robes. What form of acculturation does Mandip display?

Select one:

- Marginalisation
- Division
- Separation
- Assimilation
- Integration

Your answer is incorrect.

Acculturation refers to the process of cultural change that takes place when an individual or a group comes in to continuous contact with a culturally distinct group. Acculturation can result from immigration and can occur in either direction – hosts can get accultured; this was evident in certain places in times of Colonial rule. Four types are described according to the degree of retention and adoption of the two cultures at ethnocultural group level. Integration refers to both high retention of one's own cultural values and high adoption of the practices of the new culture.

The correct answer is: Integration

Question 42

Not answered

Marked out of 1.00

Flag question

Empathy skills are most likely to be delayed in which of the following?

Select one:

- A hearing child with one deaf parent
- A hearing child of hearing parents
- A deaf child of hearing parents

Many deaf children from hearing families have seriously delayed development of empathy and theory of mind (ToM). They often continue to fail false belief and other standard preschool ToM tests well into middle childhood. However, natively signing deaf children of deaf parents develop ToM understanding on the same early timetable as hearing children. This suggests the importance of shared conversation with family members via a mutually accessible language (either signed or spoken) for the development of ToM. In deaf children of hearing parents empathy and ToM are significantly correlated, unlike in hearing children where these two are less dependent.

Ref: Peterson CC. Empathy and theory of mind in deaf and hearing children. Journal of deaf studies and deaf education. 2016 Apr 1;21(2):141-7.

The correct answer is: A deaf child of hearing parents

Question 43

Not answered

Marked out of 1.00

Flag question

Michael Stills is 32 years old and has developed depression for the first time in in his life. You meet him and take a history. Which of the following is a precipitating factor for his depression?

Select one:

- Growing up in a large city
- The recent loss of his baby son from cot death at the age of 4 months
- Having a parent who has a diagnosis of chronic anxiety disorder
- Being married for the past 2 years
- Working as a teaching assistant

Your answer is incorrect.

Explanation:

Four 'Ps' of case formulation:

Predisposing factors: those that put a person at risk of developing a problem; may include genetics, previous life events, or temperament

Precipitating factors: a specific event or trigger to the onset of the current problem

Perpetuating factors: those that maintain the problem once it has become established

Protective factors: strengths that reduce the severity of problems and promote healthy and adaptive functioning

Ref: Racine NM et al. Systematic review: predisposing, precipitating, perpetuating, and present factors predicting anticipatory distress to painful medical procedures in children. Journal of

Marked out of 1.00

Flag question

Mrs Molloy has been struggling with feelings of depression and self-blame. She's been talking about her childhood experiences in psychotherapy and described the way her father would make her follow very strict rules. On one occasion, she returned home only a few minutes after her curfew, and she was beaten by him with a belt. The parenting style she is describing is known as

Select one:

- Indulgent
- Secure
- Permissive
- Authoritarian
- Neglectful

Your answer is incorrect.

Explanation:

Rutter has clustered parenting into four general styles. The authoritarian style, characterised by strict, inflexible rules, can lead to low self-esteem, unhappiness, and social withdrawal. Authoritarian parents tend to be restrictive and cold, and their children tend to be withdrawn or conflicted. These children generally tend to grow up with little sense of agency or personal power, with a tendency towards depression, self-doubt and fear.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 898.

The correct answer is: Authoritarian

Question **45**

Not answered

Marked out of 1.00

Flag question

You are about to go to A&E to assess a patient who is feeling suicidal. Which demographic description has got the strongest association with completed suicide?

Select one:

- Elderly female
- Young female

Your answer is incorrect.

Explanation:

In the UK, men aged 45-49 years currently present the highest suicide rate. This is thought to be linked to middle-aged men being more likely affected in recent years by economic adversity, alcoholism, and social isolation. In most countries, rates of suicide are 2-3 times higher in males than in females, which could possibly be due to a male preference for higher lethality methods and the reluctance of males to seek help.

Ref: Correia R, Jackson D. Risk to self: identifying and managing risk of suicide and self-harm. BJPsych Advances. 2020:1-11.

Turecki G et al. Suicide and suicide risk. Nature reviews Disease primers. 2019 Oct 24;5(1):1-22.

The correct answer is: Middle-aged male

Question 46

Not answered

Marked out of 4.00

Flag question

Development of Adolescent Identity.
Match the stages of adolescent identity development based on Marcia's structural interview

A 21-year-old man after a lengthy period of exploration and self-searching, feels that he has a steady and stable sense of self. He finds himself committed to his work, with a clear sense of his goals and beliefs. — Choose...

A 17-year-old man is given the choice by his parents to stop attending church if he wishes to. He doesn't wish to make an exploration of his own, but continues to attend church with his family, and to conform to their way of practicing religion. — Choose...

An 18-year-old man is in the process of developing a sense of identity when he leaves home to join the military. Here he has the opportunity to learn and develop new skills, which later on in life are incorporated into his sense of self. — Choose...

A 17-year-old woman spends most of her time outdoors, in the company of a mixed group of young people. They spend time drinking alcohol, smoking cannabis; they have no plans for the future. — Choose...

Your answer is incorrect.

Marcia distinguished four identity statuses based on the amount of exploration and commitment the adolescent experiences or has experienced

3) Identity Foreclosure: Identity issues have not been considered seriously but future commitments have been made in spite of this. In this stage teenagers accept and adopt the traditional values and cultural norms of their parents or surrounding culture, rather than exploring and forming their own.

4) Identity Achievement: Various alternatives have been carefully considered and firm future commitments have been made. In this stage the individual is said to have solved their identity crisis and is able to make commitments to goals, values and beliefs.

(Ref:https://moodle.gllm.ac.uk/pluginfile.php/94785/mod_resource/content/1/Amanda_Tillson/Unit_4/Book_chapter_Adolesence_t

The correct answer is: A 21-year-old man after a lengthy period of exploration and self-searching, feels that he has a steady and stable sense of self. He finds himself committed to his work, with a clear sense of his goals and beliefs. → Identity Achievement, A 17-year-old man is given the choice by his parents to stop attending church if he wishes to. He doesn't wish to make an exploration of his own, but continues to attend church with his family, and to conform to their way of practicing religion. → Identity Foreclosure, An 18-year-old man is in the process of developing a sense of identity when he leaves home to join the military. Here he has the opportunity to learn and develop new skills, which later on in life are incorporated into his sense of self. → Identity Moratorium, A 17-year-old woman spends most of her time outdoors, in the company of a mixed group of young people. They spend time drinking alcohol, smoking cannabis; they have no plans for the future. → Identity Diffusion

Question 47

Not answered

Marked out of 5.00

Flag question

Personality disorders
What is the most likely diagnosis for each of the following patients?

Seema is a 33-year-old doctor who irritates her friends by speaking over them at dinner parties. Her friends complain that she 'always has to be the centre of attention.' — Choose...

Tod is a 25-year-old male who works in a family-run cornershop. He forgot to lock-up after work, which resulted in the store being burgled and a financial loss for the business. Tod appeared to demonstrate no concern for his actions. This is not the first — Choose...

Sabine is a 19-year-old fashion student. She holds the self-perception of being unique and superior to her peers. She told her supervisor that she felt she was exceptional and — Choose...

carries unusual ideas.

21-year-old history student, Raphael, met up with his dissertation supervisor, who was baffled to learn that Raphael did not appear to be remotely bothered with the feedback he received.

Choose...

Your answer is incorrect.

1) In Antisocial personality disorder, individuals show a lack of concern for others, irresponsibility, irritability, aggression, inability to maintain enduring relationships and disregard and violation of others' rights

2) In narcissistic personality disorder, grandiosity, lack of empathy and the need for admiration are common traits

3) In histrionic personality disorder, individuals show a high-degree of self-dramatization, shallow affect, egocentricity, craving attention, excitement and manipulative behaviour

4) Interpersonal discomfort with peculiar ideas, perceptions, appearance and behaviour are common traits in schizotypal personality disorder

5) Emotionally cold attitude, detachment, lack of interest in others, indifference to praise or criticism, excessive introspection and fantasy are common traits of schizoid personality disorder.

The correct answer is: Seema is a 33-year-old doctor who irritates her friends by speaking over them at dinner parties. Her friends complain that she 'always has to be the centre of attention.' → Histrionic personality disorder, Tod is a 25-year-old male who works in a family-run cornershop. He forgot to lock-up after work, which resulted in the store being burgled and a financial loss for the business. Tod appeared to demonstrate no concern for his actions. This is not the first → Antisocial personality disorder, Sabine is a 19-year-old fashion student. She holds the self-perception of being unique and superior to her peers. She told her supervisor that she felt she was exceptional and that was going to be the 'next Chanel', despite scoring a satisfactory grading → Narcissistic personality disorder, Jason is a 45-year-old courier. He has a poor rapport with his work colleagues and tends to withdraw socially. Despite working for his company for fifteen years, he has never attended an out-of-work social gathering. Jason's colleagues suspect that he is into some cult worship and carries unusual ideas. → Schizotypal personality disorder, 21-year-old history student, Raphael, met up with his dissertation supervisor, who was baffled to learn that Raphael did not appear to be remotely bothered with the feedback he received. → Schizoid personality disorder

Question 48

Not answered

Marked out of 5.00

Flag question

who cries during the nighttime and, as a result, has poor sleep. She complains of feeling excessively tired during the day.	Choose...
Shilpa, an 8-year-old girl, wakes up repeatedly in the night, fearful and feeling that she is going to die	Choose...
You meet Vanessa, a four-year-old girl, in your CAMHS outpatient clinic. Her mother tells you that Vanessa regularly wakes up distressed, complaining that she has had scary dreams.	Choose...
George is a 53-year-old male who was recently prescribed aripriprazole for his schizophrenia. His wife complains that he can't keep his legs still at night and that he tends to kick her in bed	Choose...

Your answer is incorrect.

1) Hypersomnolence: recurrent episodes of excessive daytime sleepiness. It can be caused by chronic sleep deprivation, whereby an individual fails to get sufficient sleep for at least three months. This case is an example of secondary hypersomnia.
2) Akathisia is a movement disorder characterized by restlessness and an urge to be in constant motion. Behaviourally, this manifests as rocking while sitting or standing, marching on the spot, and the crossing and uncrossing of legs whilst sitting down
3) During sleep terrors, there are high levels of autonomic arousal several times at night. It is a sleep disorder in which the individual experiences feelings of terror or dread.
4) This is a case of cataplexy, a condition whereby strong emotions or laughter cause a person to suffer sudden physical collapse. The patient remains conscious. Low orexin levels can be one of the common risk factors.
5) A nightmare is an unpleasant, 'bad dream' that causes a strong emotional response.

The correct answer is: Juliana is a 46-year-old lady who suffers from acute episodes of collapse on laughter. → Cataplexy, Susie is a 27-year-old nurse. She has a three-month-old son who cries during the nighttime and, as a result, has poor sleep. She complains of feeling excessively tired during the day. → Hypersomnolence, Shilpa, an 8-year-old girl, wakes up repeatedly in the night, fearful and feeling that she is going to die → Sleep terrors, You meet Vanessa, a four-year-old girl, in your CAMHS outpatient clinic. Her mother tells you that Vanessa regularly wakes up distressed, complaining that she has had scary dreams. → Nightmares, George is a 53-year-old male who was recently prescribed aripriprazole for his schizophrenia. His wife complains that he can't keep his legs still at night and that he tends to kick her in bed → Akathisia

Question 49

Not answered

Marked out of 4.00

Flag question

A severely depressed man is prescribed a medication, which leaves him with a dry mouth. A few days later he is found at home collapsed, having taken a significant overdose.	Choose...
A 50-year-old woman has been feeling low in mood, and finding it difficult to sleep. She is prescribed a medication by her GP, which is increased over a period of 4 weeks. Subsequently she has an ECG after reporting palpitations and dizziness.	Choose...
A middle-aged man admitted to the ward with chronic psychotic illness reports some chest discomfort. An ECG is performed.	Choose...

Your answer is incorrect.

1) Amitriptyline is an antidepressant medication that causes dry mouth and QTc prolongation leading to severe ventricular arrhythmias, collapse and death.

2) At higher doses, antidepressants may cause Torsades de Pointes.

3) Prolongation of QTc interval is a common side effect of antipsychotic medication.

4) This is clozapine-induced pericarditis.

The correct answer is: A young man with first episode psychosis is prescribed a medication and three months later has a routine ECG. → Prolongation of QTc interval, A severely depressed man is prescribed a medication, which leaves him with a dry mouth. A few days later he is found at home collapsed, having taken a significant overdose. → QTc prolongation of >510ms, A 50-year-old woman has been feeling low in mood, and finding it difficult to sleep. She is prescribed a medication by her GP, which is increased over a period of 4 weeks. Subsequently she has an ECG after reporting palpitations and dizziness. → Torsades de Pointes with use of antidepressant, A middle-aged man admitted to the ward with chronic psychotic illness reports some chest discomfort. An ECG is performed. → Saddle-shaped ST elevation

Finish review

Jump to...