Started on	Saturday, 23 July 2022, 11:15 AM
State	Finished
Completed on	Saturday, 23 July 2022, 11:16 AM
Time taken	13 secs
Grade	**0.00** out of 86.00 (**0**%)

Question **1**

Not answered

Marked out of 1.00

Flag question

You are reviewing cases in an addiction clinic. The note in the file says that your next patient has previously experienced 'formication'. What kind of hallucination is formication?

Select one:

- Visceral
- Thermal
- Haptic
- Kinaesthetic
- Hygric

Your answer is incorrect.

Explanation:

A haptic hallucination is a hallucination of touch. Formication is a form of tactile (or haptic) hallucination in which there is the sensation of numerous insects crawling over the surface of the body. This is not uncommon in acute organic states, including alcohol or drug withdrawal, particularly from cocaine. It is a form of superficial hallucination. The other forms of hallucinations listed above are also superficial hallucinations. Visceral hallucination is false perceptions of internal organs such as throbbing. Kinaesthetic hallucination is false perceptions of joint muscles (e.g. twisting or vibrating). Hygric hallucination is false perception of fluid. Thermal hallucination is false perception of hot and cold.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1140.

Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 111.

Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th Edition. 2019. p. 34.

The correct answer is: Haptic

Question **2**

Not answered

Marked out of 1.00

Donna and Malcom have had concerns about their son, Stuart, who is 6 years old and doesn't seem to be developing social skills in the same way as many of his peers. He dislikes being in groups, tends to spend a lot of time alone, immersed in quite a narrow repertoire of interests. If his usual routine is disrupted he can become very upset and difficult to comfort. Donna and Malcom have been searching the internet, reading about Autistic Spectrum

Flag question

Disorder, and how it is categorised in diagnostic manuals. Which one of the following statements is true regarding diagnostic categorisation in ICD and DSM?

Select one:

- ICD-11 has a category for pervasive developmental disorders
- DSM 5 has a category for pervasive developmental disorders
- DSM 5 categorises autistic spectrum disorder (ASD) under neurodevelopmental disorders
- DSM 5 has a category for overactive disorder associated with mental retardation
- DSM 5 includes language development as a core feature of autistic spectrum disorder (ASD)

Your answer is incorrect.

DSM 5 describes three 'severity levels' associated with the diagnosis of autistic spectrum disorder. Diagnostic criteria include persistent deficits in social communication and social interaction across multiple contexts; and restricted, repetitive patterns of behaviour, interests or activities, with symptoms present in the early developmental period and causing impairment in functioning. For persistent difficulties in the social use of verbal and nonverbal communication, a DSM-5 diagnosis of social communication disorder should be considered.

The correct answer is: DSM 5 categorises autistic spectrum disorder (ASD) under neurodevelopmental disorders

Question **3**

Not answered

Marked out of 1.00

Flag question

John Tulley is known to have epilepsy. On Saturday he attended a live rock concert with some of his friends. He had one pint of beer to drink. At one point during the concert, he fell to the ground and began to shake. The appearance was in keeping with a generalised epileptic seizure. Once the shaking had stopped, John began to regain consciousness. He sat up and was moving his arms, seemingly without purpose. His speech was not coherent, and he pursed his lips repeatedly. This lasted for approximately 10 minutes. John didn't recall this having happened when he was asked about it. What is the term used for this phase?

Select one:

- Twilight state
- Petit mal
- Delirium tremens
- Absence seizure
- Acute confusional state

Your answer is incorrect.

Explanation:

The term 'twilight state' describes conditions in which there is a restriction of the morbidly changed consciousness, a break in the continuity of consciousness, and relatively well-ordered behaviour. If one keeps strictly to these criteria, then the commonest twilight state is the result of epilepsy.

Ref: Casey P, Kelly B. Fish's clinical pathology: Signs and symptoms in psychiatry, 4th

Edition. 2019. p. 107.

The correct answer is: Twilight state

Question **4**

Not answered

Marked out of 1.00

Flag question

Hana Haq is a 42-year-old woman, who has presented to her GP with features of depression following the loss of her mother from cancer. For the past month she has been waking up a few hours earlier than usual. Her appetite has decreased (she has lost half a stone in weight) and she feels generally demotivated and hopeless, taking little pleasure from her usual activities. The GP fills out the Hamilton Depression Rating Scale (HAM-D) in order to ascertain the severity of her depression. Which of the following statements is true of the HAM-D?

Select one:

- It does not allow for estimation of severity
- It is scored out of a total of 8 points
- It excludes behavioural aspects of depression
- It has a section on weight loss
- It is self-administered

Your answer is incorrect.

Explanation:

The Hamilton Depression Rating Scale is a widely used scale for measuring depression, consisting of a list of 17 items, designed to be rated by a clinician on three-point and five-point scales. The symptoms are: depressed mood, guilt feelings, suicide, insomnia (initial or early), insomnia (middle), insomnia (terminal or late), disturbed work and activities, psychomotor retardation, psychomotor agitation, anxiety (psychological), anxiety (somatic), gastrointestinal symptoms, somatic symptoms, sexual dysfunction or menstrual disturbance, hypochondriasis, weight loss, and lack of insight. The maximum score is 52, and the following interpretation of scores is often used: 10-13 mild depression; 14-17 moderate depression; 18-30 marked depression; >30 severe depression.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: It has a section on weight loss

Question **5**

Not answered

Marked out of 1.00

Flag question

Which of the following statements is true about the diagnosis of delusional disorder?

Select one:

- ICD-11 requires the symptoms to be present for 6 months
- Delusional disorder is associated with mood symptoms
- General speech and behaviour are not typically affected
- ICD-11 classifies delusional disorder under neurotic disorders
- Delusions typically vary from day to day

Your answer is incorrect.

Explanation:

Delusional disorder is characterised by the development of a delusion or set of related delusions, typically persisting for at least 3 months and often much longer, in the absence of a depressive, manic, or mixed mood episode. The delusions are variable in content across individuals, but typically stable within individuals, although they may evolve over time. Other characteristic symptoms of schizophrenia (i.e. clear and persistent hallucinations, negative symptoms, disorganised thinking, or experiences of influence, passivity, or control) are not present, although various forms of perceptual disturbances (e.g. hallucinations, illusions, misidentification of persons) thematically related to the delusion are still consistent with the diagnosis. Apart from actions and attitudes directly related to the delusion or delusional system, affect, speech, and behaviour are typically unaffected.

Ref: International Statistical Classification of Diseases and Related Health Problems (11th ed; ICD-11; World Health Organization, 2020).

The correct answer is: General speech and behaviour are not typically affected

Question **6**

Not answered

Marked out of 1.00

Flag question

Bobby is a 22-year-old man, who has been brought to A&E with his mother, following their involvement in a serious road traffic accident. His mother was very badly injured, and sadly, whilst in the A&E, she dies. You are asked to go and speak to him. You find him to be in intense emotional distress: crying, sometimes sobbing, and unable to find coherent words to express himself. He says that it all feels unreal, and it's as if he's floating outside his body. What term is most appropriate to describe what is happening to Bobby?

Select one:

- Adjustment Reaction
- First episode psychosis
- Post-traumatic Stress disorder
- Moderate Depressive Episode
- Acute Stress Reaction

Your answer is incorrect.

Explanation:

Acute stress reaction refers to the development of transient emotional, somatic, cognitive, or behavioural symptoms as a result of exposure to an event or situation (either short- or long-lasting) of an extremely threatening or horrific nature. Symptoms may include autonomic signs of anxiety (e.g. tachycardia, sweating, flushing), being in a daze, confusion, sadness, anxiety, anger, despair, overactivity, inactivity, social withdrawal, or stupor. The response to the stressor is considered to be normal given the severity of the stressor (it is not considered a disorder in ICD-11), and usually begins to subside within a few days after the event or following removal from the threatening situation.

Ref: International Statistical Classification of Diseases and Related Health Problems (11th ed; ICD-11; World Health Organization, 2020).

The correct answer is: Acute Stress Reaction

Question **7**

Not answered

Priscilla is a 19-year-old woman whose mother has insisted she come to see you because she is worried about how thin she is. Priscilla tells you that she doesn't believe herself to be

Marked out of 1.00

Flag question

too thin. In fact, she thinks that she could do with losing more weight. She exercises four times a day, restricts her intake to a couple of apples and half a piece of toast a day. She has stopped menstruating. Her BMI is 15, and her T3 levels are normal. What's the most likely diagnosis?

Select one:

- Anorexia nervosa
- Bulimia nervosa
- Klein Levin syndrome
- Hyperthyroidism
- Mixed eating disorder

Your answer is incorrect.

Explanation:

Anorexia nervosa is characterised by significantly low body weight for the individual's height, age, and developmental stage that is not due to another health condition or the unavailability of food. A commonly used threshold is body mass index (BMI) less than 18.5 in adults. Low body weight is accompanied by a persistent pattern of behaviours to prevent restoration of normal weight, which may include behaviours aimed at reducing energy intake (restricted eating) purging behaviours (e.g. self-induced vomiting, misuse of laxatives), and behaviours aimed at increasing energy expenditure (e.g. excessive exercise), typically associated with a fear of weight gain. Low body weight or shape is central to the person's self-evaluation or is inaccurately perceived to be normal or even excessive. T3 levels are not part of the diagnostic criteria; commonly, T3 levels in anorexia nervosa are low, but not in every case.

Ref: International Statistical Classification of Diseases and Related Health Problems (11th ed; ICD-11; World Health Organization, 2020).

The correct answer is: Anorexia nervosa

Question **8**

Not answered

Marked out of 1.00

Flag question

Mr. Phipps is 69 years old, with a history of depression, hypertension and hypercholesterolaemia. Last week he suffered a stroke affecting his non-dominant occipitoparietal lobe. You draw Mr. Phipps' attention to objects on a table (a watch, a spoon, and a plant) and ask him to name them. He tells you he can see a spoon, but nothing else. When you redirect his attention to the plant, he says that now he does see the plant, but that he no longer sees the spoon. What term best describes this presentation?

Select one:

- Gerstmann syndrome
- Dyslexia
- Constructional apraxia
- Simultanagnosia
- Anosognosia

Your answer is incorrect.

Simultanagnosia (also called simultagnosia) is a rare neurological disorder characterized by

the inability of an individual to perceive more than one single object at a time, and is associated with occipito-parietal lesions.

The correct answer is: Simultanagnosia

Question **9**

Not answered

Marked out of 1.00

Flag question

Miss Hunt is known to the local mental health team, having been diagnosed with a mental illness by your predecessor. The GP informs you that an ECG has been performed and shows a prolonged QT interval. You do not have the past notes available to you at this time, however you know that the following mental health condition is most associated with prolonged QT interval:

Select one:

- Depression
- Anxiety
- Post-Traumatic Stress Disorder
- Schizophrenia
- Anorexia Nervosa

Your answer is incorrect.

The starvation effects of anorexia nervosa have consequences on cardiac function and associated with prolonged QT interval. (Ref: British Heart Journal 1994; 72:69-73)

The correct answer is: Anorexia Nervosa

Question **10**

Not answered

Marked out of 1.00

Flag question

Mr. Nolan is a 35-year-old man who has suffered right cerebral hemisphere damage following a motorcycle accident. Which of the following symptoms is he most likely to experience?

Select one:

- Right upper limb hemiparesis
- Mania
- Right lower limb hemiparesis
- Depression
- Pathological crying

Your answer is incorrect.

Right hemisphere damage is associated with numerous deficits, including a flat affect and limited emotional expression. They can also experience deficits in attention, perception, learning, memory, recognition and understanding of para-verbal communication. They can demonstrate left-sided neglect and hemiparesis.

The correct answer is: Depression

Question **11**

Not answered

Marked out of 1.00

Flag question

Mrs Oona is a 42-year-old woman who attends her GP describing a sense of feeling her right arm being pulled and distorted when waking from sleep. Which of the following is the name for this phenomenon?

Select one:

- Delusion of control
- Kinaesthetic hallucination
- Visceral hallucination
- Paraschemazia
- Illusion

Your answer is incorrect.

This represents a kinaesthetic hallucination as it involves a sensory disturbance concerning the limbs. A visceral hallucination would relate to feelings perceived as arising in the viscera and internal organs. Paraschemazia is characterised by a distortion of a patient's body image. It can be caused by hallucinogenic drugs such as LSD and mescalin, epileptic auras, and sometimes migraines. A delusion of control is an abnormality of thought in which a patient believes that their thoughts can be controlled by an external force. An illusion is a sensory distortion which clear once an individual attend to it, in the case of affect-related or completion illusions, or ignores it, in the case of pareidolia.

The correct answer is: Kinaesthetic hallucination

Question **12**

Not answered

Marked out of 1.00

Flag question

Your outpatient mental health service is considering making use of a psychiatric tool in order to assess presence and severity of mental distress and illness. Someone suggests using the Brief Psychiatric Rating Scale (BPRS) as a global assessment of symptoms. Which of the following is true of the BPRS?

Select one:

- It covers 7 areas of concern
- All items are based on the individual's self-report
- It includes a section on eating disorders
- Items are rated 1-10 in terms of severity
- It includes a section on somatic concern

Your answer is incorrect.

The BPRS covers 24 sections/areas of concern, rated in severity 1 to 7. The first 14 items are based on the individual's self-report, with the remainder on observed behaviour and speech.

The correct answer is: It includes a section on somatic concern

Question **13**

Not answered

Marked out of

Which of the following describes the EEG findings of a patient with delirium?

Select one:

1.00

Flag question

- Poly-spike discharge
- 3Hz spike wave complex
- Diffuse slowing
- Localized rhythmic 5-7 Hz
- Generalized spike wave discharge

Your answer is incorrect.

Diffuse slowing on an EEG is associated with delirium. The other findings are described below: Generalised spike wave discharge is seen in tonic clonic epilepsy. 3 Hz spike wave complex is seen in absent seizures. Poly spike discharge is seen in myoclonic epilepsy. Localised rhythmic 5-7 Hz is seen in temporal lobe epilepsy.

The correct answer is: Diffuse slowing

Question **14**

Not answered

Marked out of 1.00

Flag question

Shalinee, a 30-year-old florist, has a history of epilepsy for which she takes carbamazepine. She is planning to start a family with her husband, David and seeks your advice on medication during pregnancy. Which of the following pharmacokinetic changes occurs during pregnancy?

Select one:

- Reduced absorption from intramuscular site of injection
- Increased renal elimination
- Reduced volume of distribution
- Increased plasma protein binding
- Reduced rate of metabolism

Your answer is incorrect.

Gastric emptying and small intestine motility are reduced in pregnancy due to elevation of progesterone, which tends to reduce absorption via the oral route. Total body water increases by up to 8 litres, creating a larger space within which hydrophilic drugs may distribute. Total plasma concentration of albumin-bound drugs decreases as a result of haemodilution and the subsequent fall in plasma albumin concentration and reduced plasma protein binding. Some enzymes of the hepatic cytochrome P-450 system are induced by oestrogen/ progesterone, resulting in a higher rate of metabolism (and hence elimination) of drugs. Renal blood flow is increased by 60-80% during pregnancy, and glomerular filtration rate rises by 50%, leading to enhanced elimination of drugs (Dawes 2001)

The correct answer is: Increased renal elimination

Question **15**

Not answered

Marked out of 1.00

Flag

There are numerous specific sub-types of hallucination. Which of the following would represent an extracampine hallucination?

Select one:

- A person hearing commentary voices

question

- A person complaining being able to hear people saying bad things about him from another city
- A person feeling that other people can read his mind
- A person hearing someone calling his name while falling asleep
- A person feeling cold while watching a documentary

Your answer is incorrect.

Extracampine hallucinations involve an individual perceiving a voice or auditory stimulus originating from a place which would be out of earshot and thus impossible to actually hear. The other examples describe reflex hallucinations, hypnagogic hallucinations, thought broadcast and running commentary respectively.

The correct answer is: A person complaining being able to hear people saying bad things about him from another city

Question **16**

Not answered

Marked out of 1.00

Flag question

May is a 78-year-old woman whose husband passed away three weeks ago, following a long illness. You visit her at home and she talks to you about her experience of loss. Which feature, following a loss, is usually the most salient?

Select one:

- Depression
- Anger
- Disbelief
- Numbness
- Yearning

Your answer is incorrect.

According to updated research published in the Journal of the American Medical Association, conducted by the Yale School of Medicine (JAMA 297:Feb 2001), the most salient emotion following loss is yearning. In the study, Maciejewski said yearning is the defining feature of grief and is an emotion that most clearly reflects the absence of the deceased. "Yearning is a longing for reunion with the deceased loved one, heartache about an inability to reconnect with this person. Individuals may cognitively accept the death of a loved one, but they may still pine for them and experience pangs of grief (i.e. yearning)."

The correct answer is: Yearning

Question **17**

Not answered

Marked out of 1.00

Flag question

Which of the following is not a true statement of Charles Bonnet Syndrome (CBS)?

Select one:

- A symptom of CBS is visual hallucinations of simple repeated patterns
- CBS is a form of dementia
- CBS can last from a few days to years
-

A symptom of CBS is complex visual hallucinations

CBS can be caused by cataracts

Your answer is incorrect.

Charles Bonnet Syndrome (CBS) is not a mental illness, nor a form of dementia. It is caused by reducing vision in older age such as cataracts or macular degeneration. The person experiences visual hallucinations out of the blue that can either be simple vivid patterns or more complex visual hallucinations.

The correct answer is: CBS is a form of dementia

Question **18**

Not answered

Marked out of 1.00

Flag question

Which of the following is not a component of schizotypal personality disorder?

Select one:

- Paranoid ideation
- Auditory hallucinations
- Lacking close friends
- Eccentric dress sense
- Illusions

Your answer is incorrect.

Auditory hallucinations are not associated with schizotypal personality disorder, and more associated with an acute psychotic episode.

The correct answer is: Auditory hallucinations

Question **19**

Not answered

Marked out of 1.00

Flag question

An EEG is performed on a patient of yours who was in REM sleep. Which of the following is NOT a feature of REM sleep in an EEG?

Select one:

- Theta waves
- High-voltage activity
- Slow alpha waves
- Saw-tooth waves
- Similar activity to an awake state

Your answer is incorrect.

REM sleep accounts for 25% of total sleep time in adults and is characterized by high levels of brain activity and physiological activity similar to those seen in wakefulness. The EEG will show low-voltage and mixed frequency (theta and slow alpha) activity. Saw-tooth waves are also seen.

The correct answer is: High-voltage activity

Question **20**

Not answered

Marked out of 1.00

Flag question

Mr. Lyons is a man who has been admitted to hospital with a new onset psychotic illness. He is started on an antipsychotic but then develops confusion, muscular stiffness, marked sweating and fever. Which of the following factors increases the risk of him developing this condition?

Select one:

- Concurrent clonazepam treatment
- Iron deficiency
- Obesity
- Hypothyroidism
- Older age

Your answer is incorrect.

A variety of other risk factors have emerged from epidemiological and case studies of NMS which include dehydration, physical exhaustion, exposure to heat, hyponatremia, iron deficiency, malnutrition, trauma, thyrotoxicosis, alcohol, psychoactive substances, and presence of a structural or functional brain disorder such as encephalitis or dementia.

The correct answer is: Iron deficiency

Question **21**

Not answered

Marked out of 1.00

Flag question

Mr. Parson is a man who is referred to a psychiatric clinic with concerns around his cognitive function. He is sent for an MRI, which demonstrates the 'pulvinar sign' in his thalamus. Which of the following conditions is he most likely to be suffering from?

Select one:

- Alzheimer's disease
- Lewy body dementia
- Fronto-temporal dementia
- Variant CJD
- Sporadic CJD

Your answer is incorrect.

Variant CJD is associated, in 75% of cases, with the pulvinar sign. This is bilateral high signal intensities on axial fluid attenuated inversion recovery (FLAIR) MRI. Posterior thalami involvement on sagittal T2 sequences is also common.

The correct answer is: Variant CJD

Question **22**

Not answered

Mrs. Richards is a 32-year-old woman who has been referred to a neurologist for investigation of possible narcolepsy. Which of the following investigations would be most

Marked out of 1.00

Flag question

likely to be used?

Select one:

- CT
- EEG
- Multiple sleep latency test
- MRI
- DAT scan

Your answer is incorrect.

The tests that are commonly used in diagnosing narcolepsy are the polysomnogram, the multiple sleep latency test (MSLT), and administration of the Epworth Sleepiness Scale.

The correct answer is: Multiple sleep latency test

Question **23**

Not answered

Marked out of 1.00

Flag question

Matilde is an 87-year-old female who is awaiting discharge from the inpatient unit. You review her medically and notice that her answers, although well formed, are unrelated to your questions. Her medical history includes hypertension and diabetes. Two years ago she suffered from a TIA. Comprehension of speech is associated with which part of the cerebral cortex?

Select one:

- Inferior temporal gyrus
- Medial frontal gyrus
- Posterior inferior frontal gyrus
- Fusiform gyrus
- Posterior superior temporal gyrus

Your answer is incorrect.

Wernicke's area is involved in the comprehension of written and spoken language. It is located in the posterior superior temporal gyrus of the cerebral cortex. Damage to this area, for example, through infarction, results in a receptive dysphasia whereby patients experience difficulty in understanding language. The production of speech is not impaired and so the patient is able to speak normally, although the content of their speech is unrelated to the topic of conversation.

The correct answer is: Posterior superior temporal gyrus

Question **24**

Not answered

Marked out of 1.00

Flag question

In your CAMHS outpatient clinic, you review Bobby, an 8-year-old boy recently diagnosed with ADHD. You meet with his parents to discuss your plans to commence methylphenidate (Ritalin). They are concerned about possible side effects and wish to know more about these. Which of the following is the most common side effect of methylphenidate?

Select one:

- Constipation
- Confusion
- Visual disturbance
- Anorexia
- Insomnia

Your answer is incorrect.

Methylphenidate is recommended for use as part of a comprehensive treatment programme for children with a diagnosis of severe attention deficit hyperactivity disorder (ADHD). Common side effects include reduced appetite and insomnia. Less common side effects include tics or twitches, drowsiness or dizziness, headache, abdominal pain, nausea.

The correct answer is: Insomnia

Question **25**

Not answered

Marked out of 1.00

Flag question

Anna attends a rave where she takes an unknown substance. She later describes the feeling of her limbs being distorted. Which of the following describes this phenomenon?

Select one:

- Hyposchemazia
- Aschemazia
- Paraschemazia
- Asomatognosia
- Anosognosia

Your answer is incorrect.

Paraschemazia or body image distortion is the phenomenon of experiencing limbs or other body parts are being distorted; it is commonly seen whilst under the influence of hallucinogens. Hyposchemazia and aschemazia describe the phenomenon of experiencing limbs that are diminished or absent respectively. Asomatognosia is the loss of awareness or recognition of a part of the body. Anosognosia is unawareness of neurological impairment.

The correct answer is: Paraschemazia

Question **26**

Not answered

Marked out of 1.00

Flag question

Howard is a young man admitted to the ward. He has a diagnosis of schizophrenia and whilst assessing him, he sweeps your notes off the table onto the floor. On questioning why he did that, he states that he didn't do it but that something made him. Which of the following explains the mechanism behind this experience?

Select one:

- Dyskinesia
- Altered sensorimotor feedback
- Phantom limb
- Loss of proprioception

- Anosognosia

Your answer is incorrect.

The patient in the vignette above is describing delusions of control. The phenomenon is explained by altered somatosensory feedback where there is a mismatch between the predicted motor and sensory feedback and the actual feedback. (Ref: M Wolpert, D & Ghahramani, Zoubin & Jordan, Michael. (1995). An Internal Model for Sensorimotor Integration. Science (New York, N.Y.). 269. 1880-2. 10.1126/science.7569931)

The correct answer is: Altered sensorimotor feedback

Question **27**

Not answered

Marked out of 1.00

Flag question

There are various factors, which influence group dynamics and impact upon their effectiveness. Which of the following factors can reduce the cohesiveness of a group?

Select one:

- Large group size
- Role ambiguity
- Open to new members
- Member diversity
- Members function independently

Your answer is incorrect.

Role ambiguity refers to the lack of clear definition of what is expected of an individual, with respect to the goals of their team. The more uncertainty there is about the tasks, goals and levels of performance, the less satisfied workers tend to be. But in the presence of task and social cohesion job satisfaction improves greatly.

Ref: Urien B, Osca A, García-Salmones L. Role ambiguity, group cohesion and job satisfaction. Revista Latinoamericana de Psicología. 2017 May 1;49(2):137-45.

The correct answer is: Role ambiguity

Question **28**

Not answered

Marked out of 1.00

Flag question

Attention results in the conversion of:

Select one:

- Retrieval of long term memory
- Iconic memory to echoic memory
- Short term memory into long term memory
- Working memory into short term memory
- Sensory memory into short term memory

Your answer is incorrect.

Sensory receptors in the brain are responsible for receiving sensory information which they

store for mere seconds. The brain filters this multifaceted sensory information from multiple sensory organs. When conscious attention is paid, this certain sensory information is selected and converted into short-term memory which enables us to remember it.

The correct answer is: Sensory memory into short term memory

Question **29**

Not answered

Marked out of 1.00

Flag question

Working memory can be tested at bedside using which of the following?

Select one:

- Stroop Test
- Wisconsin Card Sorting Test
- Digit span
- Cognitive Estimation Test
- Clock-drawing Test

Your answer is incorrect.

Explanation

Digit span tests working memory by asking the participant to recall sequences of numbers of increasing lengths in a forward or backward manner. It requires patients to repeat longer strings of digits increasingly as a way of assessing the ability to process relatively simple information.

Cognitive estimation is used to test problem solving. Wisconsin card sorting test is used to test set shifting. Stroop test is used to test selective attention. Clock drawing test is used to test general cognitive ability in dementia.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 54.

The correct answer is: Digit span

Question **30**

Not answered

Marked out of 1.00

Flag question

Which of the following strategies are employed in order to reduce groupthink?

Select one:

- Focusing on individual successes within the group
- Not revealing the roles of individuals within the group
- Preventing discussion in order to save time
- Allowing decisions to be made without criticism
- Appointing a 'devil's advocate'

Your answer is incorrect.

Explanation:

Groupthink is a collective pattern of defensive avoidance, characteristic of group decision making in organisations in which group members develop rationalisations supporting shared illusions of their own infallibility and invulnerability with the organisation. Characteristic

features include: (1) An illusion of invulnerability; (2) Collective efforts to rationalise in order to discount warnings; (3) An unquestioned belief in the group's inherent morality; (4) Stereotyped views of rivals and enemies; (5) Direct pressure on any member who dissents from the consensus view; (6) Self-censorship of disagreements with the apparent group consensus; (7) A shared illusion of unanimity; and (8) The emergence of self-appointed protectors of the group from adverse information that might undermine the existing complacency. Appointing a 'devil's advocate' is one strategy that can be employed to reduce groupthink.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: Appointing a 'devil's advocate'

Question **31**

Not answered

Marked out of 1.00

Flag question

Anthony Andrews is a 7-year-old boy who has been displaying some challenging behaviours at school. His teacher produces a star chart with a star sticker being awarded to Anthony for good behaviour. Which of the following forms of conditioning would this best fit with?

Select one:

- Observational learning
- Operant conditioning
- Social conditioning
- Classical conditioning
- Imprinting

Your answer is incorrect.

Explanation:

Behaviour modification is based on operant conditioning. Forms of behavioural modification include the use of secondary reinforcement, such as 'token economy', in which socially desirable/acceptable behaviours are rewarded with tokens that can be exchanged for other material items or privileges, or 'star charts' where children's good behaviour is rewarded when a certain level is achieved.

In classical conditioning, stimuli that signal significant events produce reflexive behaviour. Observational learning is learning that occurs through observing the behaviour of others. Social conditioning is the sociological process of training individuals in a society to respond in a manner generally approved by the society in general and peer groups within society. Imprinting describes the phase-sensitive period of a young animal attaching to an external stimulus, usually its mother.

Ref: Semple D, Smyth R. Oxford handbook of psychiatry. Oxford university press; 2019 Jul 30. p. 908.

The correct answer is: Operant conditioning

Question **32**

Not answered

Marked out of 1.00

Flag

You are the junior doctor at a unit for severe personality disordered patients, otherwise known as a therapeutic community. You observe group therapy and attempt to identify group dynamics based on what you have read in theory. Which of the following best describes 'group cohesion'?

Select one:

question

- The quality of the relationship that develops between each individual member and the therapist
- The process by which mere expression of ideas and conflicts is accompanied by an emotional response which produces a sense of relief
- When the group becomes self-evolving and is able to work through conflicts
- When an individual incorporates the characteristics and qualities of the group
- When the group works together towards a common goal

Your answer is incorrect.

Explanation:

Therapeutic factors in group psychotherapy:

Therapeutic factor	Definition
Abreaction	Repressed material, particularly a painful experience, is brought back to consciousness
Acceptance	Feeling of being accepted by other members of the group
Altruism	Act of one member helping another; putting another's needs before one's own
Catharsis	Expression of ideas/thoughts/suppressed material that is emotional and produces relief
Cohesion	Sense that the group is working together toward a common goal
Consensual validation	Confirmation of reality by comparing own conceptualisations with other group members
Contagion	Expression of emotion by a member stimulates awareness of similar emotion in another
Corrective familial experience	Group recreates the family of origin to work through original conflicts psychologically
Empathy	Capacity of group members to put themselves in psychological frame of another member
Identification	Unconscious defense mechanism; characteristics of another person are incorporated
Imitation	Conscious emulation/modeling of one's behaviour after another; 'role modeling'
Insight	Conscious awareness/understanding of own psychodynamics and maladaptive behaviour
Inspiration	Process of imparting a sense of optimism to group members; 'instillation of hope'
Interaction	Free and open exchange of ideas and feelings among group members
Interpretation	Group leader formulates the meaning of a patient's resistance, defense, and symbols
Learning	Acquiring knowledge about new areas, such as social skills; receiving advice and obtaining guidance
Reality Testing	Ability of the person to evaluate objectively the world outside the self
Transference	Projection of feelings, thoughts, and wishes onto the therapist
Universalisation	Awareness of patients that they are not alone in having problems; 'self-disclosure'
Ventilation	Expression of suppressed feelings, ideas, or events to other group members
	Adapted from Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition, 2022. p. 757.

SPMM COURSE

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 757.

The correct answer is: When the group works together towards a common goal

Question **33**

Not answered

Marked out of 1.00

Flag question

Cognitive Estimates Test is used to test the function of which of the following regions of the brain?

Select one:

- Occipital Cortex
- Orbitofrontal Cortex
- Olfactory Cortex
- Dorsolateral prefrontal cortex
- Anterior cingulate cortex

Your answer is incorrect.

The Cognitive Estimation Test (CET) is among a number of tests examining executive function. The dorsolateral prefrontal cortex (DLPFC), anterior cingulate cortex (ACC) and orbitofrontal cortex (OFC) all make up the prefrontal region of the frontal lobe and are all associated with executive functioning but the DLPFC is associated with, among others, problem-solving and abstract reasoning required in the CET. The ACC is associated with decision-making, motivation and response inhibition and the OFC is associated with impulse control and socially appropriate behaviours.

The correct answer is: Dorsolateral prefrontal cortex

Question **34**

Not answered

Marked out of 1.00

Flag question

Amy attends outpatient clinic complaining of discharge from her nipples. Which of the following medications is least associated with this side effect?


Select one:

- Haloperidol
- Amisulpride
- Clozapine
- Risperidone
- Sulpiride

Your answer is incorrect.

Explanation:

A side effect of antipsychotics can be hyperprolactinaemia. However, clozapine and aripiprazole are rarely associated with this side effect, and aripiprazole has been shown at low doses to reduce prolactin levels. This effect is attributed to the D2 receptor partial agonism, which may mitigate the effect of other antipsychotic medications on the pituitary gland.



Ref: Taylor DM, Barnes TRE, Young AH. The Maudsley Prescribing Guidelines in Psychiatry,

14th Edition. 2021. p. 168.

The correct answer is: Clozapine

Question **35**

Not answered

Marked out of 1.00

Flag question

You and your colleagues are discussing various aspects of the nature of psychiatry, its history and current predicaments. One of your colleagues begins to reference a well-known book by Thomas Szasz. In his seminal book, 'The Myth of Mental Illness', Thomas Szasz makes which of the following claims?

Select one:

- Symptoms are explanations for mental illness
- There is no such thing as mental illness
- All problems in living can be explained by psychochemical processes
- The body and mind are separate
- Mental illness is a sign of brain disease

Your answer is incorrect.

Explanation:

There were three major themes in Szasz's writings: 1) His contention that there is no such thing as mental illness; 2) His contention that individual responsibility is never compromised in those suffering from what is generally considered as mental illness; 3) His perennial interest in calling attention to the political nature of psychiatric diagnosis.

Ref: Benning TB. No such thing as mental illness? Critical reflections on the major ideas and legacy of Thomas Szasz. BJPsych bulletin. 2016 Dec;40(6):292-5.

The correct answer is: There is no such thing as mental illness

Question **36**

Not answered

Marked out of 1.00

Flag question

Which of the following is a component of Kohlberg's conventional stage of morality?

Select one:

- Authority
- Punishment
- Reward
- Universal ethics
- Social contract

Your answer is incorrect.

Explanation:

Lawrence Kohlberg integrated Jean Piaget's concepts and described multiple stages of moral development within three significant levels of morality. The first level is pre-conventional morality, in which punishment and obedience are the determining factors. The second level is morality of conventional role-conformity, in which children try to conform to gain approval and maintain good relationships with others and are also guided by rigid codes of 'law and order'

(authority orientation). The third and highest level is the morality of self-accepted moral principles (post-conventional), in which children voluntarily comply with rules based on a concept of universal ethical principles and make exceptions to rules in certain circumstances.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 902.

Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: Authority

Question **37**

Not answered

Marked out of 1.00

Flag question

According to Erikson's stages of psychosocial development, at which stage would a four year old be?

Select one:

- Ego identity vs role confusion
- Initiative vs guilt
- Autonomy vs shame
- Trust vs mistrust
- Industry vs inferiority

Your answer is incorrect.

Explanation:

Erik Erikson created an original and highly influential theory of psychological development and crisis occurring in periods that extended across the entire life cycle. Stage 3 takes place from about 3 years to about 5 years. Its focus is initiative versus guilt. The child's increasing mastery of locomotor and language skills expands its participation in the outside world and stimulates omnipotent fantasies of broader exploration and conquest. Guilt over the drive for conquest and anxiety over the anticipated punishment are both assuaged in the child through repression of the forbidden wishes and development of a superego to regulate its initiative. Success in this stage will result in development of the basic virtue of purpose.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. pp. 1073-1074.

The correct answer is: Initiative vs guilt

Question **38**

Not answered

Marked out of 1.00

Flag question

Mrs. Kean and her 16-year-old son Henry have been having some disagreements about his future plans. Mrs. Kean has decided which universities Henry will apply for. He protests, saying he wants to make his own decision, but Mrs. Kean replies saying that the choice is not his, the decision will be made for him by her and her husband and there's nothing he can do about it. Henry reluctantly concedes. What parenting style is Mrs. Kean exhibiting?

Select one:

- Neglectful
- Authoritative
- Authoritarian
- Permissive

- Boundaried

Your answer is incorrect.

Explanation:

Rutter has clustered parenting into four general styles. The authoritarian style, characterised by strict, inflexible rules, can lead to low self-esteem, unhappiness, and social withdrawal. Authoritarian parents tend to be restrictive and cold, and their children tend to be withdrawn or conflicted.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 898.

The correct answer is: Authoritarian

Question **39**

Not answered

Marked out of 1.00

Flag question

Melissa and her partner Jakob have recently had a baby. Jakob is looking forward to charting their son's development, and wants to draw up a time-line so that he can anticipate what new skills his son may be able to master. He is anxious to notice any clues that may suggest his son is not developing as expected. Which of the following scenarios suggests impaired development?

Select one:

- Able to walk at 13 months
- Plays with a ball at 16 months
- Holding head up at 3 months
- Able to roll at 9 months
- 2-3 words at 24 months

Your answer is incorrect.

Explanation:

By age 18 months, a child is generally using 20 single words. They will understand up to 150 words. By 24 months, their vocabulary will increase to up to 270 words.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 889.

The correct answer is: 2-3 words at 24 months

Question **40**

Not answered

Marked out of 1.00

Flag question

What type of attachment is demonstrated by a child who seems unaffected by the mother leaving and carries on playing and also unaffected by the mother's return?

Select one:

- Anxious-avoidant attachment
- Anxious-ambivalent attachment
- Disorganised attachment
- Anxious-resistant attachment


- Secure attachment

Your answer is incorrect.

Explanation:

Mary Ainsworth devised the Strange Situation procedure with 12- to 18-month infants, which categorised infant-parent relationships. The procedure involves separations and reunions with caregivers and observes response. In anxious-avoidant attachment, the child seems unconcerned at separation or reunion. It is associated with unresponsive parenting. Mary Ainsworth noted that this type of attachment often occurred when the child's needs were repeatedly not met and allows the child to maintain some proximity to the parent without experiencing maternal rejection. Note that the anxious-avoidant, anxious-resistant and disorganised categories are also described as insecure-avoidant, insecure-resistant (or ambivalent) and insecure-disorganised. Ainsworth preferred classifications using alphabetical notations (A, B, C). The disorganised category is attributed to Ainsworth's student, Mary Main.

For a short illustrative video see



Ref: Duschinsky R. Mary Ainsworth and the Strange Situation Procedure. In: *Cornerstones of Attachment Research*. New York, NY: Oxford University Press; 2020:109-210. https://www.oxfordclinicalpsych.com/view/10.1093/med-psych/9780198842064.001.0001/med-9780198842064-chapter-2. Accessed March 4, 2022.

The correct answer is: Anxious-avoidant attachment

Question **41**

Not answered

Marked out of 1.00

Flag question

Nina and her mother are spending the afternoon together. Nina's mother has been talking about families. At one point in the conversation Nina asks her mother 'what would happen if I have a baby brother?' What developmental stage, according to Piaget, is Nina most likely to be at?

Select one:

- Formal Operational
- Conventional
- Sensorimotor

- Preoperational
- Concrete Operational

Your answer is incorrect.

Explanation:

According to Piaget, the stage of formal operations lasts from about 11 years of age through the end of adolescence. The stage is so named because young persons' thinking operates in a formal, highly logical, systematic, and symbolic manner. It is characterised by the ability to think abstractly, to reason deductively, and to define concepts, and also by the emergence of skills for dealing with permutations and combinations. Adolescents attempt to deal with all possible relations and hypotheses to explain data and events during this stage. They are able to think abstractly or hypothetically, which is demonstrated by Nina's question.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1009.

The correct answer is: Formal Operational

Question **42**

Not answered

Marked out of 1.00

Flag question

Kevin is a young boy who is getting into arguments with other children at school and is occasionally pushing them over. His parents and teachers have attempted to reason with him but this has not been successful. His behaviour eventually improves when he is told that his favourite toy will be removed as long as he continues to push others. Which of the following stages of Kohlberg's moral development would best fit this scenario?

Select one:

- Obedience and punishment orientation
- Self-interest orientation
- Authority and social-order maintaining orientation
- Interpersonal accord and conformity
- Social contract orientation

Your answer is incorrect.

Explanation:

Lawrence Kohlberg integrated Jean Piaget's concepts and described multiple levels of moral development within three significant levels of morality. The first level is pre-conventional morality, in which punishment and obedience are the determining factors. The second level is morality of conventional role-conformity, in which children try to conform to gain approval and maintain good relationships with others and are also guided by rigid codes of 'law and order' (authority orientation). The third and highest level is the morality of self-accepted moral principles (post-conventional), in which children voluntarily comply with rules based on a concept of universal ethical principles and make exceptions to rules in certain circumstances.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 902.

Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: Obedience and punishment orientation

Question **43**

Not answered

Marked out of 1.00

Flag question

In the development of vision, which of the following can be seen occurring as early as the first day after birth?

Select one:

- Pupil reaction to light
- Depth perception
- Accommodation
- Convergence
- Colour vision

Your answer is incorrect.

The pupillary light reflex (PLR) whereby the diameter of the pupil is controlled according to the intensity of light occurs from birth. Depth perception, colour vision and accommodation occur at three to five months. Convergence occurs at five to seven months.

The correct answer is: Pupil reaction to light

Question **44**

Not answered

Marked out of 1.00

Flag question

Anna is 8 years old. You are conducting some exercises to determine what stage of development she appears to be at, according to Piaget's stages of development. You present her with two glasses, both of the same shape, and both holding the same amount of orange juice. You then obtain two other glasses of different shapes: one is tall and thin, another short and broad. You pour the orange juice from one of the original glasses into the tall glass, and the juice from the other original glass into the shorter glass. You now ask Anna to tell you which glass holds the most juice. What answer indicates she has reached the concrete operational stage of development?

Select one:

- 'Both glasses have the same amount of orange juice'
- 'The short glass has the most orange juice'
- 'They each have more orange juice than before'
- 'Glasses can be half empty or half full'
- 'The tall glass has the most orange juice'

Your answer is incorrect.

Explanation:

According to Piaget, the stage of concrete operations lasts from about 7 to 11 years of age. In this period, children operate and act on the concrete, real, and perceivable world of objects and events. Egocentric thought is replaced by operational thought, which involves dealing with a wide array of information outside the child. Children can now see things from someone else's perspective. Children also understand the concept of **conservation**, which is the ability to recognise that, although the shape of objects may change, the objects still maintain or conserve other characteristics that enable them to be recognised as the same. The ability of children to understand quantity is one of Piaget's most important cognitive developmental theories. Measures of quantity include measures of substance, length, number, liquids, and

area.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. pp. 1008-1009.

The correct answer is: 'Both glasses have the same amount of orange juice'

Question **45**

Not answered

Marked out of 1.00

Flag question

Which of the following of the Kohlberg's stages of moral development are paired correctly?

Select one:

- Post-conventional – authority
- Pre-conventional - punishment
- Pre-conventional – social contract
- Post-conventional - conformity
- Conventional – universal principles

Your answer is incorrect.

Explanation:

Lawrence Kohlberg integrated Jean Piaget's concepts and described multiple stages of moral development within three significant levels of morality. The first level is **pre-conventional** morality, in which **punishment** and obedience are the determining factors. The second level is morality of conventional role-conformity, in which children try to conform to gain approval and maintain good relationships with others and are also guided by rigid codes of 'law and order' (authority orientation). The third and highest level is the morality of self-accepted moral principles (post-conventional), in which children voluntarily comply with rules based on a concept of universal ethical principles and make exceptions to rules in certain circumstances.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 902.

Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: Pre-conventional - punishment

Question **46**

Not answered

Marked out of 1.00

Flag question

The Oedipus complex is associated with which stage of psychosexual development?

Select one:

- Oral
- Anal
- Latency
- Phallic
- Genital

Your answer is incorrect.

Explanation:

The phallic stage begins sometime during year 3 and continues until approximately the end of year 5. Threats of castration and the related anxiety are connected with guilt over masturbation and oedipal wishes. During this phase oedipal involvement and conflict are established and consolidated.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1062.

The correct answer is: Phallic

Question **47**

Not answered

Marked out of 1.00

Flag question

Traditionally, the view has been that boys like to play with cars and girls like to play with dolls. Which of the following is this an example of?

Select one:

- Internalisation
- Socialisation
- Stereotyping
- Indoctrination
- Normalisation

Your answer is incorrect.

Explanation:

Socialisation is the process, beginning in infancy, whereby one acquires the attitudes, values, beliefs, habits, behaviour patterns, and accumulated knowledge of one's society, through child-rearing, education, and modification of one's behaviour to conform with the demands of the society or group to which one belongs.

Indoctrination is the process of inculcating a person with ideas, attitudes, cognitive strategies or professional methodologies. A stereotype is any thought widely adopted about specific types of individuals or certain ways of behaving intended to represent the entire group of those individuals or behaviors as a whole. Normalisation refers to social processes through which ideas and actions come to be seen as "normal" and become taken-for-granted or "natural" in everyday life. Internalisation is an individual's acceptance of a set of norms and values via socialisation.

Ref: Colman AM. Oxford Dictionary of Psychology, 4th Edition. 2015.

The correct answer is: Socialisation

Question **48**

Not answered

Marked out of 1.00

Flag question

Sharon lives in the same household with her younger brother who is in a same-gender partnership. Her brother and his partner want to adopt a child. Which of the following statements regarding lesbian and gay (LG), and heterosexual adoptive parents is true?

Select one:

- Lesbian mothers, heterosexual mothers, gay fathers, and heterosexual fathers report no differences in parenting competence in adoptive families
- Heterosexual couples are more likely to adopt children than same-gender couples
- Adoptive parents of all types no longer face stigma in our society
-

LG adoptive parents face additional stigma based on sexual identity

Stigmatisation based on sexual identity has no effect on adoptive parents' mental health

Your answer is incorrect.

Explanation:

Adoptive parents of all types often face stigma related to 'non-traditional' family structures. Lesbian and gay (LG) adoptive parents often experience additional stigmatisation based on sexual identity, which in turn may negatively affect parents' mental health. Children and their adoptive sexual minority parents face stigma, overt discrimination, and microaggressions based on parent sexual orientation. Lesbian mothers, heterosexual mothers, and gay fathers all report higher parenting competence than heterosexual fathers in adoptive families. Same-gender couples may be up to seven times more likely to have adoptive children than different-gendered couples.

Ref: Farr RH, Vázquez CP. Stigma Experiences, Mental Health, Perceived Parenting Competence, and Parent–Child Relationships Among Lesbian, Gay, and Heterosexual Adoptive Parents in the United States. Frontiers in psychology. 2020 Mar 30;11:445.

The correct answer is: LG adoptive parents face additional stigma based on sexual identity

Question **49**

Not answered

Marked out of 1.00

Flag question

Which of the following Erikson's stages of development are paired correctly?

Select one:

1.5-3 years old – trust vs. mistrust

3-5 years old – initiative vs. guilt

12-18 years old – industry vs. inferiority

0-1.5 years old – autonomy vs. shame

40-65 years old – ego integrity vs. despair

Your answer is incorrect.

Explanation:

Erik Erikson created an original and highly influential theory of psychological development and crisis occurring in periods that extended across the entire life cycle. Stage 3 takes place from **about 3 years to about 5 years**. Its focus is **initiative versus guilt**. The child's increasing mastery of locomotor and language skills expands its participation in the outside world and stimulates omnipotent fantasies of broader exploration and conquest. Guilt over the drive for conquest and anxiety over the anticipated punishment are both assuaged in the child through repression of the forbidden wishes and development of a superego to regulate its initiative.

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. p. 1074.

The correct answer is: 3-5 years old – initiative vs. guilt

Question **50**

Not answered

Developmental milestones (Language)
Identify the age at which children should reach the following milestones in language

Marked out of 3.00

Flag question

development (3 questions)

1. Develops intonation and babbles as though speaking, but does not say words.

2. Uses 3-4 words together, and can change the order of words to make questions.

3. Understands puns and jokes.

Explanation:

1) 6-12 months: Babbling (consonant-vowel combinations) and responds when spoken to. 2) 2-3 years: By 3 years, children start using 3 to 4 word sentences with verbs that are generally understood by all adults. At this stage the 'wh' are common (who, why, where) 3) 6-7 years: Understands common idioms. Understands puns, humour

The correct answer is:

Develops intonation and babbling: 6-12 months

Uses 3-4 words together: 2-3 years

Understands puns and jokes: 6-7 years

Question **51**

Not answered

Marked out of 3.00

Flag question

Developmental Psychology

Various psychology and social theories have relevance to psychiatric practice. Which of the following concepts relates most closely to the below descriptions?

1. James Kearney is a 8-year-old boy who has started to demonstrate an awareness of the conservation of objects and inductive reasoning.

2. Laura Myers is a 4-year-old girl whose mother has been trying to get her to stop snatching toys from others. She has now started to respond to being denied treats if she snatches things from others.

3. Mrs Neville is a 32-year-old woman who is experiencing periods of despair and protest.

Explanation:

1) The first example describes a child in Piaget's concrete operational stage of cognitive development.

2) The second scenario is an example of moral development, with the child described being in the punishment avoidance stage, as described by Kohlberg.

3) The third vignette describes stages of grief. There are various theories around the stages of grief. Bowlby's 4 stages of grief include protest and despair.

Question **52**

Not answered

Marked out of 4.00

Flag question

Mahler's Stages of Development
For the situations given below, identify the Mahler's stage of development.

1. The infant moves autonomously and explores the outer world.

2. The infant feels as though he is completely in his own world.

3. The infant begins to gain awareness that s/he is a different entity from the mother.

Explanation:

1) Practicing occurs between 10 and 18 months. During this phase, the infant is able to crawl and eventually walk, exploring the outer world.

2) Normal autistic phase – during this stage (birth-2 months) the infant is uninterested in external stimuli, and is detached from the outer world.

3) Differentiation/hatching (5-10 months) is the stage where the child becomes aware that s/he is of a different entity from the mother.

Stage	Age	Description: Mahler's separation-individuation
Normal autism	Birth-2 months	Periods of sleep outweigh periods of arousal Reminiscent of intrauterine life
Symbiosis	2-5 months	Developing perceptual abilities enables infant to distinguish inner from outer world Mother-infant perceived as a single fused entity
Differentiation/ Hatching	5-10 months	Progressive neurological development and increased alertness Infant's attention drawn away from self to the outer world Physical and psychological distinctiveness from mother gradually appreciated
Practicing	10-18 months	Ability to move autonomously increases child's exploration of outer world
Rapprochement	18-24 months	Child slowly realises helplessness and dependence Need for independence alternates with need for closeness Child moves away from mother and comes back for reassurance
Object constancy	2-5 years	Child gradually comprehends and is reassured by permanence of mother and other important people, even when not in their presence

Adapted from Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition, 2022. pp. 892-893.

SPMM COURSE

Ref: Boland R, Verduin M. Kaplan and Sadock's Synopsis of Psychiatry, 12th Edition. 2022. pp. 892-893.

The correct answer is:

1. Infant moves autonomously - Practicing

2. Infant is in own world - Normal autistic phase

3. Infant gains awareness of being a different entity - Differentiation, Hatching

Question **53**

Not answered

Marked out of 5.00

Flag question

Theme- Memory
Match the following descriptions with the correct type of memory described

1. Remembering how to eat breakfast. [ ]

2. Remembering what you ate for breakfast. [ ]

3. Remembering what breakfast is. [ ]

4. Remembering after you have eaten breakfast that you need to pick up more cereal on your way home from work. [ ]

5. Remembering that after breakfast you took your bike out and cycled your usual route to work, and this felt tedious. [ ]

Explanation:

1) **Procedural** memory, or 'knowing how', is the unconscious memory of skills and how to do things. It is acquired through repetition and practice.

2) **Declarative** memory is memory of facts and events that can be consciously recalled (also known as explicit memory).

3) **Semantic** memory is a structured record of meanings, concepts, facts, knowledge about the world; it's to do with general factual knowledge that is shared with others, independent of personal experience.

4) **Prospective** memory refers to that which needs to be remembered in the future, and may be defined as 'remembering to remember'; it can be either event-based or time-based, and is often triggered by a cue (e.g. finishing breakfast).

5) **Episodic** memory refers to experiences and specific events in time in serial form, from which we can reconstruct the actual events that took place in our lives; it is the memory of autobiographical events, and has an emotional charge.

Question **54**

Not answered

Marked out of 5.00

Flag question

Measurement tools used in older subjects
Choose the most appropriate tool for each of the following situations in older subjects with cognitive impairment;

1. Monitoring cognitive decline (choose 2 options)
[ ]
[ ]

2. Monitoring decline in function of ADLs

3. Monitoring depression (choose 2 options)

Explanation:

1) There are many tools for measuring aspects of cognition, function and mental state in Alzheimer's disease. Though not the most sensitive/specific, the most commonly used tool to measure cognitive decline is the MMSE, however, that is not an option here. The MOCA and CAMCOG are two appropriate tools to assess cognition. The AMTS would not be sensitive enough to pick up cognitive decline, and the clock-drawing test and FAB, though assessing some aspects, are not general enough. 2) For assessment of activities of daily living, the OT department have many tools. The association don't endorse one specific tool above all in Alzheimer's but note that some aspects of each tool may need to be modified. From this list, the MOHOST is the only assessment of ADLs included. The BEHAVE-AD is an assessment of behaviour in Alzheimer's disease. 3) The MADRS and Hamilton Rating scales are both used to assess for depression in the elderly.

The correct answer is:

Monitoring cognitive decline: MOCA and CAMCOG

Monitoring decline in function of ADLs: MOHOST

Monitoring depression: Hamilton Rating Scale and MADRS

Question **55**

Not answered

Marked out of 3.00

Flag question

Electrolyte abnormality

What electrolyte abnormality could be expected in each of the following situations?

1. Rani, a 52-year-old woman under the oncologists for breast cancer complains of being excessively thirsty.

2. Jay, a 22-year-old man, was brought into A&E in a coma. His friend Josh said that they had been at a rave where he thinks Jay may have taken something. He describes Jay as being very chatty and overfamiliar. He said that because Jay had a lot of alcohol before, his friends encouraged him to drink lots of water to sober up.

3. Lena, a 19-year-old woman, is under her GP for low mood. In the past she suffered from bulimia and struggles with her weight. Her boyfriend broke up with her a few days ago. Her friend Jess brings her to A&E as Jess noticed Lena was a little confused. She also has a tremor and some numbness and is dizzy.

Explanation:

1) Hypercalcaemia - polydipsia is a common symptom of hypercalcaemia. Hypercalcaemia is

relatively common in cancer (20-30% of cancer patients develop hypercalcaemia). The most common cancers associated with hypercalcaemia are breast and lung cancer, and multiple myeloma.

2) Hyponatraemia - Jay has likely taken MDMA which results in chatty and overfamiliar behaviour, and excessive thirst. He has developed hyponatreamia secondary to over-drinking water due to this excessive thirst. MDMA also causes syndrome of inappropriate antidiuretic hormone (SIADH) which contributes to hyponatremia.

3) Metabolic alkalosis - metabolic alkalosis is associated with excessive vomiting. In this case, it is likely that Lena has started to binge and purge following the breakdown of her relationship putting her at risk of developing metabolic alkalosis.

Question **56**

Not answered

Marked out of 3.00

Flag question

Movement disorders
Choose the movement disorder that matches the description.

1. A man who kicks his wife once as he falls asleep.

2. A man on psychotropic medication for a long time who sleeps badly, wakes with a stiff back in the morning and takes at least 10 minutes to get out of bed.

3. A man who experiences slow writhing movements in peripheral limbs.

Explanation:

1) **Sleep myoclonus/hypnic jerk** is an involuntary twitch that occurs just as a person is beginning to fall asleep

2) **Parkinsonism** is associated with antipsychotic medication resulting in stiffness, bradykinesia and tremor

3) **Athetosis** is slow involuntary writing movements of the limbs caused by degeneration of the basal ganglia, as seen in Huntington's disease. It can also be secondary to stroke or trauma.

Question **57**

Not answered

Marked out of 5.00

Flag question

Personality disorder
If all the general criteria for a personality disorder were met, what personality disorder would the traits described in these vignettes be indicative of?

1. Mr Whitaker is berated by some of his acquaintances for holding onto money very tightly. He claims he has to hold onto it for future catastrophies, and will never offer to pay for someone else's drink. He is regarded as miserly and uptight.

2. Liliana is often at the centre of any party, and cannot bear to be overlooked or ignored. She makes sure she's the centre of attention by wearing provocative clothes, laughing loudly and talking copiously and vivaciously.

3. Mike always seems to have several girlfriends, often at the same time. They find him attractive and seductive, and he impresses them with his stories of evading the law for his own profit. The relationships don't last long and Mike doesn't seem bothered when they come to an end.

4. Nathan demonstrates the self-perception of being unique and superior to his peers.

5. Mary is persistently unreliable at work and has poor control of her finances.

Explanation:

1) **Anankastic** personality disorder: Traits include a preoccupation with orderliness and control (including over money), at the expense of flexibility. Rules are very rigidly maintained.

2) **Histrionic** personality disorder: Traits include excessive attention-seeking emotions including inappropriately seductive behaviour.

3) **Antisocial** personality disorder: Traits include ease on forming relationships, but not keeping them, superficial charm, lack of remorse for criminal or law-breaking activity, disregard for the rights of others.

4) **Narcissistic** personality disorder can be recognised by elevated sense of importance and significance compared to others.

5) **Emotionally unstable** personality disorder usually involves marked impulsivity and a desultory approach to life; this usually leads to difficulties with finances and at work.

Question **58**

Not answered

Marked out of 3.00

Flag question

Diagnosis of acute disturbances
In each of the below scenarios, which of the following options would be the most likely diagnosis?

1. Mrs. Greaves is a 74-year-old woman who has recently had an elective hip replacement. Whilst on the orthopaedic ward she develops confusion, dyspnoea and chest pain 3 days postoperatively.

2. Helen Honey is a 6-year-old girl whose parents take her to the GP after they and her teachers notice that she is exhibiting episodes of blank staring and unresponsiveness lasting 10 to 15 seconds, which Helen is unaware of.

3. Mrs. Ives is a 32-year-old woman with a diagnosis of bipolar affective disorder. She develops coarse tremor and ataxia after going for a run whilst staying at her holiday let in Spain during her summer holiday.

Explanation:

1) Chest pain and dyspnoea would make a **pulmonary embolus** more likely than the other options. Significant hypoxia could lead to confusion.

2) The second scenario appears to describe **absence seizures** (also known as petit mal seizures) with classic brief periods of unresponsiveness but without any movement abnormalities.

3) The third example describes likely **lithium toxicity** due to dehydration. The patient has exerted themselves in a warm environment and has experienced fluid loss as a result.

Question **59**

Not answered

Marked out of 3.00

Flag question

Anxiety disorders
In each of the below scenarios, which of the following options would be the most likely diagnosis?

1. Matthew attends a CAMHS clinic. He has recently changed schools due to experiencing several difficulties. He also refuses to go out with friends or parents because he finds it uncomfortable to eat in front of people.

2. Beth is a 22-year-old who has been experiencing a two-week history of low mood, nervousness, poor concentration and hopelessness following the breakdown of her relationship with her boyfriend three months ago.

3. Lauren is a 15-year-old who attends CAMHS clinic following a change in behaviour; her parents describe that she often blinks repeatedly or counts on her fingers. Lauren eventually confides that she repeatedly sees horrible images in her head.

Explanation:

1) Matthew has a diagnosis of social phobia or social anxiety characterised by difficulties in a school or social environment.

2) Beth is suffering from adjustment disorder, which occurs up to 3 months (usually within 1 month) after a stressor and does not persist for more than 6 months after the stressor.

3) Lauren has a diagnosis of obsessive-compulsive disorder describing obsessional intrusive thoughts and counting compulsions.

Finish review

PREVIOUS ACTIVITY
Fresh Practice Questions 2018 Set 3

NEXT ACTIVITY
Fresh Practice Questions 2020 Set 5

Jump to...

Drop here!

© 2022 SPMM Course Limited, UK.

Refund Policy | Terms & Conditions