# Pattern avoidance An explanation and proof

Yajit Jain, Deepak Narayanan and Leon Zhang

November 14, 2014

## Plan of action

- 1 Definition, Introduction and Motivation
- **2** Avoidance in  $S_n$
- **3** Avoidance in  $T_n$

Definition, Introduction and Motivation

# Lemma (Reversing Lemma)

The permutation  $\sigma$  avoids the permutation  $\pi$  iff  $\mathcal{R}(\sigma)$  avoids  $\mathcal{R}(\pi)$ .

#### Proof.

Need to only prove one direction (why?).

Suppose that  $\mathcal{R}(\sigma)$  does not avoid  $\mathcal{R}(\pi)$  if  $\sigma$  avoids  $\pi$ .

Reversing  $\mathcal{R}(\sigma)$  again produces a contradiction (why?), which implies that  $\mathcal{R}(\sigma)$  avoids  $\mathcal{R}(\pi)$  if  $\sigma$  avoids  $\pi$ .

# Corollary

For a permutation  $\pi$ ,  $s_n(\pi) = s_n(\mathcal{R}(\pi))$ .

# Definition (Flip of a sequence)

We define the *flip* of a sequence a as the sequence b with the same elements as a, but with the largest element swapped with the smallest element, the second largest element swapped with the second smallest element, etc. The flipping operator is denoted by  $\mathcal{F}$ .

### Example

$$\mathcal{F}(1324) = 4231.$$

## Example

$$\mathcal{F}(1243) = 4312.$$

# Lemma (Flipping Lemma)

The permutation  $\sigma$  avoids the permutation  $\pi$  iff  $\mathcal{F}(\sigma)$  avoids  $\mathcal{F}(\pi)$ .

#### Proof.

As before, need to only prove one direction.

Suppose  $\mathcal{F}(\sigma)$  does not avoid  $\mathcal{F}(\pi)$  if  $\sigma$  avoids  $\pi$ . (Here,  $\sigma \in \mathcal{S}_n$  and  $\pi \in \mathcal{S}_k$ )

Let  $\mathcal{F}(\sigma) = a_1 \cdots a_n$  and let  $\mathcal{F}(\pi) = b_1 \cdots b_k$ . Then there is some subsequence  $a_{i_1} \cdots a_{i_k}$  such that  $a_{i_c} < a_{i_d}$  iff  $b_c < b_d$ .

But then,  $n+1-a_{i_c}>n+1-a_{i_d}$  iff  $k+1-b_c>k+1-b_d$ .

This implies that  $\sigma$  does not avoid  $\pi$ , a contradiction (why?).

## Corollary

For a permutation  $\pi$ ,  $s_n(\pi) = s_n(\mathcal{F}(\pi))$ .

#### Lemma

The permutations of  $\{1, 2, ..., k, k+1\}$  ending in i that avoid the pattern 312 are precisely those of the form,

$$\pi_1\pi_2i$$

the concatenation of  $\pi_1, \pi_2$ , and i, where  $\pi_1$  is a permutation of  $\{1, 2, \ldots, i-1\}$  that avoids the pattern 312 and  $\pi_2$  is a permutation of  $\{i+1, \ldots, k+1\}$  that avoids the pattern 312.

#### Proof.

Assume that  $\exists$  some permutation  $\pi$  of  $\{1,2,...,k,k+1\}$  that avoids 312, ending with value i such that some integer x < i is to the right of some integer y > i. Clearly,  $\pi$  does not avoid 312. (why?)

It follows that for any permutation  $\pi$  avoiding 312 and ending in i, all x < i must be to the left of all y > i. Hence,  $\pi$  can be written as

$$\pi_1\pi_2i$$

where  $\pi_1$  is a permutation of  $\{1, 2, ..., i-1\}$  and  $\pi_2$  is a permutation of  $\{i+1, i+2, ..., k+1\}$ .

Furthermore, it is clear that the permutations  $\pi_1$  and  $\pi_2$  must avoid 312 as well. (why?)

#### Definition

The Catalan numbers are the sequence of positive integers  $C_i$  defined as follows,

$$C_0 = 1, \ C_{n+1} = \sum_{i=0}^n C_i C_{n-i} \text{ for } n \ge 0$$

#### Theorem

 $s_n(312), s_n(132), s_n(213)$ , and  $s_n(231)$  are equal to  $C_n$ , the  $n^{th}$  Catalan number.

#### Proof.

Assume that for all i from 1 to k, the number of permutations of  $\{1, 2, ..., i\}$  that avoid the order 312 as a subsequence is  $C_i$ .

The number of permutations of  $\{1, 2, ..., k, k+1\}$  that avoid 312 can be counted by enumerating through all possible values of the last term of a valid permutation.

From the above lemma, all permutations of  $\{1, 2, ..., k, k+1\}$  ending in i that avoid 312 are precisely those of the form

$$\pi = \pi_1 \pi_2 i$$

where  $\pi_1$  is a permutation of  $\{1, 2, ..., i-1\}$  that avoids 312 and  $\pi_2$  is a permutation of  $\{i+1, i+2, ..., k+1\}$  that avoids 312.

# Proof (contd.)

It follows that the total number of permutations  $\pi$  avoiding 312 and ending in i is

$$C_{i-1} \cdot C_{k-i+1}$$

Summing over all possible values of i, the total number of permutations of  $\{1, 2, ..., k + 1\}$  that avoid 312 is equal to,

$$\sum_{i=1}^{k+1} C_{i-1} \cdot C_{k-i+1} = \sum_{i=0}^{k} C_i \cdot C_{k-i} = C_{k+1}$$

By the flipping lemma and reversing lemma,  $s_n(132)$ ,  $s_n(213)$  and  $s_n(231)$  are the sequence of Catalan numbers as well.

Pattern avoidance

# Avoidance in $T_n$