

Awesome Title

Ben Adcock, Rick Archibald, Anne Gelb, Guohui Song, Rodrigo B. Platte, and Edward G. Walsh.

Abstract—

Index Terms—Fourier Data l^1 regularization Split Bregman MRI.

I. INTRODUCTION

THIS

II. METHODS

A. Measurements


$$S(k_x, k_y, t) = \iint_{\Omega} m(x, y) e^{-R^*(x, y)t + i f(x, y)t} e^{-ik_x x - ik_y y} \partial \Omega \quad (1)$$

B. Sampling

C. Fourier Frames

D. TV optimization

III. NUMERICAL



(a) *InterpolationError*

(b) *UniqueSamplePoints*

Fig. 1. (a) Interpolation Error and (b) Unique Sample Points are factors in choosing spiral MRI

Department of Mathematics, Simon Fraser University, Burnaby, BC V5A 1S6 (ben_adcock@sfu.ca)

Computer Science and Mathematics Division, Oak Ridge National Laboratory, Oak Ridge, TN 37831 (archibaldrk@ornl.gov)

School of Mathematical and Statistical Sciences, Arizona State University, Tempe, AZ, 85287 (annegelb@asu.edu)

Department of Mathematics, Clarkson University, Potsdam, NY 13699. (gsong@clarkson.edu)

School of Mathematical and Statistical Sciences, Arizona State University, Tempe, AZ, 85287 (rbp@asu.edu)

Department of Neuroscience, Brown University, Providence, RI 02912 (edward_walsh@brown.edu)




Fig. 2. Time slice




Fig. 3. Time series analysis

IV. CONCLUSION

Acknowledgments

This work is supported in part by grants NSF-DMS 1216559 and AFOSR FA9550-12-1-0393. The submitted manuscript is based upon work, authored in part by contractors [UT-Battelle LLC, manager of Oak Ridge National Laboratory (ORNL)], and supported by the U.S. Department of Energy, Office of Science, Office of Advanced Scientific Computing Research, Applied Mathematics program. Accordingly, the U.S. Government retains a non-exclusive, royalty-free license to publish or reproduce the published form of this contribution, or allow others to do so, for U.S. Government purposes.