

# Regulation of membrane scission in yeast endocytosis

Deepikaa Menon<sup>1</sup> and Marko Kaksonen<sup>1\*</sup>

\*For correspondence:

[Marko.Kaksonen@unige.ch](mailto:Marko.Kaksonen@unige.ch) (

)

<sup>1</sup>Department of Biochemistry, University of Geneva, Geneva, Switzerland

**Abstract** This is not going to elife

## Introduction (Level 1 heading)

Thanks for using Overleaf to write your article. Your introduction goes here! Some examples of commonly used commands and features are listed below, to help you get started.

Here's a second paragraph to test paragraph indents. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

## Results

**Vps1 does not influence coat or scission dynamics. Synaptojanins likely influence vesicle uncoating, but not scission dynamics.**

Endocytic membrane scission in mammalian cells is understood to be driven by constriction of the tubule neck by the Gtpase Dynamin (a bunch of dynamin papers). In yeast, it has been reported that the Dynamin-like protein Vps1 is recruited to endocytic sites (refAyscough). To test whether Vps1 influences scission, endocytic coat dynamics are observed in cells lacking Vps1 and compared against WT cells. Fig1a shows kymographs of coat protein Sla1 endogenously tagged at the N-terminus with eGFP in WT and *vps1Δ* cells. Dynamics of Sla1 in WT and *vps1Δ* cells are the same. In Fig.1b, the averaged centroid trajectory (ref2andrea)- henceforth centroid- of Sla1-eGFP is tracked in 50 endocytic sites in *vps1Δ* and wild-type cells. Inward movement of Sla1 centroid serves as a proxy for plasma membrane movement through the endocytic process (ref2andrea).  $y=0$  is set as the position of arrival of the protein complex, before inward movement begins. Centroid movement of Sla1-eGFP in wild-type cells shows a linear movement to about 150nm, and Sla1 movement in *vps1Δ* cells is the same.

Centroid tracking has shown that the number of yeast N-BAR protein Rvs167 peaks at the time of scission, and is followed by an rapid loss of fluorescent intensity, concomitant with a sharp jump of the centroid into the cytoplasm. This jump, also seen in Rvs167-GFP kymographs, is interpreted as loss of protein on the invagination on the membrane tube, causing an apparent spatial jump to the protein localized at the base of the newly formed vesicle. Kymographs of Rvs167-GFP in Vps1 deleted cells, show the same jump, indicating that vesicles are formed in the same position in Vps1

42 deletion cells as in WT cells. Hence, although Vps1 deletion leads to a growth defect at 37C (as has  
43 been shown before, supplementary Fig.1), lack of Vps1 protein does not influence the endocytic  
44 process.

#### 45 **Synaptojanins likely influence vesicle uncoating, but not scission dynamics.**


46 As an alternate to forces from Dynamin constriction, Liu et al (refliu) have proposed that an inter-  
47 action between PiP2-hydrolyzing Synaptojanins and BAR proteins could drive membrane scission.  
48 Here Rvs BAR domains would form a scaffold on the membrane tube, preventing hydrolysis of  
49 underlying PIP2. Synaptojanin would arrive at invaginated membranes, and hydrolyse unprotected  
50 PIP2. This generates a lipid boundary between BAR-protected PIP2 at the tube and hydrolyzed PIP2  
51 at the bud tip. A line tension thus formed at the interphase between the two lipid types would then  
52 generate enough force to pinch off a vesicle.

53 Of the three Synaptojanin-like proteins in yeast- Inp51, Inp52 and Inp53- Inp51 exhibits a diffuse  
54 cytoplasmic signal. Inp53 localizes to patches within the cytoplasm- cellular localization is consistent  
55 with involvement in trans-Golgi signalling (refGolgi). Inp53 was not investigated further. Inp52  
56 localizes to cortical actin patches that are endocytic sites. Two channel alignment shows that Inp52  
57 patches arrive in the late scission stage, and localizes to the bud tip, consistent with a role in  
58 membrane scission.

59 Role of Inp51 and Inp52 are tested by following Sla1-GFP and Rvs167-GFP in cells with either  
60 Inp51, Inp52, or both deleted. Retraction events do not significantly increase compared to the WT in  
61 either inp51del or inp52del cells. Magnitude and speed of coat movement in inp51del is the same  
62 as the WT. In inp52del cells, coat movement also has the same magnitude and speed, but GFP signal  
63 is persistent after membrane scission. This delay in decrease of Sla1-GFP signal is consistent with  
64 delay in vesicle uncoating rather than membrane scission. Similarly, Rvs167 disassembly has a  
65 delay, while the assembly is similar to WT, indicating a delay in removing endocytic proteins from the  
66 newly formed vesicle. Assembly of Rvs167 has a delay in inp51 deleted cells, which could indicate a  
67 defect in recruiting proteins to endocytic sites, or in progression of endocytic invaginations. Since  
68 Sla1 movement is the same, we suggest a defect in the former rather than latter.

#### 69 **Rvs deletion reduces coat movement**

70 The Rvs complex itself is known to influence scission: deletion reduces scission efficiency by 30%  
71 (ref1marko). Failed scission events are characterized by inward movement, followed by retraction  
72 of the coat protein Sla1 (ref1marko). Contribution of Rvs to the scission process however, is  
73 currently unclear. In the remaining 70% of successful invaginations, inward movement of the  
74 coat protein Sla1 also deviates from the wild-type. In Fig.1, the averaged centroid trajectory  
75 (ref2andrea) of Sla1-eGFP is tracked in rvs167deletion and wild-type cells. Time alignment is  
76 established by tracking the centroid of a second protein, here m-Cherry tagged Actin binding  
77 protein Abp1. Simultaneous tracking of GFP-tagged protein of interest and m-Cherry tagged  
78 Abp1 allows us to align all other proteins to the Abp1 reference centroid (ref2andrea). Time=0, is  
79 established as the peak of the Abp1 fluorescence intensity, which in wild-type is concomitant with  
80 the peak of Rvs167 fluorescent intensity (ref2andrea, ref3wanda), and is time in which scission  
81 occurs. Sla1 centroid in rvs167deletion cells follows the wild-type centroid movement till about  
82 60nm, after which movement slows down and scission occurs. That scission occurs at shorter  
83 invagination lengths is confirmed by formation of smaller vesicles and shorter invagination lengths  
84 in rvs167deletion cells, quantified by Correlative light and electron microscopy (CLEM) (ref3wanda).  
85 Invagination lengths of 60nm is the time window for arrival of Rvs167 (ref3wanda), indicating that  
86 coat movement of endocytic sites in rvs167deletion cells progresses normally till the expected  
87 arrival of Rvs.



**Figure 1.** A half-columnwidth image using wrapfigure, to be used sparingly. Note that using a wrap figure before a sectional heading, near other floats or page boundaries is not recommended, as it may cause interesting layout issues. Use the optional argument to wrapfigure to control how many lines of text should be set half-width alongside it.

### Rvs BAR domains recognize membrane curvature in-vivo

The curved tertiary structure and liposome binding assays of N-BAR domains have suggested that they may have a preference for curved membrane that match their own intrinsic curvature. Alternately, they may also impose their curvature on flat membrane and induce curvature formation. In-vivo, the curvature interaction of Rvs167 has not been tested. In order to do so, we delete the SH3 domain of Rvs167 observe the localization of endogenously tagged Rvs167-eGFP and BAR-eGFP and Abp1-mCherry in WT and sla2deletion cells. Sla2 acts as the molecular linker between forces exerted by the actin network and the plasma membrane. Sla2deletion cells therefore have a polymerizing actin network, but the membrane remains flat and endocytosis fails. In these cells, the full-length Rvs167 protein co-localizes with Abp1-mCherry, indicating that it is recruited to endocytic sites. BAR-eGFP localization is removed, except for rare transient patches that do not co-localize with Abp1-mCherry.

### Rvs SH3 domains contribute to curvature independent localization

We have shown that BAR domains need membrane curvature to localize. Full-length Rvs167, however, is recruited to endocytic patches in sla2deletion cells. This indicates that a second interaction, that is not the BAR-curvature dependent, recruits the protein to endocytic sites. This interaction must come from the SH3 region, showing that Rvs localization is dependent on both BAR as well as SH3 domain interactions. Absence of the SH3 domain also reduces total recruitment of Rvs and Abp1 protein, giving the SH3 domain an important and surprising role in regulating the late stage of endocytosis.

## SH3 domains are recruited by Myosin 5

SH3 domains have been shown to interact with several proteins in the actin module of endocytosis: Las17, type I myosins, and Vrp1 all have genetic or physical interactions with Rvs167 SH3 domains (Lila and Drubin, 1997; Colwill et al., 1999; Madania et al., 1999; Liu et al., 2009). We tested the interaction by studying the localization of full-length Rvs167 in cells with one of these proteins deleted, and treated with Las17 to reproduce the situation in which BAR-curvature interaction is removed. As seen in (some figure), deletion of neither Las17 nor Myo3 in combination with LatA treatment does not remove the localization of Rvs167. Deletion of Vrp1 and Myo5, with LatA treatment removes localization of Rvs167. Since Vrp1 is required for the recruitment of Myo5 (refMyo5), SH3 domains interact with Myo5 rather than Vrp1.

what about the differences in myo5 and myo3 number... if the Rvs recruitment only slightly depended on myo3 we probably wouldnt see a difference

## N-helix and GPA domains do not contribute to recruitment of Rvs or membrane movement


Etiam euismod. Fusce facilisis lacinia dui. Suspendisse potenti. In mi erat, cursus id, nonummy sed, ullamcorper eget, sapien. Praesent pretium, magna in eleifend egestas, pede pede pretium lorem, quis consectetur tortor sapien facilisis magna. Mauris quis magna varius nulla scelerisque imperdiet. Aliquam non quam. Aliquam porttitor quam a lacus. Praesent vel arcu ut tortor cursus volutpat. In vitae pede quis diam bibendum placerat. Fusce elementum convallis neque. Sed dolor orci, scelerisque ac, dapibus nec, ultricies ut, mi. Duis nec dui quis leo sagittis commodo.

## Increased BAR domain recruitment corresponds to increased membrane movement

The decreased Sla1 movement in BAR cells can be explained by a loss of interaction of the SH3 domain, or by reduced recruitment of the BAR domains. To check whether increasing the recruitment of the Rvs complex alone can rescue reduced Sla1 movement, the Rvs167 and Rvs161 ORF was duplicated endogenously (ref Huber dude) in diploid and haploid yeast cells. In diploid cells, Rvs duplication results in either 4x copies of both Rvs genes, 2x copies (WT diploid) or 1x copies, in which one gene of Rvs167 and Rvs161 are deleted. We show that amount of Rvs167 recruited to sites increases linearly, without changing either the rate of movement or total movement before scission of Sla1. Similarly, in haploid cells, increasing the gene copy of Rvs167 and Rvs161 results in increased recruitment of Rvs167, without influencing the dynamics of Sla1. Expressing two instead of one copy of the Rvs167 BAR domain alone rescues the loss of Sla1 movement in the 1x copy of BAR domain alone, as well the inward jump of BAR-GFP itself. The loss of inward movement in 1xBAR suggests that smaller vesicles are produced in these cells, confirmed by CLEM. This would in corollary indicate that the increased inward movement in 2xBAR produces WT-sized vesicles.

## Discussion

Morbi luctus, wisi viverra faucibus pretium, nibh est placerat odio, nec commodo wisi enim eget quam. Quisque libero justo, consectetur a, feugiat vitae, porttitor eu, libero. Suspendisse sed mauris vitae elit sollicitudin malesuada. Maecenas ultricies eros sit amet ante. Ut venenatis velit. Maecenas sed mi eget dui varius euismod. Phasellus aliquet volutpat odio. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Pellentesque sit amet pede ac sem eleifend consectetur. Nullam elementum, urna vel imperdiet sodales, elit ipsum pharetra ligula, ac pretium ante justo a nulla. Curabitur tristique arcu eu metus. Vestibulum lectus. Proin mauris. Proin eu nunc eu urna hendrerit faucibus. Aliquam auctor, pede consequat laoreet varius, eros tellus scelerisque quam, pellentesque hendrerit ipsum dolor sed augue. Nulla nec lacus.



**Figure 2.** A half-columnwidth image using wrapfigure, to be used sparingly. Note that using a wrapfigure before a sectional heading, near other floats or page boundaries is not recommended, as it may cause interesting layout issues. Use the optional argument to wrapfigure to control how many lines of text should be set half-width alongside it.

## Methods and Materials

Guidelines can be included for standard research article sections, such as this one.

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

## Citations

LaTeX formats citations and references automatically using the bibliography records in your .bib file, which you can edit via the project menu. Use the `\cite` command for an inline citation, like `?`, and the `\citep` command for a citation in parentheses (`?`). The LaTeX template uses a slightly-modified Vancouver bibliography style. If your manuscript is accepted, the eLife production team will re-format the references into the final published form. *It is not necessary to attempt to format the reference list yourself to mirror the final published form.* Please also remember to **delete the line** `\nocite{*}` in the template just before `\bibliography{...}`; otherwise *all* entries from your .bib file will be listed!

## Acknowledgments

Additional information can be given in the template, such as to not include funder information in the acknowledgments section.