

Regulation of membrane scission in yeast endocytosis

Deepikaa Menon^{1*} and Marko Kaksonen^{2*}

*For correspondence:

email1@example.com (FMS);
email2@example.com (FS)

[†]These authors contributed equally to this work

[‡]These authors also contributed equally to this work

Present address: [§]Department, Institute, Country; [¶]Department, Institute, Country

¹ Institution 1

Abstract This is not going to elife

Introduction (Level 1 heading)

Thanks for using Overleaf to write your article. Your introduction goes here! Some examples of commonly used commands and features are listed below, to help you get started.

Here's a second paragraph to test paragraph indents. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Results

Vps1 does not influence coat or scission dynamics. Synaptojanins likely influence vesicle uncoating, but not scission dynamics.

Endocytic membrane scission in mammalian cells is understood to be driven by constriction of the tubule neck by the Gtpase Dynamin (a bunch of dynamin papers). In yeast, it has been reported that the Dynamin-like protein Vps1 is recruited to endocytic sites (refAyscough). To test whether Vps1 influences scission, coat dynamics are observed in cells lacking Vps1 (left, Fig.1) and compared against WT cells (right, Fig.1). Centroid tracking various endocytic proteins in yeast has shown that the centroid of the yeast N-BAR protein Rvs167 peaks at the time of scission, and is followed by an rapid loss of fluorescent intensity, concomitant with a sharp jump into the cytoplasm. This jump is interpreted as loss of protein on the invagination on the membrane tube, to the fluorescent intensity at the base of the newly formed vesicle. Kymographs of Rvs167-GFP in Vps1 deleted cells however, show the same jump, indicating that vesicles are formed in the same position in Vps1 deletion cells as in WT cells. Vps1 deletion leads to a growth defect at 37C as has been shown before (supplementary Fig.1) Fig1a shows kymographs of coat protein Sla1 endogenously tagged at the N-terminus with eGFP in WT and *vps1Δ* cells. Inward movement of Sla1 in WT and *vps1del* cells are the same. In Fig.1b, the averaged centroid trajectory (ref2andrea)- henceforth centroid- of Sla1 endogenously tagged at the C-terminus with eGFP is tracked in 50 endocytic sites in *vps1deletion* and wild-type cells. Beginning of inward movement of the centroid is set as the initial position of averaged centroid. Inward movement of Sla1 centroid serves as a proxy for plasma membrane movement through the endocytic process (ref2andrea). Centroid movement

of Sla1-eGFP in wild-type cells shows a linear movement to about 150nm, and Sla1 movement in vps1del cells is the same. Position of the vesicle formed at the end of scission process is determined from the final position of the Rvs167 centroid (ref2andrea). Rvs167 centroid position in WT and vps1-del cells indicates that vesicle formation is the same as in WT cells.

Synaptojanins likely influence vesicle uncoating, but not scission dynamics.

Liu et al (refliu) have proposed that an interaction between PiP2-hydrolyzing Synaptojanins and BAR proteins could drive membrane scission. Here, Rvs BAR domains would form a scaffold on the membrane tube, preventing hydrolysis of underlying PIP2. Synaptojanin would arrive at invaginated membranes, and hydrolyse unprotected PIP2. This generates a lipid boundary between BAR-protected PIP2 at the tube and hydrolyzed PIP2 at the bud tip. A line tension thus formed at the interphase between the two lipid types would then generate enough force to pinch off a vesicle.

Of the three Synaptojanin-like proteins in yeast- Inp51, Inp52 and Inp53- Inp51 exhibits a diffuse cytoplasmic signal. Inp53 localizes to patches within the cytoplasm(refGolgi)- cellular localization is consistent with involvement in trans-Golgi signalling, and is taken to indicate lack of involvement in endocytosis, shown also in other studies (refother studies). Inp53 was not investigated further. Inp52 localizes to cortical actin patches that are endocytic sites. Two channel alignment shows that Inp52 patches arrive in the late scission stage, and localizes to the bud tip, consistent with a role in membrane scission.

Role of Inp51 and Inp52 are tested by following Sla1-GFP and Rvs167-GFP in cells with either Inp51, Inp52, or both deleted. Retraction events do not significantly increase compared to the WT in either inp51del or inp52del cells. Magnitude and speed of coat movement in inp51del is the same as the WT. In inp52del cells, coat movement also has the same magnitude and speed, but GFP signal is persistent after membrane scission. This delay in decrease of Sla1-GFP signal is consistent with delay in vesicle uncoating rather than membrane scission. Similarly, Rvs167 disassembly has a delay, while the assembly is similar to WT, indicating a delay in removing endocytic proteins from the newly formed vesicle. Assembly of Rvs167 has a delay in inp51deleted cells, which could indicate a defect in recruiting proteins to endocytic sites, or in progression of endocytic invaginations. Since Sla1 movement is the same, we suggest a defect in the former rather than latter.

Rvs deletion reduces coat movement

The Rvs complex is known to influence scission: deletion reduces scission efficiency by 30% (ref1marko). Failed scission events are characterized by inward movement, followed by retraction of the coat protein Sla1 (ref1marko). Contribution of Rvs to the scission process, and therefore, an understanding of why its absence might cause a scission defect, is currently unclear. In the remaining 70% of successful invaginations, inward movement of the coat protein Sla1 also deviates from the wild-type. In Fig.1, the averaged centroid trajectory (ref2andrea)- henceforth centroid- of Sla1 endogenously tagged at the C-terminus with eGFP is tracked in 50 endocytic sites in rvs167deletion and wild-type cells. Beginning of inward movement is set as the initial position of averaged centroid. Inward movement of Sla1 centroid serves as a proxy for plasma membrane movement through the endocytic process (ref2andrea). Time alignment is established by tracking the centroid of a second protein, here m-Cherry tagged Actin binding protein Abp1. Simultaneous tracking of GFP-tagged protein of interest and m-Cherry tagged Abp1 allows us to align all other proteins to the Abp1 reference centroid (ref2andrea). Scission time, $t=0$, is established as the peak of the Abp1 fluorescence intensity, which in wild-type is concomitant with the peak of Rvs167 fluorescent intensity (ref2andrea, ref3wanda). Centroid movement of Sla1-eGFP in wild-type cells shows a linear movement to about 150nm. Sla1 centroid in rvs167deletion cells follows the wild-type centroid movement till about 60nm, after which movement slows down and scission occurs. That scission occurs at shorter invaginations lengths is confirmed by formation of smaller vesicles and shorter invagination lengths in rvs167deletion cells, quantified by Correlative light




Figure 1. A half-columnwidth image using wrapfigure, to be used sparingly. Note that using a wrapfigure before a sectional heading, near other floats or page boundaries is not recommended, as it may cause interesting layout issues. Use the optional argument to wrapfigure to control how many lines of text should be set half-width alongside it.

91 and electron microscopy (CLEM) (ref3wanda). Invagination lengths of 60nm is the time window for
 92 arrival of Rvs167 (ref3wanda), indicating that coat movement of endocytic sites in rvs167deletion
 93 cells progresses normally till the expected arrival of Rvs.

94 Discussion

95 Morbi luctus, wisi viverra faucibus pretium, nibh est placerat odio, nec commodo wisi enim eget
 96 quam. Quisque libero justo, consectetur a, feugiat vitae, porttitor eu, libero. Suspendisse sed
 97 mauris vitae elit sollicitudin malesuada. Maecenas ultricies eros sit amet ante. Ut venenatis velit.
 98 Maecenas sed mi eget dui varius euismod. Phasellus aliquet volutpat odio. Vestibulum ante ipsum
 99 primis in faucibus orci luctus et ultrices posuere cubilia Curae; Pellentesque sit amet pede ac sem
 100 eleifend consectetur. Nullam elementum, urna vel imperdiet sodales, elit ipsum pharetra ligula,
 101 ac pretium ante justo a nulla. Curabitur tristique arcu eu metus. Vestibulum lectus. Proin mauris.
 102 Proin eu nunc eu urna hendrerit faucibus. Aliquam auctor, pede consequat laoreet varius, eros
 103 tellus scelerisque quam, pellentesque hendrerit ipsum dolor sed augue. Nulla nec lacus.

104 Methods and Materials

105 Guidelines can be included for standard research article sections, such as this one.

106 Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique,
 107 libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing
 108 semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie
 109 nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum.
 110 Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim.
 111 Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus.




Figure 2. A half-columnwidth image using wrapfigure, to be used sparingly. Note that using a wrapfigure before a sectional heading, near other floats or page boundaries is not recommended, as it may cause interesting layout issues. Use the optional argument to wrapfigure to control how many lines of text should be set half-width alongside it.

112 Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus
113 eu enim. Vestibulum pellentesque felis eu massa.

114 Citations

115 LaTeX formats citations and references automatically using the bibliography records in your .bib
116 file, which you can edit via the project menu. Use the `\cite` command for an inline citation, like
117 `?`, and the `\citep` command for a citation in parentheses (`?`). The LaTeX template uses a slightly-
118 modified Vancouver bibliography style. If your manuscript is accepted, the eLife production team
119 will re-format the references into the final published form. *It is not necessary to attempt to format*
120 *the reference list yourself to mirror the final published form.* Please also remember to **delete the line**
121 `\nocite{*}` in the template just before `\bibliography{...}`; otherwise *all* entries from your .bib
122 file will be listed!

123 Acknowledgments

124 Additional information can be given in the template, such as to not include funder information in
125 the acknowledgments section.