

STATISTICS WORKSHEET-5

Q1 to Q10 are MCQs with only one correct answer. Choose the correct option.

1. Using a goodness of fit, we can assess whether a set of obtained frequencies differ from a set of frequencies.
- a) Mean
 - b) Actual
 - c) Predicted
 - d) Expected

Ans:- d) Expected

2. Chisquare is used to analyse
- a) Score
 - b) Rank
 - c) Frequencies
 - d) All of these

Ans:-c) Frequencies

3. What is the mean of a Chi Square distribution with 6 degrees of freedom?
- a) 4
 - b) 12
 - c) 6
 - d) 8

Ans:-c) 6

4. Which of these distributions is used for a goodness of fit testing?
- a) Normal distribution
 - b) Chisquared distribution
 - c) Gamma distribution
 - d) Poisson distribution

Ans:-b) Chisquared distribution

5. Which of the following distributions is Continuous
- a) Binomial Distribution
 - b) Hypergeometric Distribution
 - c) F Distribution
 - d) Poisson Distribution

Ans:- F Distribution

6. A statement made about a population for testing purpose is called?
- a) Statistic
 - b) Hypothesis
 - c) Level of Significance
 - d) TestStatistic

Ans:-b) Hypothesis

7. If the assumed hypothesis is tested for rejection considering it to be true is called?
- a) Null Hypothesis
 - b) Statistical Hypothesis
 - c) Simple Hypothesis
 - d) Composite Hypothesis

Ans:-a)Null Hypothesis

8. If the Critical region is evenly distributed then the test is referred as?
- a) Two tailed
 - b) One tailed
 - c) Three tailed
 - d) Zero tailed


Ans:-a)Two tailed

9. Alternative Hypothesis is also called as?
- a) Composite hypothesis
 - b) Research Hypothesis
 - c) Simple Hypothesis
 - d) Null Hypothesis

Ans:- b)Research Hypothesis

10. In a Binomial Distribution, if 'n' is the number of trials and 'p' is the probability of success, then the mean value is given by _____
- a) np
 - b) n

Ans:-a)np



FLIP ROBO