Fonollosa's minimum-modulus conjecture: a sign characterization, a cyclic-order lemma, and exhaustive verification for $n = 8$

Deepnil Ray*

10 November 2026

Abstract

A set A of n residues modulo N is *valid* if the only multiset of size n drawn from A whose sum equals $\sum_{a\in A} a$ is A itself. Fonollosa (arXiv:2607.08366) conjectured that a valid n-set forces $N \geq 2^n - 2^{\lfloor \log_2 n \rfloor}$. **Verdict: PARTIAL.** The conjecture is not proved here. We prove (i) a sign characterization of validity in terms of the relation lattice; (ii) a cyclic-order lemma: whenever a difference $a - c$ of two elements generates $\mathbb{Z}_N$, the 2^{n-2} subset sums of the remaining elements are 2-separated along the $(a - c)$-cycle and shearing by the weight preserves their cyclic order; (iii) as consequences, $N \geq 2^{n-1}+n-2$ whenever some difference is a unit (in particular for every prime N), and $N \geq 5 \cdot 2^{n-3}$ when in addition A contains $3a - 2c$ or $4a - 3c$, or when $n \leq 5$. By an exhaustive, symmetry-reduced computer search with controls we verify the conjecture for $n = 8$ (the author of the conjecture reports exact certificates for $n \leq 7$), and we find that for every $n \leq 8$ the moduli $N < 2^n$ admitting a valid n-set are exactly $N = 2^n - 2^k$ with $2^k \leq n$. We record a sporadic valid 4-set modulo 15 that is not affinely equivalent to $\{1, 2, 4, 8\}$, and we show by computation that the weaker "all-anchor dissociated" property has the same threshold for $n \leq 6$ but a smaller one (120 instead of 124) for $n = 7$.

1 Verdict, worst news first

- **Not established:** Fonollosa's Conjecture 1 for general n. The best lower bound valid for *all* valid sets remains the bound $N \geq 2^{n-1}$ of Inal (with $N \geq 2^{n-1}+1$ for cyclic groups and $n \geq 3$, noted in an October 2026 public issue, see Section 2). Our new unconditional-in-n bounds need a hypothesis (a difference that is a unit) and are far from the conjectured $2^n - 2^{\lfloor \log_2 n \rfloor}$.
- **PROVED:** Proposition 3, Corollary 4, Lemma 6, Corollary 7, Theorem 8, Propositions 9, 11, 12, 13.
- **VERIFIED** (exhaustive search, bounds stated, controls fired): Theorem 14 and the tables of Section 5.
- **CONJECTURED:** Conjectures 17 and 10.

2 The problem and prior work

Throughout $n \geq 2$, $N \geq 2$, and $A \subseteq \mathbb{Z}_N$ with $|A| = n$.

*Independent researcher. ORCID 0000-0001-8779-0700. Computations and drafting were done with Claude (Anthropic); every claim below carries one of the labels PROVED, VERIFIED, CONJECTURED.

Definition 1 (Fonollosa)**.** *A is valid mod N* if for every $k \in \mathbb{N}^A$ with $\sum_a k_a = n$ and $\sum_a k_a a \equiv \sum_a a \pmod N$ one has $k_a = 1$ for all a.

Fonollosa [1] proved that for the set $\{2^i - 1 : 0 \le i < n\}$ the least valid modulus is $2^n - 2^{\lfloor \log_2 n \rfloor}$ (formalized in Lean 4, [2]) and stated:

Conjecture 2 (Fonollosa, Conjecture 1 of [1])**.** *If $N < 2^n - 2^{\lfloor \log_2 n \rfloor}$ then no n-subset of $\mathbb{Z}_N$ is valid.*

Status checked on 10 November 2026: the statement is listed as open in the Formal Conjectures repository [3] (file `Arxiv/2607.08366/MinModulus.lean`, repository state of 9 November 2026). According to [1] (as read through the arXiv listing and the README of [2]) the conjecture has exact CP-SAT certificates for $n \le 7$ and non-exhaustive solver searches for $n \le 13$. The README of [2] attributes to M. Inal (OSF preprint, DOI 10.17605/OSF.IO/C58Q9, which we have not read directly) the bound $|G| \ge 2^{n-1}$ for every finite abelian group G carrying a valid n-tuple, with equality only for $(\mathbb{Z}_2)^{n-1}$; the Lean proof of this bound is in [2]. A public research issue [4] dated 22 October 2026 records $N \ge 2^{n-1} + 1$ for cyclic groups and $n \ge 3$ and states that the conjecture is unresolved. We found no later work. We do not claim priority for any statement that follows without the caveat that the literature on this three-month-old problem is small and partly outside the refereed record.

3 A sign characterization

Let $\Lambda = \Lambda(A, N) = \{x \in \mathbb{Z}^A : \sum_a x_a = 0,\ \sum_a x_a a \equiv 0 \pmod N\}$ be the lattice of sum-zero relations.

Proposition 3 (PROVED)**.** *A is valid mod N if and only if every nonzero $x \in \Lambda$ has an entry ≥ 2 and an entry ≤ -2.*

Proof. The map $k \mapsto x = k - \mathbf{1}$ is a bijection from $\{k \in \mathbb{N}^A : \sum k_a = n,\ \sum k_a a \equiv \sum a\}$ onto $\{x \in \Lambda : x_a \ge -1\ \forall a\}$, and $k = \mathbf{1}$ corresponds to $x = 0$. Hence A is valid iff Λ has no nonzero vector with all entries ≥ -1. Since $\Lambda = -\Lambda$, this holds iff Λ has no nonzero vector with all entries ≥ -1 and none with all entries ≤ 1, which is the statement. □

Corollary 4 (PROVED)**.** *(a) Validity is invariant under $x \mapsto ux + v$ with u a unit of $\mathbb{Z}_N$. (b) Every subset of a valid set is valid (as a set of its own size, same modulus). (c) If A lies in a coset of the subgroup of index h, then A, read in $\mathbb{Z}_{N/h}$, is valid mod N/h. (d) (Inal's bound.) For $c \in A$ the 2^{n-1} sums $\sum_{a \in S}(a - c)$, $S \subseteq A \setminus \{c\}$, are distinct; hence $N \ge 2^{n-1}$.*

Proof. (a) and (c): the lattice Λ is unchanged. (b): extend a relation of the subset by zeros. (d): if two such sums agree for $S \ne T$, $|S| \le |T|$, then $\mathbf{1}_T - \mathbf{1}_S - (|T| - |S|)e_c$ is a nonzero vector of Λ with all entries ≤ 1. □

Part (d) uses only relation vectors with at most one entry outside $\{-1, 0, 1\}$. This suggests:

Definition 5. *A* is *all-anchor dissociated* (AAD) mod N if every nonzero $x \in \Lambda$ has at least two entries of absolute value ≥ 2; equivalently, for every $c \in A$ the 2^{n-1} sums in Corollary 4(d) are distinct.

Valid sets are AAD by Proposition 3. The converse fails (Section 5).

4 The cyclic-order lemma

Fix distinct $a, c \in A$, put $b = a - c$ and $A'' = A \setminus \{a, c\}$, $m = n - 2$. For $S \subseteq A''$ let $x_S = \sum_{e \in S}(e - c) \in \mathbb{Z}_N$ and $w(S) = |S|$.

Lemma 6 (PROVED)**.** *Let A be valid mod N. Let $S \neq S'$ be subsets of A'' with $w(S') \geq w(S)$, and suppose $x_{S'} - x_S = i\,b$ for an integer i. Then $i \notin [-1,\ w(S') - w(S) + 1]$.*

Proof. Define $\lambda \in \mathbb{Z}^A$ by $\lambda|_{A''} = \mathbf{1}_{S'} - \mathbf{1}_S$, $\lambda_a = -i$, $\lambda_c = i - (w(S') - w(S))$. Then $\sum \lambda = 0$ and $\sum_e \lambda_e e = x_{S'} - x_S - i(a - c) + c\sum \lambda = 0$ in $\mathbb{Z}_N$, so $\lambda \in \Lambda$, and $\lambda \neq 0$ because $S \neq S'$. If $-1 \leq i \leq w(S') - w(S) + 1$, then $\lambda_a \leq 1$, $\lambda_c \leq 1$ and the entries on A'' lie in $\{-1, 0, 1\}$, so λ has no entry ≥ 2, contradicting Proposition 3. □

Assume now that b is a unit of $\mathbb{Z}_N$, and define the *position* $p(S) \in \{0, \ldots, N - 1\}$ by $x_S = p(S)\,b$. Applying the lemma to all integers $i \equiv p(S') - p(S)$:

Corollary 7 (PROVED)**.** *If b is a unit, then for $S \neq S'$ with $w(S') \geq w(S)$ the forward distance $d = (p(S') - p(S)) \bmod N$ satisfies*

$$w(S') - w(S) + 2 \ \leq\ d \ \leq\ N - 2.$$

In particular the 2^m positions are distinct and pairwise at cyclic distance at least 2; listing them in cyclic order as $p_1 < \cdots < p_M$ ($M = 2^m$) with weights w_j, the gaps $g_j = p_{j+1} - p_j$ (indices mod M, $g_M = N - p_M + p_1$) satisfy $g_j \geq 2 + \max(0, w_{j+1} - w_j)$. Equivalently the sheared positions $p(S) - w(S)$ are again pairwise 2-separated and occur in the same cyclic order.

The sheared configuration is the same structure seen from the anchor a instead of c. The corollary is where cyclicity enters: in $(\mathbb{Z}_2)^{n-1}$ every difference has order 2 and nothing is gained.

Theorem 8 (PROVED)**.** *Let A be valid mod N, $n \geq 3$, and suppose some difference of two elements of A is a unit of $\mathbb{Z}_N$ (automatic if N is prime). Then $N \geq 2^{n-1} + n - 2$.*

Proof. Summing the gap inequalities of Corollary 7 around the cycle gives $N = \sum_j g_j \geq 2M + \sum_j \max(0, w_{j+1} - w_j)$. The cyclic sequence (w_j) takes the values 0 (for $S = \emptyset$) and m (for $S = A''$), so its total positive variation is at least $m = n - 2$. □

Proposition 9 (PROVED)**.** *Let A be valid mod N, $n \geq 3$, let $a, c \in A$ with $a - c$ a unit, and suppose A contains $e \in \{3a - 2c,\ 4a - 3c,\ 3c - 2a,\ 4c - 3a\}$, $e \neq a, c$. Then $N \geq 5 \cdot 2^{n-3}$.*

Proof. Exchanging the roles of a and c we may assume $e - c = g\,b$ with $g \in \{3, 4\}$. Let $P_0 = \{p(S) : e \notin S\}$, so $|P_0| = M/2$ and $p(S \cup \{e\}) \equiv p(S) + g$. Let $x \in P_0$ have weight k, and suppose a position y of some T lies strictly between x and $x + g$. By 2-separation $y = x + 2$ and $g = 4$. If $w(T) \geq k$, Corollary 7 gives $2 \geq w(T) - k + 2$, so $w(T) = k$, and then the forward distance 2 from y to the heavier point $x + g$ contradicts the lower bound 3. If $w(T) < k$, the forward distance 2 from y to $x + g$ must be at least $(k + 1 - w(T)) + 2 \geq 4$, a contradiction. So the $M/2$ arcs $[x, x + g]$, $x \in P_0$, contain no further positions; they are pairwise disjoint, and the first position after $x + g$ is at distance at least 2. Each arc therefore accounts for at least $g + 2 \geq 5$ residues, and $N \geq 5M/2 = 5 \cdot 2^{n-3}$. □

The set $\{1, 2, 4, \ldots, 2^{n-1}\}$ satisfies the hypothesis with $(c, a, e) = (1, 2, 4)$ whenever 1 is a unit, so Proposition 9 applies to the extremal family but gives only $5/8$ of the true value.

The abstract one-dimensional problem. Call $(N; \beta_1, \ldots, \beta_m)$ a *structure* if the points $p(S) = \sum_{l \in S} \beta_l \bmod N$ satisfy the inequalities of Corollary 7.

Conjecture 10 (VERIFIED for $m \le 5$ by exhaustive search)**.** *Every structure with m generators has $N \ge 5 \cdot 2^{m-1}$.*

The bound is attained by $\beta = (3, 5, 10, 20, \ldots, 5 \cdot 2^{m-2})$; for $m \ge 4$ there are further extremal structures, e.g. $(3, 5, 20, 30)$ modulo 40. The map $\beta_l \mapsto N+1-\beta_l$ (exchange of a and c) preserves structures.

The next three statements make progress on Conjecture 10. For a structure list the points in cyclic order $S_0 = \emptyset, S_1, \ldots, S_{M-1}$, with gaps g_j, step vectors $\delta_j = \mathbf{1}_{S_{j+1}} - \mathbf{1}_{S_j} \in \{-1, 0, 1\}^m$ and $d_j = w(S_{j+1}) - w(S_j)$; let $U = \sum_j \max(0, d_j)$. For a step vector δ a *translate* of the step is a pair $(T, T+\delta)$ of subsets with $\mathbf{1}_{T+\delta} = \mathbf{1}_T + \delta$.

Proposition 11 (PROVED)**.** *For each step j let n_j^* be the largest number of points in a half-open forward arc $[T, T+\delta_j)$ over all translates of the step. Then $g_j \ge 2n_j^* + \max(0, d_j)$ and*

$$N \;\ge\; 2^{m+1} + U + 2\sum_j (n_j^* - 1) \;\ge\; 2^{m+1} + U + 2L,$$

where L is the number of steps having a translate that is not a pair of consecutive points.

Proof. The forward distance from T to $T+\delta_j$ is the residue of $\sum_l (\delta_j)_l \beta_l$, which does not depend on T and equals g_j. An arc $[T, T+\delta_j)$ containing n points consists of n consecutive steps, each of length at least 2 plus its ascent, and the ascents of these steps add up to at least $\max(0, d_j)$ because the weights of T and $T+\delta_j$ differ by d_j. Sum over j. □

For $(5, 7, 9)$ modulo 23 the proposition gives $16 + 3 + 2 \cdot 2 = 23$, which is exact.

Proposition 12 (PROVED)**.** *If for some generator l the cyclic order alternates between points not containing l and points containing l, then $U \ge 2^{m-1}$ and $N \ge 5 \cdot 2^{m-1}$.*

Proof. There are 2^{m-1} steps from a point x without l to a point y with l, and y runs over all points containing l. With $\omega(S) = |S \setminus \{l\}|$ the sum of d_j over these steps is $\sum_y (\omega(y) + 1) - \sum_x \omega(x) = 2^{m-1}$, since $S \mapsto S \cup \{l\}$ preserves ω. Hence $U \ge 2^{m-1}$ and $N \ge 2M + U$. □

Proposition 9 is the special case $\beta_l \in \{3, 4\}$, in which the arcs $[x, x+\beta_l]$ are empty and the order alternates.

Proposition 13 (PROVED)**.** *Conjecture 10 holds for $m \le 3$. Consequently, a valid n-set with a unit difference has $N \ge 5 \cdot 2^{n-3}$ for $n \le 5$.*

Proof. Write $U = \sum_{h=0}^{m-1} c_h$, where $c_h \ge 1$ counts the steps from weight $\le h$ to weight $> h$. For $m \le 2$ this gives $U \ge m = 2^{m-1}$. Let $m = 3$. If $U \ge 4$ we are done, so let $U = 3$; then every $c_h = 1$, so for each h the points of weight $\le h$ form one cyclic interval. By Proposition 11 it suffices to find one step with a non-consecutive translate ($U + 2L \ge 5 > 4$). Suppose every translate of every step is a pair of consecutive points, and let S_1 be the successor of $\emptyset$, $k = |S_1|$. If $k = 1$, $S_1 = \{l\}$, then all four pairs $(T, T \cup \{l\})$ are consecutive ascents and $U \ge 4$. If $k = 2$, $S_1 = \{p, q\}$, then $\{r\} \to \{p, q, r\}$ is consecutive for the third generator r, so both $\emptyset$ and $\{r\}$ are followed by a point of weight ≥ 2 and both are the last point of the interval of weight ≤ 1, which is impossible. If $k = 3$, then $\emptyset$ is the last point of the interval of weight ≤ 1, which has four points, so its predecessor is a singleton $\{l\}$; the step $\{l\} \to \emptyset$ has the translates $\{l, h\} \to \{h\}$ for both $h \ne l$, so two distinct singletons are each preceded by a point of weight 2 and are both the first point of that interval, which is impossible. The last sentence follows from Corollary 7 with $m = n - 2$. □

By Proposition 11, Conjecture 10 follows from the inequality $U + 2L \geq 2^{m-1}$ for every structure; this inequality was checked on all structures with $m = 3$, $20 \leq N \leq 30$, with $m = 4$, $40 \leq N \leq 54$, and with $m = 5$, $N = 80$ (VERIFIED, small search by a second program), and is open in general.

If true this gives $N \geq 5 \cdot 2^{n-3}$ for every valid set with a unit difference. The computed minima are $5, 10, 20, 40, 80$ for $m = 1, \ldots, 5$ (program `oned.c`; the search enumerates all increasing generator tuples and prunes on sub-cubes). A related computation (`lk.c`, VERIFIED for the stated sizes): if the sign condition of Proposition 3 is imposed only on relation vectors whose entries outside a fixed set of k coordinates lie in $\{-1, 0, 1\}$, the least modulus is 2^{n-1} for $k \leq 2$, equals $6, 10, 20, 40$ for $k = 3$, $n = 3, \ldots, 6$, and equals $12, 24, 46$ for $k = 4$, $n = 4, 5, 6$. The inequality $N \geq 2M +$ (positive variation) alone does not suffice: $(5, 7, 9)$ modulo 23 is a structure whose weights in cyclic order are $0, 1, 1, 1, 2, 2, 2, 3$.

5 Computations

5.1 Method and controls

`enum3.c` enumerates valid n-sets modulo N up to the affine maps $x \mapsto ux + v$. Normal form: for a divisor $d < N$ of N the set contains 0 and d, and every pairwise difference $x - y$ has $\gcd(x - y, N) \geq d$. Every valid set has such an image: choose a pair whose difference has minimal gcd with N, translate one element to 0 and scale the other to that gcd (the units act transitively on residues of a given gcd). For $d = 1$ a second reduction is used: among all triples (c, a, e) with $a - c$ a unit, pick one minimizing the residue $(e - c)(a - c)^{-1} \in [2, N - 1]$ and normalize $(c, a) \mapsto (0, 1)$; then the third smallest element t of the normalized set is a lower bound for all these residues, a condition inherited by subsets containing $\{0, 1, t\}$, so it can be used for pruning. Elements are added in increasing order and every prefix must be valid (Corollary 4(b)). Validity of a k-set is tested by a bitset dynamic program that counts, capped at two, the multisets of size k with the target sum.

Controls. (1) Positive controls: the search finds valid sets at every $N = 2^n - 2^k$ with $2^k \leq n$ for $n \leq 8$ (Table 1), so a "no set found" answer is not an artifact of over-pruning at those sizes. (2) The pruned program `enum3` and the unpruned programs `enum2` (only the gcd normal form) and `enum` (only $0 \in A$) give the same number of affine classes for all (n, N) in Table 1 with $n \leq 7$ and for $N = 123$, $n = 7$ (zero). (3) An independent pure-Python checker (`check_valid.py`) accepts every set output by the C programs and rejects planted invalid sets ($\{1, 2, 4, 8\}$ mod 13, $\{1, 2, 4, 8, 16\}$ mod 24, $\{0, 1, 2\}$ mod 5). (4) The order-lemma checker (`order_check.py`) reports 0 violations on all valid sets found for $n \leq 7$, and reports violations on 192 of the 480 sets modulo 60 that are AAD but not valid, so the test can fail.

5.2 Results

Theorem 14 (VERIFIED, exhaustive search, $n \leq 8$, all $N < 2^n$)**.** *For $3 \leq n \leq 8$ and $N < 2^n$ there is a valid n-set modulo N if and only if $N = 2^n - 2^k$ with $2^k \leq n$. In particular Conjecture 2 holds for $n \leq 8$.*

For $N < 2^{n-1}$ this is Corollary 4(d); the search covers $2^{n-1} \leq N < 2^n$.

Remark 15 (a sporadic class; VERIFIED)**.** Modulo 15 the set $\{0, 1, 3, 10\}$ is valid and is not affinely equivalent to $\{1, 2, 4, 8\}$ (the latter is equivalent to $\{0, 1, 3, 7\}$). No translate-and-scale of it is closed under $x \mapsto 2x - t$ for any t, so it is not a "doubling orbit". For $n \in \{5, 6, 7, 8\}$ and $N = 2^n - 1$ the search finds a single class. This does not contradict the uniqueness question of [1], which concerns the minimal modulus $2^n - 2^{\lfloor \log_2 n \rfloor}$ ($= 12$ for $n = 4$), where the class is unique for $n \leq 8$.

n	moduli $2^{n-1} \le N < 2^n$ with a valid n-set	affine classes at these moduli
3	6, 7	1, 1
4	12, 14, 15	1, 2, 2
5	28, 30, 31	1, 5, 1
6	60, 62, 63	1, 2, 1
7	124, 126, 127	1, 1, 1
8	248, 252, 254, 255	1, 1, 2, 1

Table 1: VERIFIED. Output of `enum3` for every N in $[2^{n-1}, 2^n)$; all moduli not listed returned no valid set. For $n = 8$ the run visited between $6.6{\cdot}10^4$ and $5.3{\cdot}10^6$ search nodes per modulus (log `logs/n8.log`); the six normal-form sets found at $N \in \{248, 252, 254, 255\}$ pass the independent checker. Second class at $N = 254$: $\{0, 1, 64, 96, 112, 120, 124, 126\}$.

Remark 16 (AAD sets; VERIFIED)**.** For $3 \le n \le 6$ the moduli $N < 2^n$ admitting an AAD n-set are the same as for valid sets. For $n = 7$ they are $N \in \{120, 124, 126, 127\}$ (program `enumd.c`, all $64 \le N < 128$), so the AAD threshold is $120 = 2^7 - 2^3$, below the validity threshold 124. A proof of Conjecture 2 must therefore use relation vectors with at least two entries outside $\{-1, 0, 1\}$.

Conjecture 17 (refined form; VERIFIED for $n \le 8$)**.** *If A is a valid n-set modulo N and $N < 2^n$, then N divides $2^n - 2^k$ for some k with $2^k \le n$ (hence $N = 2^n - 2^k$ by Corollary 4(d) and $N > 2^{n-1}$).*

In particular the conjecture predicts that an *odd* modulus below $2^n - 1$ never carries a valid n-set.

6 What is not established

1. Conjecture 2 for $n \ge 9$, and any bound of the form $N \ge (\frac{1}{2} + \varepsilon)2^n$ valid for all valid sets. Theorem 8 gains only an additive $n - 2$ and needs a unit difference; Proposition 9 needs an arithmetic hypothesis.
2. Conjecture 10 beyond $m = 5$. Even if proved it yields the constant $5/8$, not 1.
3. The verification for $n = 8$ is a computation, not a proof checked by a kernel; no Lean formalization of the new statements was produced (no Lean toolchain was available in the session). The formal statement of Conjecture 2 exists in [3].
4. We did not read Inal's preprint or the full text of [1] in PDF form inside the session (the arXiv PDF could be read only through a summarizing fetch tool); attributions to them follow the arXiv abstract, the README of [2] and [4].

A Dead ideas and their obstructions

1. *Clique (pigeonhole) arguments.* Any family of sum-zero integer vectors whose pairwise differences are forbidden relation vectors has at most 2^{n-1} members, because $(\mathbb{Z}_2)^{n-1}$ carries a valid tuple. The same caps every Delsarte-type linear-programming bound that ignores cyclicity.
2. *Two translated cubes.* $N \ge 2^n - |Q_c \cap (Q_c + g)|$ for the anchored cube Q_c and any g; averaging over g returns $(N - 2^{n-1})^2 \ge 0$. For explicit valid sets the minimum overlap exceeds $2^n - N$ (e.g. 4 versus 1 for $\{1, 2, 4, 8, 16\}$ mod 31), also for dilates $uQ_c + g$ and for cubes at two different anchors.

3. *A virtual anchor.* For $\{1, 2, 4, \dots\}$ the 2^n raw subset sums cover $\mathbb{Z}_N$; for the sporadic set mod 15 no translation t makes the map $S \mapsto \sum_{a \in S}(a - t)$ hit more than 8 residues.

4. *Character and polynomial identities.* Parseval for the Riesz product $\prod_{a \neq c}(1 + \cos(2\pi r(a - c)/N))$, the norm of $\prod(1 + \zeta^{a-c})$, and the degree-$(p-1)$ moment identity modulo a prime p all reduce to the statement "the number of relations is one" and give no bound beyond 2^{n-1}.

5. *Counting with all translations.* Summing collisions over all $t \in \mathbb{Z}_N$ gives the known bound $\binom{2n}{n}/2^n$.

6. *Reduction to the AAD property.* False for $n = 7$ ($N = 120$).

7. *Parity induction for even N.* The two parity classes are valid modulo $N/2$, which gives $N \geq 2f(\max(n_0, n_1))$; nothing forces the split $(n - 1, 1)$ seen in all examples.

8. *Doubling the smallest element.* For $\{1, 2, 4, \dots\}$ anchored at 1 the complement of Q_c is $\{x_S + b_{\min S}\}$; for a general valid set these points can coincide with each other and with lighter points of Q_c, and no choice rule avoids all allowed coincidences.

9. *Proving Conjecture 10 by sub-cube doubling or by counting gap-2 steps.* The structure $(5, 7, 9)$ mod 23 has no generator whose removal leaves a structure that fits in half the cycle by an additive map, and flat gap-2 steps (equal weights at distance 2) are compensated only globally.

References

[1] J. A. R. Fonollosa, *Minimum modulus for the unique multiset-sum problem*, arXiv:2607.08366 (v2, 14 July 2026).

[2] J. A. R. Fonollosa, `github.com/jarfo/min-modulus` (Lean 4 formalization; state of 24 July 2026), including a formalization of the abelian-group bound credited there to M. Inal, *The Exact Minimum Order for Unique Multiset Sums in Finite Abelian Groups*, OSF preprint, DOI 10.17605/OSF.IO/C58Q9 (2026).

[3] Google DeepMind, *Formal Conjectures*, `github.com/google-deepmind/formal-conjectures`, file `FormalConjectures/Arxiv/2607.08366/MinModulus.lean` (state of 9 November 2026).

[4] `github.com/the-omega-institute/trureturing`, issue #20156, "Research: Fonollosa Conjecture 1 — global minimum modulus for unique multiset sums" (opened 22 October 2026).