

Sem-I \rightarrow complex Nos.

$$Z = \underset{\substack{\uparrow \\ \text{Real}}}{a} + i \underset{\substack{\downarrow \\ \text{Im.}}}{b}, \quad a, b \in \mathbb{R}$$

GATE

Sem-III complex variables : derivative of complex variable.
 Sem-IV complex Integration :-

Complex variables

$$Z = x + iy, \quad x, y \in \mathbb{R}$$

$$Z = f(x, y)$$

$$|Z| = \sqrt{x^2 + y^2}, \quad \theta = \tan^{-1} \left[\frac{\text{Im}}{\text{Re}} \right] = \tan^{-1} \left[\frac{y}{x} \right]$$

$$\text{Let } w = f(z) = Z^2 = (x + iy)^2 = x^2 + 2ixy + i^2 y^2 \quad \because i^2 = -1$$

$$= (x^2 - y^2) + i(2xy)$$

$$= u(x, y) + i v(x, y)$$

$$\boxed{w = f(z) = u + iv} \rightarrow \text{C.V.}$$

$$Z = x + iy \rightarrow \text{C.V.}$$

$$\text{Modulus of } f(z) = |f(z)| = \sqrt{u^2 + v^2}$$

$$\text{Argument} = \text{Amplitude} = \theta = \tan^{-1} \left[\frac{\text{Im}}{\text{Re}} \right] = \tan^{-1} \left[\frac{v}{u} \right]$$


Defⁿ: Analytic Function / Regular Function / Entire Fun /
 Holomorphic Function.

IF $f(z) = u + iv$ is differentiable at each & every point
 in given Region of convergence (ROC), Then $f(z) = u + iv$
 is called as an Analytic Function. $z = 2 + 3i = (2, 3)$

$$\text{Ex. } f(z) = \frac{z^3 - 4z^2 + 3z - 7}{(z-2)(z+3)(z-7)(z+10)}$$

$$f(z) = \infty = \frac{\quad}{\quad}$$

$$\boxed{z = 2, -3, 7, -10}$$



Cauchy's Riemann equl's [C.R. equl's]

IF $f(z) = u + iv$ is an Analytic Fun and $\frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}, \frac{\partial v}{\partial x}, \frac{\partial v}{\partial y}$ are exists, Then

$$\textcircled{i} \quad \boxed{\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}} \quad \text{or} \quad \boxed{U_x = V_y}$$

$$\textcircled{ii} \quad \boxed{\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}} \quad \text{or} \quad \boxed{U_y = -V_x}$$

These conditions (equl's) are called Cauchy's Riemann equl's or C.R. equl's

Analytic \longleftrightarrow C.R. equl exists

II] C.R equations in polar form:-

IF $f(z) = u + iv = u(r, \theta) + iv(r, \theta)$ is an Analytic Fun

and $\frac{\partial u}{\partial r}, \frac{\partial u}{\partial \theta}, \frac{\partial v}{\partial r}, \frac{\partial v}{\partial \theta}$ are exists, Then

C.R. equl's
in
polar form

$$\textcircled{i} \quad \boxed{\frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial v}{\partial \theta}} \quad \text{or} \quad \boxed{U_r = \frac{1}{r} V_\theta}$$

$$\textcircled{ii} \quad \boxed{\frac{\partial v}{\partial r} = -\frac{1}{r} \frac{\partial u}{\partial \theta}} \quad \text{or} \quad \boxed{V_r = -\frac{1}{r} U_\theta}$$

Ex-① $f(z) = z^2$ Is an Analytic Fun or Not?

→ Step ① $f(z) = z^2 = (x + iy)^2 = (x^2 - y^2) + i(2xy)$

$f(z) = z^2 = u + iv$

where $u = x^2 - y^2$

and $v = 2xy$

step ② $\frac{\partial u}{\partial x} = 2x - 0 = 2x$

; $\frac{\partial v}{\partial x} = 2y$

$\frac{\partial u}{\partial y} = 0 - 2y = -2y$

; $\frac{\partial v}{\partial y} = 2x$

$$\frac{\partial u}{\partial y} = 0 - 2y = -2y \quad ; \quad \frac{\partial v}{\partial y} = 2x$$

step (III) $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} = \boxed{2x} \Rightarrow \boxed{\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}}$
 $\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x} = \boxed{-2y} \Rightarrow \boxed{\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}}$ } C.R. eqns are exist.

Then $f(z) = z^2$ is an Analytic Fun.

1 Find a, b, c, d, e If $f(z) = (ax^3 + bxy^2 + 3x^2 + cy^2 + x) + i(dx^2y - 2y^3 + exy + y)$ is an analytic function?

→ We have $f(z) = u + iv$ is an Analytic Fun (given)

✓ $u = ax^3 + bxy^2 + 3x^2 + cy^2 + x$ — (i) } given

✓ $v = dx^2y - 2y^3 + exy + y$ — (ii)

C.R. eqns are exist.

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \text{--- (iii)}$$

$$\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x} \quad \text{--- (iv)}$$

$$\begin{cases} \frac{\partial u}{\partial x} = 3ax^2 + by^2 + 6x + 0 + 1 \\ \frac{\partial v}{\partial y} = dx^2 - 6y^2 + ex + 1 \end{cases}$$

But by eqn (iii)

$$3ax^2 + by^2 + 6x + 1 = dx^2 - 6y^2 + ex + 1$$

$$\Rightarrow \boxed{3a = d}$$

$$\therefore \boxed{b = -6}$$

$$\Rightarrow \boxed{b = -6}$$

$$\therefore \boxed{e = 6}$$

$$\Rightarrow \boxed{6 = e}$$

From eqn (i) & eqn (ii)

$$\frac{\partial u}{\partial y} = 2bxy + 2cy$$

$$\frac{\partial v}{\partial x} = 2dxy + ey$$

By eqn (iv) $\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$

$$(2bxy + 2cy) = -(2dxy + ey)$$

$$\Rightarrow 2b = -2d \Rightarrow \boxed{b = -d}$$

$$\Rightarrow -6 = -d \Rightarrow \boxed{d = 6}$$

$$\Rightarrow 2c = -e \Rightarrow 2c = -6 \Rightarrow \boxed{c = -3}$$

We have $3a = d$
 $3a = 6$
 $\Rightarrow \boxed{a = 2}$

$$\therefore \boxed{\text{Ans: } a=2, b=-6, c=-3, d=6, e=6}$$

2. Find the constants k , if $f(z) = \frac{1}{2} \log(x^2 + y^2) + i \tan^{-1} \frac{kx}{y}$ is an Analytic function.

→ We have $f(z) = u + iv \Rightarrow$ Analytic Fun (given)
 Then C.R. eqns are exist.

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \text{--- (i)}$$

$$\boxed{\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}} \quad \text{--- (ii)}$$

$$u = \frac{1}{2} \log(x^2 + y^2) \quad \text{--- (iii)}$$

$$v = \tan^{-1} \left(\frac{kx}{y} \right) \quad \text{--- (iv)}$$

$$\frac{\partial u}{\partial x} = \frac{1}{2} \left(\frac{1}{x^2 + y^2} \right) \times (2x + 0) = \frac{x}{x^2 + y^2} \quad \text{--- (v)}$$

$$\frac{\partial v}{\partial y} = \frac{1}{\left[1 + \left(\frac{kx}{y} \right)^2 \right]} \times \left[\frac{kx(-1)}{y^2} \right] = \frac{-kx}{\left[\frac{y^2 + k^2 x^2}{y^2} \right] y^2} = \frac{-kx}{k^2 x^2 + y^2} \quad \text{--- (vi)}$$

By eqn (i) $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$

$$\frac{x}{x^2 + y^2} = \frac{-kx}{k^2 x^2 + y^2}$$

$$\Rightarrow \boxed{k = -1}$$

Dr. Uday Kashid (PhD, Mathematics)

4. Show that $f(z) = e^{2z} - z$ is an analytic function ? [Biom-Dec-19, (5M)]

→ We have $f(z) = e^{2z} - z = e^{2(x+iy)} - (x+iy)$
 $2x \quad i(2y)$

$$= e \cdot e^{-x-iy}$$

$$= e^{2x} [\cos(2y) + i \sin(2y)] - x - iy$$

$$\therefore e^{i\theta} = \cos\theta + i\sin\theta$$

$$f(z) = [e^{2x} \cos(2y) - x] + i[e^{2x} \sin(2y) - y] = u + iv$$

$$\therefore u = e^{2x} \cos(2y) - x \quad \text{--- (i)}$$

$$v = e^{2x} \sin(2y) - y \quad \text{--- (ii)}$$

$$\frac{\partial u}{\partial x} = 2e^{2x} \cos(2y) - 1$$

$$\frac{\partial v}{\partial y} = 2e^{2x} \cos(2y) - 1$$

$$\frac{\partial u}{\partial y} = -2e^{2x} \sin(2y) - 0$$

$$\frac{\partial v}{\partial x} = 2e^{2x} \sin(2y) - 0$$

$$\Rightarrow \boxed{\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}}$$

$$\Rightarrow \boxed{\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}}$$

C.R. eqns are exist
Hence
 $f(z)$ is an Analytic fun.

4. Prove that an Analytic function with constant modulus is constant function. [Dec-08]

→ We have $f(z) = u + iv$ is an Analytic Function. Then
C.R eqns are always exist.

$$\boxed{\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}} \quad \text{--- (i)}$$

$$\boxed{\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}} \quad \text{--- (ii)}$$

$$|f(z)| = \sqrt{u^2 + v^2} = \text{constant} = k \text{ (say)}$$

$$u^2 + v^2 = k^2 \quad \text{--- (iii)}$$

diff eqn (iii) w.r.t x

$$2u \frac{\partial u}{\partial x} + 2v \frac{\partial v}{\partial x} = 0$$

$$u \frac{\partial u}{\partial x} + v \frac{\partial v}{\partial x} = 0 \quad \text{--- (iv)}$$

diff eqn (iii) w.r.t y

$$2u \frac{\partial u}{\partial y} + 2v \frac{\partial v}{\partial y} = 0$$

$$\Rightarrow u \frac{\partial u}{\partial y} + v \frac{\partial v}{\partial y} = 0 \quad \text{--- (v)}$$

By C.R. eqns from eqn (i) & (ii)

$$-u \frac{\partial v}{\partial x} + v \frac{\partial u}{\partial x} = 0 \quad \text{--- (vi)}$$

$$v \frac{\partial v}{\partial x} + u \frac{\partial u}{\partial x} = 0 \quad \text{--- (vii) [copied]}$$

$u = \text{const}$
∴ derivative $u = 0$
 $\frac{\partial u}{\partial x} = 0, \frac{\partial u}{\partial y} = 0$

$$v \frac{\partial v}{\partial x} + u \frac{\partial u}{\partial x} = 0 \quad \text{--- (iv) [copied]}$$

Multiply eqⁿ (vi) by v and multiply eqⁿ (iv) by u and Addition

$$\begin{array}{r} -uv \frac{\partial v}{\partial x} + v^2 \frac{\partial u}{\partial x} = 0 \\ + uv \frac{\partial v}{\partial x} + u^2 \frac{\partial u}{\partial x} = 0 \\ \hline 0 + (u^2 + v^2) \frac{\partial u}{\partial x} = 0 \end{array}$$

$$\begin{aligned} \Rightarrow (u^2 + v^2) \frac{\partial u}{\partial x} &= 0 \\ K^2 \frac{\partial u}{\partial x} &= 0 \\ \Rightarrow \boxed{\frac{\partial u}{\partial x} = 0} &\text{--- (A)} \end{aligned}$$

But by C.R eqⁿ $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} = 0$

$$\boxed{\frac{\partial v}{\partial y} = 0} \quad \text{--- (B)}$$

Dr. Uday Kashid (PhD, Mathematics)

Multiply eqⁿ (vi) by u and eqⁿ (iv) by v and Subtraction

$$-u^2 \frac{\partial v}{\partial x} + uv \frac{\partial u}{\partial x} = 0$$

$$-v^2 \frac{\partial v}{\partial x} + uv \frac{\partial u}{\partial x} = 0$$

$$-(u^2 + v^2) \frac{\partial v}{\partial x} + 0 = 0$$

$$\Rightarrow -K^2 \frac{\partial v}{\partial x} = 0 \quad \Rightarrow \boxed{\frac{\partial v}{\partial x} = 0} \quad \text{--- (C)}$$

But C.R eqⁿ $\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$

$$\boxed{\frac{\partial u}{\partial y} = -0 = 0} \quad \text{--- (D)}$$

By eqⁿ (A), (B), (C), & (D)

$$\frac{\partial u}{\partial x} = 0$$

$$\frac{\partial u}{\partial y} = 0$$

$$\frac{\partial v}{\partial x} = 0$$

$$\frac{\partial v}{\partial y} = 0$$

$$\boxed{u = \text{constant}}$$

$$\boxed{v = \text{constant}}$$

$$f(z) = u + iv = \text{const.}$$

Orthogonal Trajectory:

Dr. Uday Kashid (PhD, Mathematics)

Orthogonal Trajectory: In mathematics, if every member of one family of curves that intersect to each member of another family of curves at right angles (see figure) is called as Orthogonal trajectory in between both family of curves.




Fig.1. Family of concentric Circles and Straight lines passing through origin.




Fig.2. Family of ellipse and parabola

Theorem:: If $f(z) = (u + i v)$ is an Analytical function then the curves $u(x, y) = C_1$ and $v(x, y) = C_2$ represents family of orthogonal trajectories to each other.

Milne-Thompson Method:- (MTM)

Q. IF we know $u = u(x, y)$ Then How to find an Analytic Funⁿ $f(z) = u + i v$ $\rightarrow u = \text{given}$

step (I) $f(z) = u + i v \rightarrow$ Analytic Funⁿ and $u = \text{given}$

step (II) diff w.r.t. x ; $f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$

step (III) By C.R eqns $\left[\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x} \right]$

$$f'(z) = \left[\frac{\partial u}{\partial x} \right] - i \left[\frac{\partial u}{\partial y} \right]$$

$$\checkmark f'(z) = \phi_1(x, y) - i \phi_2(x, y)$$

$$u = x^2 + 2xy + y^2$$

$$\frac{\partial u}{\partial x} = 2x + 2y = \phi_1(x, y)$$

$$\frac{\partial u}{\partial y} = 0 + 2x + 2y = \phi_2(x, y)$$

step (IV) By MTM, put $x = z$ and $y = 0$

$$f'(x + i y) = \phi_1(x, y) - i \phi_2(x, y)$$

$$f'(z + i 0) = \phi_1(z, 0) - i \phi_2(z, 0)$$

$$f'(z) = \phi_1(z) - i \phi_2(z)$$

step (V) Integrate w.r.t. z

Milne-Thompson Th^m
IF we put $x = z$ and $y = 0$ in derivative part
Then, value of eqn is Not change.
7/11/15

step (v) Integrate w.r.t. z

$$\int f'(z) dz = \int \phi_1(z) dz - i \int \phi_2(z) dz + C$$

$$f(z) = \psi_1(z) - i \psi_2(z) + C \rightarrow \text{Analytic}$$

Dr. Uday Kashid (PhD, Mathematics)

Case-II] If $V = V(x, y) = \text{given}$

step (i) $f(z) = u + iv \rightarrow \text{Analytic Fun}$

$$(i) f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

step (ii) By C.R eqn $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$

$$f'(z) = \frac{\partial v}{\partial y} + i \frac{\partial v}{\partial x}$$

$$f'(z) = f'(x+iy) = \phi_1(x, y) + i \phi_2(x, y)$$

step (iii) By MTM, put $x = z$ and $y = 0$

$$f'(z+i0) = \phi_1(z, 0) + i \phi_2(z, 0)$$

step (iv) Integrating w.r.t z

$$\int f'(z) dz = \int \phi_1(z) dz + i \int \phi_2(z) dz + C$$

$$f(z) = \psi_1(z) + i \psi_2(z) + C \rightarrow \text{Analytic}$$

Dr. Uday Kashid (PhD, Mathematics)

Ex.1 Construct an Analytic function $f(z) = (u + iv)$ If $u = e^x \cos(y)$ [EXTC-May-19, (6M)]

\rightarrow step (i) we have $f(z) = u + iv \rightarrow \text{Analytic Fun}$

$$\text{step (ii) diff. w.r.t. } x; f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

step (iii) By C.R eqn $\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$

$$f'(z) = \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y}$$

But we have $u = e^x \cos y$

$$\frac{\partial u}{\partial x} = e^x \cos y \quad \text{and} \quad \frac{\partial u}{\partial y} = -e^x \sin y$$

$$f'(z) = (e^x \cos y) - i(-e^x \sin y)$$

step-iv) By Milne-Thompson method, put $x=z$ and $y=0$

$$f'(x+iy) = (e^x \cos y)_{\substack{x=z \\ y=0}} + i(e^x \sin y)_{\substack{x=z \\ y=0}}$$

$$f'(z+io) = e^z \cos(0) + i(e^z \sin(0))$$

$$f'(z) = e^z(1) + 0 = e^z$$

step-v) Integrate w.r.t z

$$\int f'(z) dz = \int e^z dz + C$$

$$\boxed{f(z) = e^z + C}$$

→

$$f(z) = e^{x+iy} + C = e^x \cdot e^{iy} + C = e^x [\cos y + i \sin y] + C$$

$$= e^x \cos y + i e^x \sin y + i\alpha$$

$$\text{let } \boxed{C = i\alpha}$$

$$= (e^x \cos y) + i(e^x \sin y + \alpha)$$

$$= u + iv$$

$$\Rightarrow u = e^x \cos y$$

$$\boxed{v = e^x \sin y + \alpha}$$

→ { Harmonic conjugate of u
orthogonal trajectories of u

Ex.2 Construct an Analytic function $f(z)$ if $v = e^x [x \sin(y) + y \cos(y)] = c$ [BIOM-Nov-18, (6M)]

→ step-i) We have $f(z) = u + iv \rightarrow$ Analytic fun

$$\textcircled{ii} f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

$$\textcircled{iii} \text{ By C.R eqn } \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$$

Dr. Uday Kashid (PhD, Mathematics)

$$f'(z) = \frac{\partial v}{\partial y} + i \frac{\partial v}{\partial x}$$

$$\text{We have } v = e^x [x \sin y + y \cos y] - c = 0$$

$$\frac{\partial v}{\partial x} = e^x [x \sin y + y \cos y] + e^x [\sin y + 0] - 0$$

$$= e^x [x \sin y + y \cos y + \sin y]$$

$$\checkmark \frac{\partial v}{\partial y} = e^x [x \cos y + 1 \cos y - y \sin y] - 0$$

$$f'(z) = e^x [x \cos y + \cos y - y \sin y] + i e^x [x \sin y + y \cos y + \sin y]$$

step (iv) By MTM, put $x=z$ & $y=0$

$$f'(z) = e^z [z(1) + (1) - 0] + i e^z [z(0) + 0 + 0]$$

$$f'(z) = e^z (z+1) + 0$$

step (v) Integrate w.r.t z

$$f(z) = \int (z+1) e^z dz + \kappa$$

$$= \{(z+1)[e^z] - (1)[e^z]\} + \kappa$$

$$f(z) = e^z [z+1-1] + \kappa = z e^z + \kappa$$

Dr. Uday Kashid (PhD, Mathematics)

Ex.3 Find orthogonal trajectories of family of curves $e^x [\cos(y)] - xy = c$ [EXTC-Nov-18, (6M)]

→ Assume $u = e^x \cos y - xy - c = 0$ → given

Step (i) $f(z) = u + iv$ → Analytic

(ii) $f'(z) = \frac{\partial u}{\partial x}$

(iii) By C-R eqn $\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$; $f'(z) = \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y}$

$$\frac{\partial u}{\partial x} = e^x \cos y - y - 0$$

$$\frac{\partial u}{\partial y} = -e^x \sin y - x - 0$$

$$f'(z) = (e^x \cos y - y) - i(-e^x \sin y - x)$$

(iv) By MTM, put $x=z$ & $y=0$

$$f'(z) = [e^z(1) - 0] + i(e^z(0) + z)$$

$$= e^z + iz$$

(v) Integrate w.r.t z

$$\int f'(z) dz = \int e^z dz + \int iz dz + \kappa$$

$$f(z) = e^z + i \frac{z^2}{2} + \kappa$$

$$f(z) = e^{x+iy} + \frac{i}{2}(x+iy)^2 + \kappa$$

$$= e^x e^{iy} + \frac{i}{2}(x^2 + 2ixy - y^2) + \kappa$$

$$\begin{aligned}
&= e^x e^{iy} + \frac{i}{2}(\underline{x^2 + 2ixy - y^2}) + \kappa \\
&= e^x [\cos y + i \sin y] + \frac{i}{2}(x^2 - y^2) - xy + \kappa \\
&= e^x \cos y + i e^x \sin y + \frac{i}{2}(x^2 - y^2) - xy + \kappa \\
&= [e^x \cos y - xy - c] + i \left[\frac{(x^2 - y^2)}{2} + e^x \sin y \right] \quad \text{let } \kappa = -c \\
&= u + iv \\
u &= e^x \cos y - xy - c \\
v &= e^x \sin y + \frac{(x^2 - y^2)}{2} \Rightarrow \begin{cases} \text{orthogonal trajectories of } u \\ \text{harmonic conjugate of } u \end{cases}
\end{aligned}$$

CASE-3

IF $(u-v) = \text{given}$

Dr. Uday Kashid (PhD, Mathematics)

step (i) We have $f(z) = u + iv \rightarrow \text{Analytic Fun} \rightarrow \text{---(i)}$

**step (ii) Multiple 'i' to Both side of step (i)

$$if(z) = iu + i^2v = iu - v \rightarrow \text{---(ii)}$$

***step (iii) Add step (i) & step (ii)

$$f(z) + if(z) = (u + iv) + (iu - v)$$

$$\underline{(1+i)f(z)} = \underline{(u-v)} + i \underline{(u+v)}$$

$$\text{step (iv)} \quad F(z) = U + iV \rightarrow \text{---(case-1)}$$

We have given $(u-v)$ i.e. $U = (u-v) = \text{given}$

$$\text{step (v)} \quad F'(z) = \frac{\partial U}{\partial x} + i \frac{\partial V}{\partial x}$$

$$\text{step (vi)} \quad \text{By C-R eqn} \quad \frac{\partial U}{\partial y} = -\frac{\partial V}{\partial x}$$

$$F'(z) = \frac{\partial U}{\partial x} - i \frac{\partial U}{\partial y}$$

step (vii) By MTM, put $x = z$ & $y = 0$

$$F'(z) = \left(\frac{\partial U}{\partial x} \right)_{x=z} - i \left(\frac{\partial U}{\partial y} \right)_{x=z}$$

$$F'(z) = \left(\frac{\partial U}{\partial x} \right)_{\substack{x=z \\ y=0}} - i \left(\frac{\partial U}{\partial y} \right)_{\substack{x=z \\ y=0}}$$

step-(vii)

Integrate w.r.t. z

$$F(z) = \int \left(\frac{\partial U}{\partial x} \right)_{\substack{x=z \\ y=0}} dz - i \int \left(\frac{\partial U}{\partial y} \right)_{\substack{x=z \\ y=0}} dz + C_1$$

$$(1+i)f(z) = \phi_1(z) - i\phi_2(z) + C_1$$

divide by $(1+i)$

$$f(z) = \frac{1}{(1+i)} \phi_1(z) - \frac{i}{(1+i)} \phi_2(z) + \frac{C_1}{(1+i)}$$

Dr. Uday Kashid (PhD, Mathematics)

CASE-(4) IF $(u+v)$ = given

step-① $f(z) = u + iv \rightarrow$ Analytic —①

①

$$if(z) = iu - v$$

②

$$\text{Add ① \& ② } (1+i)f(z) = (u-v) + i(u+v)$$

\rightarrow case-②

step-④

$$F(z) = U + iV$$

We have given $V = (u+v) = \text{given}$

step-⑤

$$F'(z) = \left(\frac{\partial U}{\partial x} \right) + i \frac{\partial V}{\partial x}$$

step-⑥

$$\text{By C-R eqn } \frac{\partial U}{\partial x} = \frac{\partial V}{\partial y}$$

$$F'(z) = \frac{\partial V}{\partial y} + i \frac{\partial V}{\partial x}$$

step-⑦

By MTM, put $x=z$ and $y=0$

$$F'(z) = \left(\frac{\partial V}{\partial y} \right)_{\substack{x=z \\ y=0}} + i \left(\frac{\partial V}{\partial x} \right)_{\substack{x=z \\ y=0}}$$

step-(viii)

Integrate w.r.t z

$$F(z) = \int \left(\frac{\partial V}{\partial y} \right)_{\substack{x=z \\ y=0}} dz + i \int \left(\frac{\partial V}{\partial x} \right)_{\substack{x=z \\ y=0}} dz + C,$$

$$F(z) = \int \left(\frac{\partial u}{\partial y} \right) \frac{1}{z^2} dz + \int \left(\frac{\partial v}{\partial x} \right) \frac{1}{z^2} dz$$

$$(1+i)f(z) = \phi_1(z) + i\phi_2(z) + C_1$$

$$\left[f(z) = u+iv = \frac{\phi_1(z)}{(1+i)} + \frac{i}{(1+i)}\phi_2(z) + \frac{C_1}{(1+i)} \right]$$

Ex.4 Find an analytic function $f(z)$ such that $u - v = (x - y)(x^2 + y^2 + 4xy)$

→ step-① we have $f(z) = u+iv \rightarrow$ Analytic fun

step ② $if(z) = iu - v$

step ③ Add, $(1+i)f(z) = (u-v) + i(u+v)$

Dr. Uday Kashid (PhD, Mathematics)

step ④ $F(z) = U + iV$

we have given $U = u-v = (x-y)(x^2+y^2+4xy)$ — given

step ⑤ $F'(z) = \frac{\partial U}{\partial x} + i \frac{\partial V}{\partial x}$

step ⑥ By C.R. eqns, $\frac{\partial U}{\partial y} = -\frac{\partial V}{\partial x}$

$$F'(z) = \frac{\partial U}{\partial x} - i \frac{\partial U}{\partial y} \longrightarrow$$

$$U = (u-v) = (x-y)(x^2+y^2+4xy)$$

$$\frac{\partial U}{\partial x} = (1-0)(x^2+y^2+4xy) + (x-y)(2x+0+4y)$$

$$\frac{\partial U}{\partial y} = (0-1)(x^2+y^2+4xy) + (x-y)(0+2y+4x)$$

$$F'(z) = \left[1(x^2+y^2+4xy) + (x-y)(2x+4y) \right] - i \left[-(x^2+y^2+4xy) + (x-y)(2y+4x) \right]$$

step ⑦ By M.T.M. put $x=z$ & $y=0$

$$F'(z) = \left[(z^2+0+0) + (z-0)(2z+0) \right] - i \left[-(z^2+0+0) + (z-0)(0+4z) \right]$$

$$= (z^2+2z^2) - i(-z^2+4z^2)$$

$$= 3z^2 - i(3z^2)$$

$$F'(z) = (1-i)3z^2$$

step ⑧ Integrate w.r.t z

$$\int F'(z) dz = (1-i)3 \int z^2 dz + C_1$$

$$f(z) = (1-i)z^3 + C_1$$

$$F(z) = (1-i) \Im\left(\frac{z^3}{3}\right) + C_1$$

$$** (1+i)f(z) = (1-i)z^2 + C_1$$

$$f(z) = \frac{(1-i)}{(1+i)} z^2 + \frac{C_1}{(1+i)} = \frac{(1-i)}{(1+i)} \times \frac{(1-i)}{(1-i)} z^2 + C = \frac{(1-i)^2}{1^2 - i^2} z^2 + C = \frac{(1-2i-1)}{2} z^2 + C$$

$$\boxed{f(z) = -i z^2 + C}$$

Ex.5. Find an analytic function $f(z)$ such that $u + v = \cos x \cosh y - \sin x \sinh y$

$$\rightarrow f(z) = u + iv \rightarrow \text{Analytic Fun}$$

$$if(z) = iu - v$$

$$(1+i)f(z) = (u-v) + i(u+v)$$

$$F(z) = u + iv$$

$$\text{But we have } v = (u+v) = \cos x \cosh y - \sin x \sinh y$$

$$F'(z) = \left(\frac{\partial u}{\partial x}\right) + i \frac{\partial v}{\partial x}$$

$$\text{By C.R eqn } \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$$

$$F'(z) = \frac{\partial v}{\partial y} + i \frac{\partial v}{\partial x} \quad \text{--- (1)}$$

$$\frac{\partial v}{\partial x} = -\sin x \cosh y - \cos x \sinh y$$

$$\frac{\partial v}{\partial y} = \cos x \sinh y - \sin x \cosh y$$

$$\text{By MTM } \left. \begin{aligned} \left(\frac{\partial v}{\partial x}\right)_{x=z, y=0} &= -\sin z (1) - \cos z (0) = -\sin z \\ \left(\frac{\partial v}{\partial y}\right)_{x=z, y=0} &= \cos z (0) - \sin z (1) = -\sin z \end{aligned} \right\}$$

$$F'(z) = \left(\frac{\partial v}{\partial y}\right)_{x=z, y=0} + i \left(\frac{\partial v}{\partial x}\right)_{x=z, y=0}$$

$$= -\sin z + i(-\sin z)$$

$$f'(z) = -\sin z (1+i)$$

Integrate w.r.t z

$$f(z) = (1+i) \cos z + C_1$$

$$(1+i)f(z) = (1+i) \cos z + C_1$$

$$f(z) = \cos(z) + \frac{C_1}{(1+i)} = \cos z + C$$

Dr. Uday Kashid (PhD, Mathematics)