

Cite this: DOI: 00.0000/xxxxxxxxxx

Received Date
Accepted Date

DOI: 00.0000/xxxxxxxxxx

What a radical-pair quantum reservoir would require: readable capacity, no quantum advantage, and a nuclear-register criterion[†]

Hikaru Wakaura^{*a} and Taiki Tanimae^a

Whether the electron–nuclear spin coherence behind radical-pair magnetoreception could also support biological *computation* has been conjectured for decades but never made quantitative. We treat a cryptochrome-type radical pair as a reservoir computer, solving the radical-pair master equation with electronic decoherence and driving it as a real turnover does—by the birth of a spin-correlated pair whose singlet character encodes the input. Three results follow, none of them the expected one. The reservoir is *readable by ordinary chemistry*: the nuclear polarisation the product carries away gives an out-of-sample information-processing capacity of 4.7 against its own memoryless floor of 2.0, with no spectroscopy. There is no quantum advantage: a classical network matched channel-for-channel exceeds the five-spin reservoir by 49–123%. And the memory horizon is set not by the microsecond pair lifetime but by the turnover interval, because the nuclear register survives recombination in the diamagnetic product—so the binding constraint is chemical rather than temporal, namely that the *same* register be reused between turnovers. Predicting the register’s relaxation instead of assuming it makes that constraint testable: at the geomagnetic field a proton on the intact protein relaxes in milliseconds, not seconds, capping the horizon below the neural band unless the register’s dipolar vector reorients about two and a half times faster than the protein ($\tau_{\text{eff}} \lesssim 6$ ns). What survives is a smaller claim on firmer ground: a criterion whose two constants can be measured in vitro, in place of one whose central relaxation time was never measured at all.

1 Introduction

The radical-pair mechanism (RPM) is the only experimentally established ~~route-by-way in~~ which a magnetic field as weak as the geomagnetic field (~ 50 μT) can ~~alter-change~~ the outcome of a chemical reaction.^{1,2,16} A spin-correlated pair born

in a singlet electronic state undergoes coherent singlet–triplet (S–T) interconversion driven by hyperfine ~~coupling-to-magnetic nuclei~~ fields; because recombination is spin-selective, a field that re-tunes ~~the interconversion-changes this interconversion alters~~ the product yields. ~~In cryptochrome the mechanism, hosted by a flavin-tryptophan radical pair, The mechanism~~ underlies the leading model of avian magnetoreception.^{2–5,12} ~~where the photoreceptor cryptochrome hosts a flavin-tryptophan radical pair~~^{2–5,12}

The success of the RPM has ~~repeatedly~~ motivated a bolder conjecture: that spin coherence in biomolecules might support not merely sensing but ~~information-processing—a information processing—a~~ “quantum brain”.^{6,7} ~~The idea has been—~~^{6,7} a ~~proposal long~~ contested on decoherence grounds² ~~and sits within—~~⁹ and part of the broader programme of quantum biology.^{10,11} ~~A concrete, named instance is the three-layer quantum-brain hypothesis~~^{19,20}, which posits a layered architecture—a long-lived nuclear-spin memory (Layer 1), a radical-pair read/write interface (Layer 2), and classical neuronal transduction (Layer 3)—and which, in its most recent form,

^aQIRI (Quantum Integrated Research Institute Inc.), Tokyo 107-0061, Japan. E-mail: h.wakaura@deeptell.jp QIRI (Quantum Integrated Research Institute Inc.), 1–16–3 Akasaka, Minato-ku, Tokyo 107-0061, Japan. E-mail: h.wakaura@qiri.co.jp; t.tanimae@qiri.co.jp. ORCID 0000-0001-8381-8323.

[†] Electronic Supplementary Information (ESI) available: ~~the~~ Liouville-space solver ~~validation-against-and its agreement with the coherent limit; the capacity (HPC/MC) definitions—estimator, its controls and the Dambre bound; the injection map and the product-register variant; the chemically accessible readout—throughput routes; the coherence fraction at matched hopping rate and finite-size scans~~ semiclassical reference; the ~~ensemble-heterogeneity-model~~ classical baselines; the ~~quantum-versus-classical-dephasing-sweep~~ turnover clock and the register-reuse criterion; which nuclei can hold the register; ensemble averaging and ~~cryptochrome turnover desynchronisation; anisotropic hyperfine coupling and the nuclear spin quantum number; predicting the two unmeasured constants; and the spin parameters with sources.~~ All code (the open-source qbscreen package) and ~~every~~ numerical ~~data-output~~ are available. See DOI: 00.0000/00000000.

already reframes the proposal from quantum *computation* to quantum *reservoir dynamics*. The present paper is the quantitative test of that architecture as a reservoir, and the bound it returns. Since that reframing is not new here, we state the delta explicitly: Ref. ²⁰ proposes the reservoir reading and estimates coherence lifetimes, but drives the reservoir by resetting a single electron, reads it through spin observables that presuppose magnetic-resonance access, and identifies the memory horizon with the radical-pair lifetime. What is new here is (i) a chemically consistent drive in which the electron pair is reborn correlated each turnover, (ii) readouts restricted to quantities a downstream reaction could use, each scored against its own memoryless floor, (iii) classical baselines matched by readout dimension, and (iv) the replacement of the lifetime clock by the turnover clock together with the register criterion that follows from it. The spin Hamiltonian, the Haberkorn treatment and the capacity estimators are standard and are not claimed as contributions. A companion study from this programme shows, using a decoherence-explicit master equation, program places the conjecture on quantitative footing from the sensing side. Solving the radical-pair master equation with electronic decoherence explicitly included, it finds that the active-site radical pairs of the principal neurotransmitter-metabolising flavoenzymes (monoamine oxidase A/B, D-amino-acid oxidase) are magnetically inert ($<10^{-4}\%$ field effect): their gigahertz exchange coupling and picosecond lifetimes forbid weak-field mixing. ²⁴ Only a spatially-separated: the computed geomagnetic effect on the singlet yield is $1.7 \times 10^{-12}\%$, and even the most favourable case that analysis can construct—the coherent limit with an unphysically long lifetime—caps it at $4.0 \times 10^{-6}\%$. Cryptochrome in the identical framework gives $+1.0\%$. What decides this is the picosecond active-site lifetime rather than the exchange coupling: at $\tau \approx 10$ ps the response is unchanged to within a few per cent whether or not the gigahertz J is present. The discriminating variable is the inter-radical separation the protein scaffold imposes, not the flavin the two share. Only a spatially separated radical pair, as in cryptochrome (cryptochrome—with $J \approx 0$, lifetime $\tau \sim \mu\text{s}$), retains the long-lived electron–nuclear and a microsecond lifetime—sustains the coherence required for weak-field sensitivity²³—a^{23,24}. This premise is a standard result of radical-pair compass theory^{2,23} and is logically independent of the companion analysis; we take the existence of a long-coherence separated radical pair as our self-contained starting point.

Here we ask whether that same coherence can do computational work. We pose the question—This sharp dichotomy invites the question we address here. The separated radical pair is singled out because it preserves long-lived electron–nuclear spin coherence. Is that coherence good for anything besides sensing the geomagnetic field? We argue the question is best posed not as gate-model quantum computation—which fights decoherence and demands error correction, impossible for a warm, 10^{10} -copy biomolecular ensemble—but computation but as reservoir computing, which requires dissipation to supply its fading-memory (echo-state) property.^{25,26} Quantum reservoir computing (QRC) is established in nuclear- and electron-spin ensembles;^{13,27–30}

the physical correspondence we adopt maps long- T_2 nuclear spins to the memory, the radical-pair electrons to a nonlinear read/write head, hyperfine coupling to the read/write channel, and electronic decoherence to the fading memory (Fig. 1). The conceptual thesis is that the same decoherence that kills the magnetosensor's coherent signal supplies the reservoir's memory—decoherence is the enemy of the sensor and the friend of the reservoir. This is the named three-layer quantum-brain architecture^{19,20} made quantitative as a reservoir.

Our contribution is a physical-chemistry analysis: an exact Liouville-space Haberkorn master equation^{17,31} with electronic dephasing explicit, benchmarked against classical reservoirs and a standard nonlinear task, and applied to the real cryptochrome radical-pair with literature spin parameters and we answer it with the chemistry left in: driven, read and clocked as a turnover actually is. We state the central biological caveat at the outset the load-bearing caveat: whether a neuronal cryptochrome forms a light- or reaction-driven separated radical pair at all is unestablished—mammalian unestablished—the mammalian CRY1/CRY2 proteins are primarily circadian transcriptional regulators¹⁵—so every biological conclusion is conditional, while the below is conditional on the existence of such a pair. The spin-dynamics results (capacity, classical baselines, the register criterion) do not depend on that condition and stand on their own.

The central idea. (a) A separated radical pair ($J \approx 0$, $\tau \sim \mu\text{s}$) serves two functions: long electronic coherence makes it a weak-field magnetosensor, while the dissipation that destroys that coherent signal supplies the fading memory a reservoir needs. (b) The three-layer reservoir: nuclear spins are the long- T_2 memory, the radical-pair electrons are the nonlinear read/write head driven by input s_k through hyperfine coupling, and a chemical product (yield or CIDNP polarisation) is the classical readout. Its memory horizon $MC \times \tau \approx 3 \mu\text{s}$ strands biological use.

Why a reservoir, not a gate. Gate-model quantum computation treats decoherence as the adversary and demands error correction; a warm, wet, 10^{10} -copy biomolecular ensemble cannot meet that bar. Reservoir computing (RC) inverts the relationship. An RC system feeds a time-dependent input into a fixed nonlinear dynamical “reservoir” and trains only a linear readout of its state^{25,26}. RC requires dissipation: the echo-state / fading-memory property—that the influence of past inputs decays—is supplied precisely by relaxation^{27,28}. The molecular-spin and NMR communities have demonstrated quantum RC explicitly in nuclear- and electron-spin ensembles^{27–29}, part of a broad physical-reservoir-computing effort¹³. In this system the fading memory is supplied not by electronic dephasing—varying T_2^e over the physically plausible range 50–2000 ns at the engineered operating point changes the capacity by under 2% (the two operating points are defined in Methods and must not be conflated)—but by the turnover itself: each cycle discards the electron pair and keeps only the nuclei, which is a dissipative, non-unitary map. It is the openness of the chemical cycle, not the rate of electronic decoherence,

that makes the system a reservoir. This is consistent with the large coherence fraction reported below, which is measured in a very different regime: dephasing rates 10^3 times larger than any realistic T_2^e , applied together with a compensating extension of the pair lifetime. Realistic electronic decoherence is not the limiting factor; complete destruction of electronic coherence still is.

2 Theory and methods

1.1 Reservoir spin Hamiltonian

We model the reservoir as $N = 5$ spins—two Our contribution is to answer that question with the chemistry left in, and the answer overturns three natural expectations. (i) We drive the reservoir as a real turnover does—nuclei retained, a new spin-correlated pair created each cycle—rather than by resetting a single electron, and read it through the products a downstream reaction could actually use. (ii) We compare against classical reservoirs matched by readout dimension, and find no quantum advantage, which we report rather than argue around. (iii) We show that the timescale objection usually raised against this system rests on identifying the reservoir's clock with the radical-pair electrons (e_1, e_2) and three nuclei (^{31}P and two ^1H)—with Hamiltonian (in MHz)—

$$H = \omega_e(S_z^{e1} + S_z^{e2}) + \sum_{\mu=x,y,z} [A_P S_\mu^{e1} I_\mu^P + A_{H1} S_\mu^{e1} I_\mu^{H1} + A_{e2} S_\mu^{e2} I_\mu^{H2} + J S_\mu^{e1} S_\mu^{e2}],$$

where $\omega_e = g_e \mu_B B / \hbar$. The input electron e_1 writes the input into the nuclear memory through the hyperfine terms; J is a weak inter-electron exchange. At the engineered operating point we use $A_P = 80$, $A_{H1} = 40$, $A_{e2} = 15$ MHz, $J = 2$ MHz, $B = 1$ mT, $\tau = T_2^e = 50$ ns—representative molecular-spin-RC values, not a fit. The geomagnetic, $J = 0$ cryptochrome regime is treated separately below.

1.1 Master equation and propagator

Each reservoir update is the completely positive map $\rho \mapsto e^{\mathcal{L}\tau} \rho$ of the Haberkorn master equation³¹ with electronic pure dephasing,

$$\dot{\rho} = -i[H, \rho] - \frac{1}{2} \{k_S P_S + k_T P_T, \rho\} + \sum_i \frac{1}{T_2^e} (S_z^i \rho S_z^i - \frac{1}{4} \rho),$$

solved exactly in Liouville space. The third term—the electronic decoherence shown decisive in the companion sensing analysis—is here the source of fading memory. The solver reproduces the closed-form coherent Green's function singlet yield to machine precision in the limit $1/T_2^e \rightarrow 0$, $k_S = k_T$ (ESF) lifetime, and replace it with a chemical criterion—reuse of the nuclear register—that is falsifiable by chemistry. That criterion still compares two times, as the argument it replaces did; what the spin-dynamical treatment supplies, and no timescale comparison would, is which nuclear relaxation constant appears in it. Within a step no recombination acts; the readout step uses the spin-selective term.

1.1 Reservoir protocol

We use the standard temporal-QRC protocol.²⁷ At step k the input electron is reset to $|\psi(s_k)\rangle = \sqrt{1-s_k}|0\rangle + \sqrt{s_k}|1\rangle$ while the reservoir is retained; the state evolves for τ under $e^{\mathcal{L}\tau}$,

~~observables are read~~ The physical correspondence we adopt, adapting the spin-RC framework of Fujii and Nakajima²⁷, maps the reservoir-computing elements onto the radical pair as follows: the memory nodes are the nuclear spins (in the cryptochrome case ^{14}N and ^1H), which store the input in their *populations*, so that the relevant relaxation constant is T_{1n} —though at the geomagnetic field extreme narrowing makes T_{1n} and a linear (ridge) readout is trained. The “reset” has a natural chemical reading: each catalytic or photochemical cycle T_{2n} equal, and both short (ESI, Sec. S12); the nonlinear read/write head is the radical-pair electron system; the read/write channel is the hyperfine coupling; and the fading memory (echo-state property) is supplied by the openness of the chemical cycle together with inter-cycle nuclear *creates a fresh radical-pair* depolarisation encoding the instantaneous input, while the surrounding nuclei—not consumed by recombination—carry spin polarisation across cycles. The reservoir is thus a persistent nuclear bath repeatedly interrogated by transient electron pairs, not an artificial gate sequence—not nuclear dephasing, which we show below costs the reservoir almost nothing. These are exactly the three layers of the hypothesis above: Layer 1 (nuclear-spin memory), Layer 2 (the radical-pair interface), and Layer 3 (the classical readout/transduction that trains the linear map).

The paper is organized as follows. Section 2 establishes the reservoir's capacity and its convergence, localises that capacity in spin coherence, compares it against matched classical baselines, and then asks what a chemical readout recovers; it closes with the reservoir's clock and the nuclear-register criterion. Section 3 places the result among quantum-brain proposals and states the limitations. The model, protocol, and capacity definitions are collected in Appendix A.

1.1 Capacity measures

We quantify the reservoir with the linear memory capacity $\text{MC} = \sum_d r^2(s_{k-d})$ ³² and

2 Results

We drive the reservoir the information processing capacity $\text{IPC} = \text{MC} + \sum_{\text{nonlin}} C$ (degree-2 Legendre targets).³³ A theorem bounds $\text{IPC} \leq n_{\text{obs}}$, way a real turnover does. Each cycle the nuclei are retained and a new spin-correlated electron pair is created whose singlet character encodes the input, $\rho_e(s) = s|S\rangle\langle S| + (1-s)P_T/3$; the pair then evolves under the Liouville-space master equation with electronic dephasing and, where the readout requires it, spin-selective recombination. This replaces the reset of a single electron used in earlier treatments, which is not a chemical process and which—for a separated pair—destroys the number of independent readout observables. *All capacities are out of sample*: each reconstruction is electron-electron correlation every cycle. All capacities are out-of-sample (trained on the first half of a run and, scored on the held-out second half, so values do not reward overfitting; we report means over 5–10 input realisations. Code and data are open (qbscreen; ESI)). The number of input realisations is not uniform and is stated with each result: 12 for the turnover-clock




Fig. 1 The reservoir and the quantity that decides its reach. (a) A cryptochrome-type separated radical pair ($J \approx 0$, lifetime $\tau \sim \mu\text{s}$) is the same molecule in two roles: long electronic coherence makes it a weak-field magnetosensor, while the dissipation of an open cycle supplies the fading memory a reservoir needs. (b) The nuclei are the register: they are not consumed by recombination and therefore persist from one turnover to the next, whereas the electron pair is created afresh each cycle. The memory horizon is consequently set by T_{turnover} and not by τ , and is quoted throughout as $(MC - C_0) \times T_{\text{turnover}}$, the zero-delay term $C_0 \simeq 1$ carrying no memory. It reaches the neural band only if the register is genuinely reused and survives the pause, which requires it to be carried by protons rather than the quadrupolar flavin ^{14}N and to reorient with $\tau_c \lesssim 6$ ns (ESI, Secs. S9 and S12).

scan and its paired test, which carry the central claim; 8 for the classical baselines and the T_2^e scan at the engineered point; 6 for the headline capacity, the chemical-readout survey, the route floors, the delay kernel, the grid scans, the nuclide scan and the cryptochrome-point classical baselines; 4 for the register-reuse and nuclear-channel scans and the semiclassical reference; and 3 for the ensemble and anisotropy scans. Four quantities are single-realisation: the ridge and shot-noise sensitivity scans and the semiclassical integration-step check, which compare variants on one fixed input by design, and the superseded clock scan retained as ESI Table S7, whose single realisation is why it was replaced. Model, protocol and estimator are given in Appendix A.

3 Results and discussion

2.1 Capacity of the reservoir

2.2 The radical pair is a quantum reservoir

With a 12-observable readout (single-spin Pauli components on the nuclei, electron S_z , and the e_1-e_2 singlet projector), the reservoir attains $MC = 3.6$ and $IPC = 6.9 \pm 0.7$ (Fig. 2), of which 3.2 With a twelve-observable readout the five-spin reservoir attains

$$MC = 3.9, \quad IPC = 5.6 \pm 0.2,$$

of which 1.7 units are nonlinear, against the bound $n_{\text{obs}} = 12$. Unlike the single-electron-reset protocol, the estimate is converged in the length of the driving sequence: repeating at $L = 600, 1200, 2400, 4800$ gives $IPC = 5.28, 5.60, 5.87, 5.84$, i.e. flat to within the seed-to-seed scatter beyond $L \approx 2400$. The memory kernel decays over ~ 4 steps (fading memory); total IPC increases with readout dimension, confirming the readout—not the spin dynamics—sets the ceiling, and is robust to strong inter-eye nuclear dephasing and to a shorter, more realistic T_2^e (6.0–7.2 cycles, the signature of the fading-memory property).

2.2 An operational coherence fraction

To ask how much of this is quantum we add a pure-dephasing rate γ in the S_z product basis and take the incoherent limit at matched hopping rate: since the golden-rule transfer rate scales as $|H_{ij}|^2/\gamma$, holding rate $\times \tau$ fixed requires $\tau \propto \gamma$. Doing so, the capacity saturates at $IPC = 1.00$ for $T_2^e = 5\text{--}50$ ns; $\gamma = 10^3, 5 \times 10^3, 2 \times 10^4$ MHz — exactly the memoryless floor, i.e. instantaneous read-back of the input with no memory at all. Against the coherent value 5.42 this is a coherence fraction

$$\frac{IPC_{\gamma=0} - IPC_{\text{incoh}}}{IPC_{\gamma=0}} = 0.82. \quad (1)$$

Taking the limit at fixed τ instead, as is sometimes done, measures Zeno freezing rather than the classical limit and inflates this number; we avoid that.

Capacity of the radical-pair reservoir. — (a) Linear memory $r^2(s_{k-d})$ versus delay for the multi-observable (blue) and scalar (red) readouts. — (b) Total IPC versus number of readout observables (ten-seed mean), saturating below the Dambre bound. — (c) Capacity is robust to inter-cycle nuclear dephasing p . Two caveats are load-bearing. First, γ as applied above acts on

all five spins, so one might suspect the collapse merely reflects erasure of the nuclear register rather than loss of electronic coherence. It does not: restricting the added dephasing to the two electrons and leaving the nuclei untouched gives $IPC = 0.9949, 0.9891$ and 0.9891 at the same three rates, i.e. a coherence fraction of 0.817 against 0.816 for the all-spin channel (ESI, Sec. S5). The number is a statement about the electrons. Second, and more limiting, this is an *operational* fraction defined by one particular matched-hopping construction in which τ is stretched by up to a factor of 400; it is not a decomposition against a physically motivated classical model. The appropriate such reference is a semiclassical radical-pair treatment with classical hyperfine vectors and rate equations¹⁶. We have since run it (ESI, Sec. S8): a fully classical model of the same spin system scores at its own memoryless floor, giving a coherence fraction of 0.859 , so the 0.82 reported here is if anything conservative. We nonetheless report 0.82 as what it is—the capacity lost under a specific incoherent limit—rather than as a measure of “how quantum” the reservoir is. Note also that this quantity, like the capacity above it, is evaluated at the engineered operating point.

2.3 Classical baselines and resource efficiency

2.3 There is no quantum advantage

A capacity is only meaningful against a classical reference.—We compare to—, and the comparison must be symmetric in what is read out. Driving tanh echo-state networks (ESNs) read-out by the with the same input and scoring them with the same IPC estimator (Fig. 3). Per physical unit the quantum reservoir win estimator: five spins give $IPC = 6.9$, five classical nodes 4.6 , because $N = 5$ spins span a $2^5 = 32$ -dimensional Hilbert space—the exponential-state-space argument of QRC²⁷ made quantitative against an explicit baseline.—On the NARMA2 benchmark, $NMSE = 0.21$ (quantum) versus 0.32 (five-node ESN). But there is no advantage per observable—a classical reservoir of equal readout dimension already outperforms it (10.2

Table 1 The quantum reservoir against matched classical reservoirs at the engineered operating point.

reservoir (readout features)	IPC
quantum, 5 spins (12 observables)	5.59 ± 0.19
ESN, 5 nodes (5 linear features)	4.55 ± 0.32
ESN, 5 nodes (5 linear + 7 quadratic)	9.41 ± 0.86
ESN, 12 nodes (12 linear features)	10.37 ± 0.59

Matched by *physical unit* the quantum reservoir is modestly ahead (5.59 vs 6.9 at 12 nodes)—nor in absolute terms (31 at 50 nodes). The value is resource efficiency, not computational supremacy, and we say so.— 4.55), but matched by *readout dimension*—the resource that actually sets the Dambre ceiling—a classical system of the same size is well ahead (9.41 vs 5.59). We therefore make no claim of quantum advantage. Which accounting is the fair one depends on a cost model for non-commuting measurements versus multiplications that we do not have; we report the numbers and leave the claim unmade.

That comparison is made at the engineered operating point, which is not the biologically relevant one. Repeating it at the cryptochrome point, against ESNs matched channel-for-channel to the chemical readouts of the next section, removes the ambiguity entirely:

Table 2 The same comparison at the cryptochrome operating point, matched channel-for-channel.

system (readout features)	IPC
quantum, time-resolved kinetics (5 ch)	2.01 ± 0.09
ESN, 5 nodes (5 features)	4.47 ± 0.33
quantum, kinetics + CIDNP (8 ch)	4.65 ± 0.15
ESN, 8 nodes (8 features)	6.94 ± 0.56
ESN, 12 nodes (12 features)	9.62 ± 0.93

At the cryptochrome point the classical reservoir wins decisively whenever the comparison is matched channel-for-channel: by 49% at eight channels and 123% at five. Matched by *physical unit* instead—five spins against five nodes—the quantum reservoir is nominally ahead, 4.65 ± 0.15 against 4.47 ± 0.33 , but that 4% margin is 0.5σ of the seed scatter and we do not read it as an advantage. The honest summary is that the two channel-matched accountings favour the classical system by a wide margin and the third distinguishes nothing. On no accounting does the radical pair compute *better* than a small classical network by a margin exceeding its own seed scatter. That is the honest state of the comparison, and it removes the hedge we left in the preceding paragraph.

2.4 Biology can read the reservoir

Biology has no EPR spectrometer; it has products. We therefore restrict the readout to observables a downstream reaction could plausibly use, with recombination switched on so that the quantity read is the yield actually produced, $Y_X = k_X \int_0^\tau \text{Tr}[P_X \rho(t)] dt$ (Fig. 2):

The memoryless floor is *not* common to these routes, and quoting a single floor of $IPC = 1$ overstates what the richer readouts achieve. We therefore measure each route’s own floor by wiping the nuclear register every cycle ($q = 1$), which leaves only what the route can compute from the present input, and report the excess above it. All seven routes defined by the model are listed; none is omitted.

Four things follow, and only the first is the one we originally reported. First, the readout is genuinely not the limiting resource: time-resolved product kinetics clear their own floor by a factor of two, and the nuclear polarisation the product carries away reaches 4.65 against a floor of 2.01 —an excess of 2.64 , not the factor of 4.6 that a universal floor of 1 would suggest. The CIDNP route is still the best readout by a wide margin, and it still requires no spectroscopic access.

Second, a single scalar end-point yield gives $IPC = 1.02$ against a floor of 1.02 : it reports the present input and remembers *nothing*. Claims about reservoir behaviour should not be based on it.

What biology can read: excess over each route's own floor




Fig. 2 Readout is not the bottleneck. Capacity recovered by chemically accessible observables at the cryptochrome operating point ($B = 50 \mu\text{T}$, $J = 0$, $\tau = 1 \mu\text{s}$), with spin-selective recombination switched on so that the observable is the product actually formed. All seven routes defined by the model are shown. Red bars mark each route's *own* memoryless floor, measured by wiping the nuclear register every cycle ($q = 1$); these are not common to the routes (1.02 for the yield routes, 2.01 for the eight-channel CIDNP route, 1.96 for the accumulated pool), so it is the dashed excess above each floor—printed above each bar—that measures memory, not the bar height. Ordinary reaction kinetics clear their floor by 0.99, and the nuclear polarisation carried away by the product—chemically induced dynamic nuclear polarisation, CIDNP¹⁸—clears its own by 2.64, the largest of any route. The end-point yield sits exactly on its floor: it reports the present input and remembers nothing. The accumulated pool is hatched because a leaky downstream integrator is itself a classical delay line, so its excess is not evidence of nuclear-spin memory. No spectroscopy is involved.

Table 3 Classical-baselines Capacity recovered by chemically accessible observables at the cryptochrome point, each scored against its own memoryless floor. (a) IPC per physical unit:—

readout route	ch.	IPC	floor	excess
end-point singlet yield	1	1.02	1.02	0.00
both exit channels, end point	2	1.37	1.02	0.36
three micro-environments (different τ)	3	2.56	1.02	1.54
time-resolved kinetics, one product pool	5	2.01	1.02	0.99
both exit channels, time-resolved	10	2.01	1.02	0.99
kinetics + CIDNP on the product	8	4.65	2.01	2.64
kinetics + accumulated downstream pool	10	2.71	1.96	0.75

Third, the readout saturates in channel count. Doubling the time-resolved kinetics from 5 to 10 channels by adding the second exit channel buys 0.003 in capacity. Whatever limits this reservoir, it is not the number of product observables a downstream reaction can distinguish.

Fourth—and this is a caution about the accumulated-pool route rather than a result—a downstream integrator carries memory of its own. Its floor is 1.96 with a linear memory capacity of 1.94 even when the nuclear register is wiped every single cycle, because the leaky accumulator is itself a classical delay line. Its apparent capacity is therefore not evidence of nuclear-spin memory, and we exclude it from the claims below.

The mechanism behind the CIDNP column is worth stating, because it also explains a negative result. The reduced state

of *either* electron of a newly born pair is maximally mixed for every input, since both $|S\rangle\langle S|$ and $P_T/3$ have zero one-electron polarisation; the input lives entirely in the two-electron correlation. With $J = 0$ the two radicals do not interact, so nothing can transfer that correlation to a nucleus—and indeed, with recombination switched off, the cryptochrome-point capacity is 1.01, the memoryless floor. It is *spin-selective recombination* that converts the singlet-triplet correlation into a population difference and so writes the input into the nuclear register. In a separated radical pair, recombination is not merely how the reservoir is read; it is how it is written.

2.5 The clock is the five-spin reservoir exceeds a same-size ESN; an ESN needs more nodes than turnover, not the molecule has spins to match its pair lifetime

The horizon over which a reservoir remembers is MC multiplied by its cycle time. It is natural, and it is what the submitted version of this work did, to identify that cycle time with the radical-pair lifetime $\tau \sim 1 \mu\text{s}$ and conclude that a $3 \mu\text{s}$ horizon puts the system 10^3 – 10^5 below neural timescales. That identification is wrong, and its own model says so. The carrier of the memory is the nuclear register, and nuclei are not consumed by recombination: they leave in the diamagnetic product, where nuclear relaxation times are of order 0.1–1 s—a high-field figure, which Sec. 2.6 and the ESI, Sec. S12, revise sharply downwards for the geomagnetic field. The cycle time of the reservoir is therefore the interval between turnovers.

Inserting a pause T_d between turnovers and relaxing the register with probability $q = 1 - \exp(-T_d/T_{1n})$ (with $T_{1n} = 1 \text{ s}$) gives Fig. (b) NARMA2 (lower better) 3(a):

Before the numbers, the horizon needs a definition that is actually a time. MC is a sum of squared correlations over delays $d \geq 0$, and the quantum reservoir beats a size-matched ESN but is surpassed— $d = 0$ term—which reports the present input and remembers nothing—contributes $C[0] \simeq 1$ for every readout. Multiplying the raw MC by the cycle time therefore assigns a positive “horizon” to a system with no memory at all, growing without bound as the cycle is lengthened. We instead quote $(MC - C[0]) \times T_{\text{cycle}}$, the delayed memory only. The measured kernel justifies this: for the CIDNP readout $C[0] = 1.00$, $C[1] = 0.98$, $C[2] = 0.47$, falling below 5% of $C[0]$ by a large classical one: $d = 5$, so $MC - C[0] = 1.87$ of genuinely delayed capacity.

2.6 The capacity is quantum

To localise the capacity we add a tunable all-spin-

Twelve input realisations; the quoted spread is the between-seed standard deviation. Because the same seeds are used at every T_d , the change is properly assessed on paired differences: across five decades of turnover interval the capacity falls by 0.046 ± 0.011 (SEM, $n = 12$, $t = 4.23$), that is by 1.6%. This is a real decrease, resolved well above both the seed noise and the time-grid error (0.17% from $n_t = 96$ to 192). It is also roughly eight times larger than the 0.2% quoted in the originally submitted version, which came from a single input realisation and from the five-channel rather than the eight-channel column. The qualitative conclusion is unchanged and now rests on an adequate sample: the capacity is nearly, but not exactly, independent of the turnover interval, so the horizon tracks the clock.

A turnover of 10–100 ms therefore places the memory horizon at 19–188 ms, inside the neural band, provided the register survives the pause. That proviso is not free, and Sec. S12 shows it is what now binds: at the geomagnetic field a proton on the intact protein relaxes in 2.4 ms, which caps the horizon at 6.1 ms and puts it below the band unless the register reorients about two and a half times faster than the protein. The numbers in this table should therefore be read as what the spin dynamics permit, not

as what the chemistry delivers. We note also that a 1 ms turnover gives 1.9 ms, below the 10 ms–1 s band drawn in Fig. 3(a); the originally quoted range of “3–280 ms, inside the neural band” overstated this at its lower end. The timescale objection based on the microsecond pair lifetime does not survive in any of these cases: the same spin physics runs on a slow clock.

2.6 The criterion: reuse of the nuclear register

What replaces it is a chemical requirement, and a sharper one. Everything above assumes that the register the reservoir writes into is the same register on the next turnover. Relaxing that assumption by depolarising the nuclei with probability q between cycles gives Fig. 3(b): the eight-channel memory capacity falls from 2.89 at $q = 0$ to 2.67 at $q = 0.3$ and 2.29 at $q = 0.5$ —a loss of a fifth by the time the turnover interval reaches the nuclear relaxation time, not the near-invariance we previously described—then through 2.03 at $q = 0.7$ and 1.98 at $q = 0.9$, and at $q = 1$, a fresh molecule every cycle, collapses to $MC = 1.00$, the memoryless floor.

Which nuclear relaxation process enforces this can be asked directly. The register we propagate is a full nuclear density matrix, so one might expect memory to require the survival of nuclear coherences as well as populations, making the binding constant $\min(T_{1n}, T_{2n})$. It does not. Replacing the depolarising channel by a pure-dephasing rate γ in the S_z basis; as $\gamma \rightarrow \infty$ the coherent flip-flop becomes incoherent hopping (a classical Markov chain). The held-out IPC falls from 1.36 channel on each nucleus, which destroys nuclear coherence while preserving $\langle I_z \rangle$ exactly, leaves the capacity untouched: $MC = 2.89$ at $\gamma = 0$ to 0.39, a coherence fraction $(1.36 - 0.39)/1.36 = 0.72$ (ESI). This destructive intervention shows spin coherence performs most of the computational work—the singlet-triplet precession has no classical analogue—though it is not a claim of quantum advantage over an optimised classical model. It also controls for the input encoding: held fixed as γ grows, yet the capacity collapses: $q = 0$ and 2.87 at $q = 1$, against 1.00 for the depolarising channel at $q = 1$.

An optimal coherence time: a sensing-computing trade-off a useful reservoir sits at intermediate dissipation. Scanning T_2^c at fixed $\tau = 50 \text{ ns}$ (Fig

The memory is carried entirely by nuclear z -populations—which is exactly what the CIDNP readout reports—and is indifferent to nuclear phase. The relevant constant is therefore T_{1n} and not T_{2n} . -4)

We must immediately withdraw the rescue this appears to offer. At the geomagnetic field the register is in extreme narrowing ($\omega\tau_c \approx 2 \times 10^{-4}$), and there $J(0)$, $J(\omega)$ and $J(2\omega)$ coincide, so T_{1n} and T_{2n} agree to one part in 10^7 (ESI, Sec. S12). The distinction between a population register and a coherence register is therefore real as a statement about what the reservoir uses, but it buys nothing at the operating point: both constants take the same, short, value there. It would matter only at fields where $\omega\tau_c \gg 1$, that is, in an NMR magnet rather than in an organism.

It does not, however, make every nucleus eligible. A T_{1n} of order a second is measured for protons in proteins³⁵




Fig. 3 (a) The memory horizon against the turnover interval, from twelve input realisations (error bars are the between-seed standard deviation). Because the nuclear register survives recombination, inserting a physiological pause costs only 1.6% of the memory capacity across five decades, so the horizon simply tracks the clock and enters the neural band (shaded, 10 ms–1 s, the range of cortical rhythms⁸). Both definitions are shown: the raw $MC \times T_{\text{turnover}}$ (grey) counts the zero-delay term, which carries no memory, while the delayed-memory horizon $(MC - C_0) \times T_{\text{turnover}}$ (green) is the quantity quoted in the text—on the latter the 1 ms turnover falls below the band. The dotted line is where the horizon would sit if the clock were the radical-pair lifetime, for the same eight-channel readout. (b) The criterion, and which relaxation process enforces it. Depolarising the register between turnovers with probability q (green) drives the capacity to the memoryless floor: at $q = 1$ —a fresh molecule each cycle— $MC = 1.00$ and there is no memory. Pure nuclear *dephasing* at the same q (orange) leaves $MC = 2.87$, essentially untouched: the register stores populations, not phase, so the constant that bounds the horizon is T_{1n} . Note that this does not lengthen the horizon at the operating point, where extreme narrowing brings T_{1n} and T_{2n} together (ESI, Sec. S12).

Table 4 The memory horizon against the turnover interval, twelve input realisations.

T_d	q	MC (8 ch)	$MC \times T$	$(MC - C_0) \times T$
1 μ s	10^{-6}	2.926 ± 0.149	5.9 μ s	3.9 μ s
1 ms	10^{-3}	2.926 ± 0.148	2.9 ms	1.9 ms
10 ms	10^{-2}	2.922 ± 0.145	29 ms	19 ms
100 ms	9.5×10^{-2}	2.880 ± 0.116	288 ms	188 ms

Table 5 Which nuclear relaxation process enforces the criterion.

q	MC (8 ch), depolarising	MC (8 ch), dephasing
0.0	2.890	2.890
0.5	2.288	2.899
0.9	1.977	2.867
1.0	1.002	2.865

—at high field, the caveat of the preceding paragraph—but ^{14}N is quadrupolar, and at the measured flavin quadrupole couplings and a protein rotational correlation time it relaxes in ~ 0.1 – 1μ s—faster than any plausible turnover. Two of the three nuclei in our model are the flavin ^{14}N , and they are therefore wiped every cycle whatever the turnover rate. Restricting the register to the tryptophan proton leaves $\text{IPC} = 2.78$ against a floor of 2.01, an excess of 0.78 where the full register gave 2.64; the time-resolved kinetic readout, which needs all three nuclei, falls exactly to its floor (ESI, Sec. S9). The criterion survives on a proton register, at 29% of the capacity and through the CIDNP

channel alone. We regard this as the single most important correction in the present version, because it replaces an assumed relaxation time with a measured one at the cost of most of the capacity.

This is the IPC rises to a maximum of 7.2 near $T_2^e \sim \tau$ and declines for longer coherence, where the echo-state property degrades. Weak-field sensing, by contrast, improves monotonically with T_2^e . The two functions of the same molecule therefore optimise at different coherence times and cannot be maximised at once: cryptochrome ($T_2^e \sim \mu$ s) constraint that actually decides whether a radical pair can serve as a biological reservoir:

The same nuclear register must be re-used across successive turnovers, and successive turnovers must be driven by distinct input samples. The register must be carried by protons—the flavin ^{14}N is quadrupolar and cannot hold one—and must reorient with $\tau_c \leq 6$ ns, since at the geomagnetic field a proton whose dipolar vector reorients only with the protein relaxes in 2.4 ms

on the intramolecular term alone, 1.6 ms with the intermolecular bath. Given all of that, the memory horizon is $(MC - C_0) \times T_{\text{turnover}}$, bounded above by $(MC - C_0) \times T_{\text{In}}$; without it, the system retains nothing.

This criterion is a statement about a spin register. We should be exact about how much of it needed the spin dynamics. The quantity $MC - C_0 \simeq 1.9$, and with it the horizon, comes from the reservoir calculation. The constants that decide whether the horizon reaches the neural band divide unevenly. The proton T_1 of 2.4 ms and the photon budget are relaxation theory and photophysics and would have followed without the master equation. The 6.1 ms cap and the $\tau_c \leq 6$ ns requirement would not: both are obtained by feeding $q = 1 - \exp(-T_d/T_{\text{In}})$ into the measured $MC(q)$ curve, which is tuned for sensing, past its computing optimum. To our knowledge this sensing-computing trade-off, controlled by a single open-system parameter, has not previously been identified in a quantum reservoir: itself an output of the master equation. We previously claimed all four were independent of the spin dynamics; that was wrong for the last two. The population-versus-coherence finding is real, but at the operating point it changes none of those numbers. We therefore claim the spin dynamics for the capacity, not for the criterion's parameters. We should not, however, overstate the distinction we originally drew. Once q is parameterised as $q = 1 - \exp(-T_d/T_{\text{In}})$, the criterion's quantitative content is a comparison of two times, T_{turnover} against T_{In} , just as the argument it replaces compared T_{turnover} against τ . What changes is not the form of the argument but which constant appears in it, and that constant is the longer one—by about 2×10^3 for a protein-bound proton at the geomagnetic field, and by $\sim 4 \times 10^5$ for a register that reorients with $\tau_c = 0.1$ ns. The genuinely non-temporal content is the binary chemical requirement that the pair be regenerated on the same molecule.

Computing and sensing optimise at different coherence times. Reservoir IPC versus T_2^S at fixed $\tau = 50$ ns: computing peaks at $T_2^S \sim \tau$ and declines, whereas weak-field sensing improves monotonically (arrow). It is falsifiable by chemistry: it fails if the radical pair is regenerated on a fresh molecule each turnover, or if T_{In} in the diamagnetic intermediate is faster than the turnover rate. We note that neither T_{turnover} nor T_{In} has been measured in vivo for a neuronal cryptochrome, so as applied to neurons this is a conditional whose antecedents are not yet accessible. The in vitro route below is proposed because it makes both of them controllable.

2.7 The readout bottleneck

Biology has no EPR spectrometer; it reads a chemical product. Restricting to physically accessible channels,

2.7 The register is the engineered point recovers $\text{IPC} = 0.60$ (end-point yield), 2.40 (adding time-resolved kinetics) product, and 3.64 (adding CIDNP¹⁸ nuclear polarisation on the diamagnetic product). Thus the “one number” objection is false in principle—reaction kinetics alone help—but the CIDNP channel presupposes a downstream biological reader of nuclear polarisation, for which there is no direct evidence; we treat the kinetic-only value as conservative and the CIDNP value as an optimistic bound, that helps

The results above propagate to the next turnover the nuclear state of the pair that did not recombine, renormalised. That is a defensible reservoir, but it is not the object the argument of this paper is about: our claim is that the register leaves in the diamagnetic product. The two are disjoint sub-populations of each turnover. We therefore reimplemented the cycle so that the state handed forward is the nuclear density matrix actually carried away by the recombined molecule,

$$\sigma_{\text{nuc}} \propto \int_0^\tau [k_S \text{Tr}_e(P_S \rho P_S) + k_T \text{Tr}_e(P_T \rho P_T)] dt, \quad (2)$$

with the electron pair reborn unpolarised, and recomputed every readout route.

Cryptochrome

Table 6 Capacity with the survivor register against the physically faithful product register.

readout route	ch.	survivor	product
end-point singlet yield	1	1.02	1.01
time-resolved kinetics	5	2.01	2.74
both exit channels, time-resolved	10	2.01	2.76
kinetics + CIDNP on the product	8	4.65	4.91

The chemically correct register is the better one. Time-resolved kinetics rise from 2.01 to 2.74 and the CIDNP route from 4.65 to 4.91; the linear memory capacity rises correspondingly, from 1.97 to 2.44 and from 2.87 to 3.12. The conclusions of this paper are therefore conservative as originally computed. We report the survivor-register numbers throughout because they are the ones the released code produced, and give Eq. (2) as the physically faithful variant. Every result that reduces the paper's claims—the proton-only register, the collapse of the kinetic readout under turnover jitter, and the anisotropy cost—has been re-run with the product register and survives (ESI, Sec. S4). The one quantitative difference is that the kinetic readout degrades more gracefully under half a cycle of jitter with the product register, retaining 18% of its excess at half a cycle where the survivor register retains under 1%; it reaches zero near one and a half cycles. No conclusion in this work depends on the choice.

2.8 What the numbers depend on

Three sensitivities deserve to be stated, because two of them were not.

Discretisation and sampling. The linear memory capacity—the

quantity the turnover-clock claim is about—converges in the time grid: MC = 2.889, 2.866, 2.861 for $n_t = 48, 96, 192$, a change of 0.17% across the last refinement, against the 1.6% effect being claimed. The five-channel capacity is less well converged (1.96, 1.97, 1.95) and we do not base claims on its third digit. The originally submitted turnover-clock scan used a single input realisation and quoted four significant figures; that was not supportable, and the scan is repeated here with twelve.

Regularisation. The linear readout is a ridge with $\lambda = 10^{-6}$ on unstandardised features, and this choice is load-bearing: the time-resolved channels are nested partial integrals of one signal, so the Gram matrix is ill-conditioned ($\kappa \approx 10^9$), and the recovered capacity runs from 4.53 at $\lambda = 10^{-12}$ to 1.98 at $\lambda = 10^{-6}$ to 1.00 at $\lambda = 10^{-1}$ for the kinetic route. Standardising the features raises the values substantially (to 4.34 at $\lambda = 10^{-6}$). The reported capacities are thus *conservative* with respect to this choice rather than inflated by it, but they are not choice-independent, and we no longer present them as though they were.

Counting statistics. The physically meaningful regulariser is the noise on a real product pool. Adding Gaussian shot noise of relative size $N^{-1/2}$ to each channel, for a pool of N molecules, leaves the CIDNP capacity at 4.54 for $N = 10^{10}$ and 4.54 for $N = 10^8$, falling to 3.93 only at $N = 10^6$. At the copy numbers relevant to a cell the readout is not shot-noise limited. This does not address the distinct question of whether 10^{10} copies driven by a common input but not phase-locked to a common turnover schedule retain the single-molecule capacity after pooling; that is computed in Sec. S10 and reported under Limitations below, where it turns out to be the more severe of the two effects.

3 Discussion

What changed, and why it matters. Three claims that are natural to make about this system—that a scalar chemical readout is too impoverished to be useful, that a spin reservoir of a few spins beats a classical system of the same size, and that microsecond radical-pair lifetimes put biological computation out of reach—all fail when the model is driven as a real turnover and compared symmetrically. The readout is not the bottleneck; the quantum advantage is not there, and at the biologically relevant operating point no accounting we can construct puts the radical pair ahead by more than its own seed scatter; and the timescale gap is an artifact of identifying the reservoir clock with the geomagnetic field. The biologically decisive test uses the real radical pair: $J = 0$, $B = 50 \mu\text{T}$, literature hyperfine couplings ($A_{\text{N5}} \sim 14$, $A_{\text{Trp}} \sim 40 \text{ MHz}$),^{21,22} $\tau = 1 \mu\text{s}$. Notably Lee *et al.*²¹ simulate this pair at exactly these conditions, grounding our choices. The molecule is *intrinsically* a working reservoir; IPC = 5.4 with full readout (Fig. 5). But with the biologically accessible readout the capacity collapses to ≈ 0 : at $50 \mu\text{T}$ with $J = 0$ the singlet yield is nearly field-independent, so the kinetic channels are degenerate—a spin-chemical result only the master equation reveals. The remaining obstruction is the clock: the memory horizon $\text{MC} \times \tau \approx 3.2 \mu\text{s}$ (robust to the definition; ESI) is 10^3 – 10^5 times shorter than synaptic ($\sim 10 \text{ ms}$), spike-train ($\sim 100 \text{ ms}$) pair lifetime rather than the turnover interval. What remains is a chemical criterion, together with the structural fact

that the register stores populations rather than coherence and is therefore governed by T_{1n} . That fact does not, however, rescue the biological reading as we first supposed: at the geomagnetic field extreme narrowing makes T_{1n} and behavioural (seconds) timescales.⁸ The gap is two-sided: any biological input also varies only on millisecond timescales, so within one memory window the input is effectively constant. This is a quantitative, falsifiable bound: any neural reservoir-computing role for cryptochrome must live within a microsecond memory window. T_{2n} equal, and both short, so what the criterion now demands is a register that reorients fast enough to lengthen them.

The real molecule computes, but on the wrong clock. (a) IPC of a cryptochrome-realistic $\text{FAD}^{\bullet-}/\text{Trp}^{\bullet+}$ pair ($J = 0$, $B = 50 \mu\text{T}$, full readout) versus T_2^c : $\text{IPC} \approx 5.4$. (b) The memory horizon $\text{MC} \times \tau \approx 3 \mu\text{s}$ is 10^3 – 10^5 times shorter than neural timescales (log scale).

Relation to prior work. Quantum reservoir computing in spin ensembles is established, including the use of dissipation as a resource and the exponential-state-space argument.^{13,27–30} We do not claim these as new, and—having found no advantage under a symmetric comparison—we do not add to them. The question of whether nuclear polarisation accumulates usefully across successive cryptochrome photocycles has been asked before, by Wong *et al.*³⁶, who found that the gain available in an idealised model does not survive conditions approaching those in vivo. Our criterion is the chemical statement of what would have to be true for it to survive, and our conclusions are consistent with theirs: the reservoir works only in a narrow and specific corner of parameter space. Photo-CIDNP itself has been observed in a flavoprotein⁴¹ though not yet in cryptochrome, and the underlying polarisation mechanism is standard¹⁸. Our contribution is to take a chemically realised radical pair, drive and read it as chemistry actually would, and convert a long-standing qualitative conjecture^{6,7} into a criterion that experiment can test. The decoherence objection usually cited against such proposals⁹ does not apply directly here: it evaluates superpositions of ion positions and tubulin dipole states, not nuclear spins, which is precisely why nuclear-spin proposals were advanced in reply to it. The objection that does apply is the one this work inherits—nuclear relaxation in the diamagnetic product—and the relevant history is discouraging. The long ^{31}P coherence times underpinning Ref. 6 are theoretical estimates rather than measurements, and have since been bounded at 37 min and expected at “a few seconds” by spin-dynamics calculation³⁷, while ^{31}P singlet order in pyrophosphate is measured to relax faster than T_1 ³⁴. We therefore ground our own relaxation assumption on measured proton T_1 values in proteins rather than on estimate, and restrict the register accordingly (ESI, Sec. S9). The result also sharpens why CIDNP matters here: in a separated pair it is not an optional extra channel but the mechanism by which information reaches the nuclei at all.

3.1 The feasibility frontier and the framework

Assembling the results,

Limitations. The model is a five-spin reduction. Two of its simplifications have now been relaxed, with opposite outcomes. Treating ^{14}N correctly as spin-1 rather than spin-1/2 increases the excess capacity by 28%, so that approximation was conservative. Anisotropy is not: scanning an axial hyperfine tensor at fixed isotropic average costs roughly two-thirds of the capacity by $\eta = 0.5$, and the capacity vanishes exactly at the Ising point where the perpendicular components go to zero, which identifies those components as what carries the memory (ESI, Sec. S11). That scan is not a calculation with the published tensor sets of the cryptochrome pair^{21,22}, which we still do not treat, so the ~~separated radical pair clears the physics bar for QRC (fading memory, nonlinear capacity, 72% coherence-derived), but biological use requires (limitation is narrowed rather than removed—but the direction is unfavourable and we report it as such. The capacity estimator uses degree-2 targets, so higher-order nonlinear capacity is not counted. Whether a neuronal cryptochrome forms a separated radical pair at all is unestablished—CRY1/CRY2 are primarily circadian transcriptional regulators¹⁵—so the biological reading of our criterion is conditional. The spin-dynamics results do not depend on that condition.~~

Four further conditions bound the biological reading. Three we have now quantified, and each cost the paper something.

(i) ~~multichannel spin-resolved readout (a single scalar gives $\text{IPC} \lesssim 1$), (ii) timescale matching (the binding constraint: microsecond memory versus millisecond-second neural dynamics), and (iii) a mechanism to exploit ensemble heterogeneity. The realistic, mechanism-agnostic estimate is a small but nonzero reservoir, $\text{IPC} \approx 1.5$ Which nuclei hold the register. Not all of them. Nuclear dephasing turns out to be irrelevant—the register is a population register—so the binding constant is $T_{1\rho}$ as assumed. But $T_{1\rho}$ is not one number. Protein-bound ^1H relaxes on the second timescale in the measurements available³⁵—all of them at NMR field strengths, a caveat item (iv) below makes decisive—whereas ^{14}N is quadrupolar and, at the measured flavin quadrupole couplings and a protein correlation time, relaxes in ~ 0.2 – 3.5 μs . The capacity/readout/memory horizon lens is a reusable framework: applied to the active-site flavoenzymes²⁴ it excludes them at the first axis (picosecond horizon); applied to Fisher's Posner-molecule proposal,⁶ where ^{31}P nuclear spins carry second-scale coherence, the timescale test is passed and the binding constraint shifts to entanglement and readout. The same three axes locate the decisive bottleneck case by case. $0.3 \mu\text{s}$ at the geomagnetic field and in single microseconds even at 9.4 T ^{38,39}—faster than any physiological turnover at either. Two of our three nuclei are ^{14}N , and they cannot hold a register at any physiological turnover rate. Restricting the register to the proton leaves 29% of the excess capacity and destroys the time-resolved kinetic readout entirely. Of the two routes we tested under this restriction, CIDNP is the one that survives; the other five routes were not re-evaluated with a restricted register (ESI, Sec. S9). The turnover-clock argument survives, on measured rather than assumed relaxation data; the capacity does not survive intact.~~

3.1 Relation to prior work and outlook

~~Spin-based QRC and the dissipation-as-resource and exponential-state-space arguments are established;^{27–30}~~

(ii) *Ensemble averaging.* Pooling $\sim 10^{10}$ copies is safe against heterogeneity—of the couplings and, separately, of the relaxation rate—but not against desynchronisation. Letting τ_c vary log-normally over a sixteen-fold range, so that copies retain different fractions of their register, leaves the CIDNP excess at 2.04 against 1.96 for a uniform pool: capacity does not fall, and the whole effect is under five per cent. A mean-field control at the pool's own mean survival accounts for only a third of even that small difference, so a heterogeneous pool is not simply a homogeneous one at its mean $T_{1\rho}$; a single effective $T_{1\rho}$ is a working approximation rather than an identity (ESI, Sec. S10). The kinetic readout's memory falls away with the spread of copy phases: with the product register its excess declines from 1.72 at zero jitter to 0.32 at half a cycle, 0.18 at one cycle and 0.02 at one and a half, and with the survivor register it is already at 0.01 by half a cycle. Two cautions attach to reading those numbers as a smooth decay. The survivor sequence is not monotonic—it returns to 0.11 at one cycle—and the register-wiped floor itself rises steeply beyond one cycle (from 1.01 to 1.86 for the product register), because pooling copies that sit whole cycles apart is itself a delay line. The excess is the right quantity precisely because it subtracts that, but the underlying capacities do not fall monotonically and we do not claim these as new. What is new is (i) a chemically realised radical-pair reservoir benchmarked against explicit classical baselines, (ii) the sensing-computing coherence-time trade-off in a molecule that doubles as a calibrated field sensor; they do. CIDNP retains 40–49% of its excess even at two cycles of jitter, depending on the register, because it integrates over the cycle rather than sampling it (Sec. S10). Synchronised turnover is thus a requirement of the kinetic readout specifically. Both restrictions—nuclide and (iii) ~~desynchronisation~~—leave CIDNP as the surviving one of the two routes tested under each.

(iii) *Readout signal-to-noise.* The ridge parameter is an implicit noise model and the recovered capacity depends on it (Sec. 2.8); the reported values are conservative with respect to that choice, and adding Poisson shot noise for a pool of 10^8 molecules or more leaves them essentially unchanged.

(iv) *Falsifiability.* Neither T_{turnover} nor $T_{1\rho}$ has been measured for a neuronal cryptochrome, but both can be predicted from measured constants, and doing so sharpens the criterion into something testable without a neuron (ESI, Sec. S12). Two results follow, and the first is adverse. Nuclear relaxation at the geomagnetic field is in the ~~conversion of the quantum-brain conjecture into a falsifiable timescale bound plus a transferable framework. The quantum-information content is testable now in vitro: covalent extreme-narrowing regime, where $1/T_1 \propto \tau_c$ is maximal, so the second-long $T_{1\rho}$ we assumed is a high-field value: a proton on a 60 kDa protein relaxes in 2.4 ms at $50 \mu\text{T}$ against 9.0 s at 9.4 T. What sets the ceiling is not whether the register is chemically bound to the protein but the effective correlation time of its dipolar vector, $\tau_{\text{eff}} = S^2 \tau_c + (1 - S^2) \tau_{\text{int}}$,~~

together with the dipolar sum $\Sigma = \sum_j S_j^2 r_j^{-6}$ over intramolecular and intermolecular partners. For a vector that reorients only with the intact 60 kDa protein ($S^2 = 1$, $\tau_c = 15.2$ ns) the intramolecular geminal term alone gives $T_1 \leq 2.4$ ms and a ceiling of < 6.1 ms; adding the intermolecular proton bath—whose size we calibrate on the water row of our own validation table, at 54% of the intramolecular rate—shortens these to 1.6 and 4.0 ms. That row is an empirical anchor rather than an independent check: f is defined as its residual, so the agreement it then shows is an identity. Fast internal motion lengthens T_1 : the intramethyl H-H vector of the flavin $8\alpha\text{-CH}_3$ has the exact geometric value $S^2 = 1/4$, because that vector does *not* co-rotate rigidly with the protein even though the methyl is covalently attached, and the intramethyl coupling alone would give $T_1 = 5.0$ ms and a ceiling of 11.3 ms—inside the band. That escape does not survive the bath: it moves the methyl's own boundary from 19.3 to 10.8 ns. The criterion is therefore quantitative and vector-specific rather than a statement about binding: the register's dipolar vector must lose orientational memory about two and a half times faster than the intact protein ($\tau_{\text{eff}} \lesssim 6$ ns), a factor that would be 1.6 if the intermolecular bath were neglected. Second, the drive cannot be light: with the measured flavin extinction coefficient, a single flavin turns over once per 7.7 s in full sunlight and once per some hundreds of years inside a skull, so a photochemical cycle fails by about ten orders of magnitude and the regeneration must be catalytic. The criterion that survives is correspondingly specific—the pair regenerated catalytically on a register with $\tau_c \lesssim 6$ ns, at a turnover interval that depends on how mobile that register is—9.8–27 ms at $\tau_c = 5$ ns, widening to 5.6–134 ms at 1 ns and 5.4–1340 ms at 0.1 ns—and both of its parameters are measurable in vitro by relaxation dispersion and steady-state kinetics.

What would settle it. The criterion is addressable in vitro before it is addressable in a neuron. Covalently linked donor–bridge–acceptor radical pairs have synthetically tunable J and τ exchange coupling and lifetime and are read by transient EPR and optical/CIDNP spectroscopy,^{2,14,42} so the predicted IPC ≈ 7 , the 72% coherence fraction, and the $T_2^s \sim \tau$ optimum are all directly measurable.^{2,14,42} Repeatedly cycling such a system and asking whether the product yield of turnover k depends on the input at turnover $k-1$, $k-2$, ... measures the memory horizon directly, and varying the interval between cycles tests the register-reuse criterion where it is sharpest.

4 Conclusions

A cryptochrome

Driven and read as a chemical turnover rather than as an abstract quantum channel, a cryptochrome-type radical pair is intrinsically a capable quantum reservoir (IPC = 6.9 engineered, 5.4 at the geomagnetic field; 72% coherence-derived) but is biologically non-functional as one: the accessible product readout yields ≈ 0 capacity and the microsecond memory horizon falls 10^3 a reservoir whose output ordinary chemistry can recover: time-resolved product kinetics give IPC = 2.0 against a floor of 1.0, and the nuclear polarisation carried off by the product

reaches 4.7 against a floor of 2.0, with no spectroscopic access required. Under a specific matched-hopping incoherent limit 82% of that capacity is lost, and the loss is attributable to the electrons rather than to erasure of the nuclear register—and yet there is no quantum advantage: at the biologically relevant operating point a classical network matched channel-for-channel exceeds the radical pair. The obstacle usually invoked, a microsecond memory horizon, does not survive scrutiny: the register is nuclear and survives recombination, so the clock is the turnover interval, and a 10^{-10} below every neural timescale. We thus place a quantitative, falsifiable bound on the reservoir-computing route to a “quantum brain” and supply a reusable capacity/readout/timescale framework for any 100 ms turnover places the horizon at 19–188 ms—so long as the register itself lasts that long, which for a vector reorienting only with the whole protein nucleus at the geomagnetic field it does not. What decides the question instead is whether the same nuclear register is reused from one turnover to the next; if it is not, the capacity falls to a memoryless MC = 1. Because that register stores its information in nuclear populations rather than in nuclear coherence, the constant that bounds it is a nuclear T_1 . That distinction, which we first took to be what rescued the horizon, in fact buys nothing at the operating point: the geomagnetic field puts the register in extreme narrowing, where T_{1n} and T_{2n} are equal. It cannot, however, be any nucleus. The flavin ^{14}N is quadrupolar and relaxes in microseconds, so the register must be carried by protons; that restriction removes seven tenths of the excess capacity—the IPC itself falls by two fifths—and leaves the product's nuclear polarisation as the only readout we tested that still works, a conclusion reached independently by asking what survives when the copies in a 10^{10} molecule pool fall out of step. We offer the criterion as the thing to test, in the sharpened form that the relevant constant is the proton T_1 of the diamagnetic product at the geomagnetic field, which our own calculation puts at 2.4 ms—1.6 ms once the intermolecular bath is restored—for a nucleus whose dipolar vector reorients only with the protein, rather than at the second suggested by high-field measurements—so that the register must reorient at least two and a half times faster than the protein for the horizon to reach the neural band at all. The turnover interval of a neuronal cryptochrome has not been measured.

A Model and methods

A.1 Reservoir Hamiltonian

We model the reservoir as $N = 5$ spins—two radical-pair system. The same spin chemistry that makes cryptochrome a magnetosensor makes it electrons (e_1, e_2) and three spin-1/2 nuclei—with Hamiltonian (in MHz). At the cryptochrome point the three nuclei carry the hyperfine couplings of $^{14}\text{N}5$ and $^{14}\text{N}10$ on FAD $^{\bullet-}$ and H β 1 on TrpH $^{\bullet+}$ (ESI, Table S19), with ^{14}N modelled as spin-1/2; no ^{31}P is present in the cryptochrome model, and the ^{31}P -based proposal of Ref. ⁶ is therefore *not* the

system tested here.

$$H = \omega_e (S_z^{e1} + S_z^{e2}) + \sum_{\mu=x,y,z} \left[A_P S_\mu^{e1} I_\mu^P + A_{H1} S_\mu^{e1} I_\mu^{H1} + A_{e2} S_\mu^{e2} I_\mu^{H2} + J S_\mu^{e1} S_\mu^{e2} \right], \quad (3)$$

where $\omega_e = g_e \mu_B B / h$ is the electron Zeeman frequency. Each radical carries its own nuclei—they are not shared, which is what a spatially separated pair means—and the exchange J links the two electrons. At the engineered operating point we use $A_{e1a} = 80$, $A_{e1b} = 40$, $A_{e2a} = 15$ MHz, $J = 2$ MHz and $B = 1$ mT; representative molecular-spin-RC values chosen to give effective hyperfine writing, not a fit to a specific protein. For the cryptochrome point we set $J = 0$, $B = 50$ μ T and $\tau = 1$ μ s, with hyperfine magnitudes representative of the dominant FAD $^{\bullet-}$ and TrpH $^{\bullet+}$ couplings^{21,22}. The capacity is not fine-tuned to these numbers: at the engineered point it varies by under 2% over $T_2^e = 50$ –2000 ns, and that insensitivity is estimator-robust (a full-rank readout gives -1.7% over the same range). Below the operating point the two estimators disagree: the ridge readout recovers a steep fall (-47% from 50 to 5 ns), but this is an artifact of the fixed absolute ridge penalising the small-amplitude features at short T_2^e — a quantum reservoir, with decoherence the enemy of full-rank readout gives $+2.5\%$ for the same comparison (cf. the Regularisation paragraph of Sec. 2.8). The insensitivity claimed here is therefore to T_2^e at or above τ , and we make no claim below the operating point. The register criterion of Sec. 2.6, however, is *not* independent of the hyperfine assignment—it is carried by the perpendicular tensor components and vanishes with them (ESI, Sec. S11).

A.2 Master equation and propagator

Each reservoir update is the completely positive, trace-preserving map $\rho \mapsto e^{\mathcal{L}\tau} \rho$ generated by the Haberkorn master equation³¹ (whose spin-selective recombination operator we adopt in its standard form¹⁷) with electronic pure dephasing,

$$\frac{d\rho}{dt} = -i[H, \rho] - \frac{1}{2} \{k_S P_S + k_T P_T, \rho\} + \sum_i \frac{2}{T_2^e} \left(S_z^i \rho S_z^i - \frac{1}{4} \rho \right), \quad (4)$$

solved exactly in Liouville space. Couplings are quoted in MHz throughout; H is multiplied by 2π before propagation, so that $e^{\mathcal{L}\tau}$ acts in angular frequency (rad μ s $^{-1}$) with τ in μ s, and the rates k_S , k_T and $2/T_2^e$ are in μ s $^{-1}$. The convention is pinned to physics rather than to internal consistency: a lone electron propagated by the same code at 1 mT precesses at $g_e \mu_B B / h = 28.0$ MHz (test_units.py). The coefficient $2/T_2^e$ makes the first function and single-electron transverse coherence decay as e^{-t/T_2^e} , so T_2^e is the single-electron transverse relaxation time in the usual sense. The third term is the electronic decoherence that the companion sensing analysis showed to be decisive; it is not,

however, what supplies the fading memory here, which comes from the openness of the chemical cycle and from inter-cycle nuclear depolarisation. We verified that, in the limit $1/T_2^e \rightarrow 0$ and $k_S = k_T$, the Liouville-space solver reproduces the closed-form coherent Green's-function singlet yield to better than 10^{-9} (ESI). Whether recombination acts within the update step depends on which result is being computed; the two conventions are set out under “Reservoir protocol” below.

A.3 Reservoir protocol

We adapt the temporal-QRC protocol of Ref.²⁷, but the drive is *not* the friend of the second—a duality testable in synthetic radical pairs independently of any neurobiological claim—single-electron reset used there. At step k the nuclei are retained and the electron pair is recreated as a spin-correlated pair whose singlet character encodes the input,

$$\rho_e(s_k) = s_k |S\rangle\langle S| + (1 - s_k) P_T / 3, \quad (5)$$

so that the reservoir state entering cycle k is $\rho_e(s_k) \otimes \sigma_{\text{nuc}}^{(k)}$. The state then evolves for the pair lifetime τ under $e^{\mathcal{L}\tau}$; a set of observables is read out, and a linear (ridge) readout is trained on them. Inputs s_k are i.i.d. uniform on $[0, 1]$; the first 80–100 steps are discarded as washout. The washout discards initial transients, so the reported capacities are independent of the initial spin state—a consequence of the fading-memory (echo-state) property.

Data availability

Two operating points are used and must not be conflated. The engineered point ($J = 2$ MHz, $B = 1$ mT, $\tau = T_2^e = 50$ ns) is a molecular-spin-RC reference and is where the twelve-observable capacity, the coherence fraction and the classical baselines of Secs. 2 are evaluated; it lies outside the weak-field separated-pair regime that motivates the biological reading, and we flag every number taken there. The cryptochrome point ($J = 0$, $B = 50$ μ T, $\tau = 1$ μ s, $T_2^e = 1$ μ s) is where the chemically accessible readouts, the turnover clock and the register criterion are evaluated. Recombination is likewise not uniform across the paper: the twelve-observable capacity is computed with $k_S = k_T = 0$ within the update step, whereas every chemical-readout, clock and criterion result is computed with spin-selective recombination acting throughout the cycle ($k_S = 1$, $k_T = 0.2$ μ s $^{-1}$).

The single-electron reset of Ref.²⁷, in which only the input electron is replaced by a product state while the reduced state of the remainder is retained, is used in this work *only* as a control. It is not a chemical process, and for a separated pair ($J = 0$) it destroys the electron–electron correlation every cycle: since $\langle P_S \rangle = \frac{1}{4} - \langle \mathbf{S}_1 \cdot \mathbf{S}_2 \rangle$ with $\langle \mathbf{S}_2 \rangle = 0$, the singlet probability is pinned at exactly $1/4$ independently of the input, and the measured capacity is identically zero (ESI, Sec. S3). All results reported below use Eq. (5).

A.4 Capacity measures

We quantify the reservoir with two information-theoretic

benchmarks. The linear *memory capacity*³² $MC = \sum_{d \geq 0} r^2(s_{k-d})$ sums the squared multiple correlation for reconstructing delayed inputs. The *information processing capacity*³³ adds nonlinear functions of the input history, $IPC = MC + \sum_{\text{nonlinear}} C$, evaluated here through degree-2 Legendre targets (diagonal $P_2(s_{k-d})$ and cross $s_{k-d_1}s_{k-d_2}$ terms). A fundamental theorem³³ bounds the total capacity by the number of linearly independent readout observables, $IPC \leq n_{\text{obs}}$; we use this bound throughout as a correctness anchor. It is an asymptotic bound for an orthonormal target family and is violated by a few per cent in finite samples, so satisfying it does *not* demonstrate that the readout observables are independent, and we make no such argument anywhere (ESI, Sec. S2).

All results are ~~produced by the open-source reproducible from the open~~ qbscreen package (MIT licence; ~~); the module list is given under Data and code availability below. We note there that ensemble.py, quantum_vs_classical.py and the run_reservoir driver in reservoir.py still implement the superseded single-electron reset and are retained only as controls, so no result reported here is taken from them; the capacity estimators in reservoir.py are used throughout (293/293 tests pass).~~

Data and code availability

All analyses are implemented in the open qbscreen package, https://github.com/deeptell-inc/new_Q_brain. The corrected correlated-pair-birth injection is in qbscreen/corrected_injection.py; the chemically accessible readout routes, the turnover-clock scan and the ~~numerical outputs underlying every figure and table are provided as JSON files in the repository,~~ register-reuse criterion are in qbscreen/readout_routes.py; the protocol-consistent capacity, coherence fraction and classical baselines are in qbscreen/final_numbers.py; the master-equation solver is in qbscreen/master_equation.py. Numerical outputs are in simulation_results/. The capacity bound and the agreement of the solver with the closed-form coherent limit are checked in qbscreen/tests/ (293 tests, all passing). Twenty-nine tests in test_panel_claims.py bind the headline quantities and load-bearing claims, and a further one hundred and forty in test_table_rows.py bind the capacity and standard-deviation cells of every table in the main text and the ESI, with the expected values transcribed from the manuscript rather than from the data files. The opposite direction is checked by test_printed_cells.py, which parses the numeric cells back out of the .tex: the headline tables are matched column by column against their data files, and every other printed table cell must equal some shipped value. The suite therefore fails if either side moves. The spin-parameter table (S19) is now bound as well, by test_spin_parameters.py: it parses the printed values and compares them to the dictionaries the Hamiltonian is built from, checking the millitesla column as the megahertz column divided by the electron gyromagnetic ratio rather than as an independent input. Derived-time and ~~the central claims are covered by an automated test suite. A standalone Data Availability Statement accompanies this submission. The study is~~

~~entirely computational; all capacities are reported out of sample.~~ raw-IPC columns remain unbound. Hand-typed cross-document references to supplement tables, which $\LaTeX$ cannot check across separate documents, are bound against supplementary.aux by test_cross_document_refs.py. This closes the drift that produced the error the present revision began with, in which a figure quoted from the five-channel column was printed beside the eight-channel table. The $\LaTeX$ sources these tests read accompany the submission with the freeze manifest and are not distributed in the public repository; on a clone those tests skip rather than fail.

Conflicts of interest

There are no conflicts to declare.

Author contributions

H.W. conceived the study, developed the master-equation and reservoir code, ran the simulations, and wrote the manuscript. T.T. contributed to the spin-dynamics formulation and the analysis of the nuclear-register criterion. Both authors reviewed the results and approved the manuscript.

Acknowledgements

We thank the reviewers of the companion sensing study, whose insistence on a decoherence-explicit treatment shaped the methodology used here. All [numerical](#) results are reproducible from the open-source qbscreen package.

Notes and references

- 1 U. E. Steiner and T. Ulrich, *Magnetic field effects in chemical kinetics and related phenomena*, Chem. Rev. **89**, 51 (1989).
- 2 P. J. Hore and H. Mouritsen, *The radical-pair mechanism of magnetoreception*, Annu. Rev. Biophys. **45**, 299 (2016).
- 3 T. Ritz, S. Adem, and K. Schulten, *A model for photoreceptor-based magnetoreception in birds*, Biophys. J. **78**, 707 (2000).
- 4 K. Maeda *et al.*, *Magnetically sensitive light-induced reactions in cryptochrome are consistent with its proposed role as a magnetoreceptor*, Proc. Natl. Acad. Sci. USA **109**, 4774 (2012).
- 5 J. Xu *et al.*, *Magnetic sensitivity of cryptochrome 4 from a migratory songbird*, Nature **594**, 535 (2021).
- 6 M. P. A. Fisher, *Quantum cognition: The possibility of processing with nuclear spins in the brain*, Ann. Phys. **362**, 593 (2015).
- 7 C. M. Kerskens and D. López Pérez, *Experimental indications of non-classical brain functions*, J. Phys. Commun. **6**, 105001 (2022).
- 8 G. Buzsáki and A. Draguhn, *Neuronal oscillations in cortical networks*, Science **304**, 1926 (2004).
- 9 M. Tegmark, *Importance of quantum decoherence in brain processes*, Phys. Rev. E **61**, 4194 (2000).
- 10 N. Lambert, Y.-N. Chen, Y.-C. Cheng, C.-M. Li, G.-Y. Chen, and F. Nori, *Quantum biology*, Nat. Phys. **9**, 10 (2013).

- 11 [J. Cao et al., Quantum biology revisited, Sci. Adv. **6**, eaaz4888 \(2020\).](#)
- 12 [H. G. Hiscock et al., The quantum needle of the avian magnetic compass, Proc. Natl. Acad. Sci. USA **113**, 4634 \(2016\).](#)
- 13 [G. Tanaka et al., Recent advances in physical reservoir computing: A review, Neural Netw. **115**, 100 \(2019\).](#)
- 14 [M. R. Wasielewski et al., Exploiting chemistry and molecular systems for quantum information science, Nat. Rev. Chem. **4**, 490 \(2020\).](#)
- 15 [C. L. Partch, C. B. Green, and J. S. Takahashi, Molecular architecture of the mammalian circadian clock, Trends Cell Biol. **24**, 90 \(2014\).](#)
- 16 [C. R. Timmel, U. Till, B. Brocklehurst, K. A. McLauchlan, and P. J. Hore, Effects of weak magnetic fields on free radical recombination reactions, Mol. Phys. **95**, 71 \(1998\).](#)
- 17 [J. A. Jones and P. J. Hore, Spin-selective reactions of radical pairs act as quantum measurements, Chem. Phys. Lett. **488**, 90 \(2010\).](#)
- 18 [P. J. Hore and R. W. Broadhurst, Photo-CIDNP of biopolymers, Prog. Nucl. Magn. Reson. Spectrosc. **25**, 345 \(1993\).](#)
- 19 [H. Wakaura, Systematic screening of brain neurotransmitter enzymes for radical-pair mechanism competence identifies candidates for magnetic field sensitivity, Research Square preprint, doi:10.21203/rs.3.rs-9278975/v1 \(2025\).](#)
- 20 [H. Wakaura, Motional narrowing rescues entanglement in biological radical-pair systems: quantum error correction and reservoir computing beyond the breaking threshold, Research Square preprint, doi:10.21203/rs.3.rs-9493052/v1 \(2026\).](#)
- 21 [A. A. Lee, J. C. S. Lau, H. J. Hogben, T. Biskup, D. R. Kattinig, and P. J. Hore, Alternative radical pairs for cryptochrome-based magnetoreception, J. R. Soc. Interface **11**, 20131063 \(2014\).](#)
- 22 [E. Schleicher et al., Selective \$^{13}\text{C}\$ labelling reveals the electronic structure of flavocoenzyme radicals, Sci. Rep. **11**, 18395 \(2021\).](#)
- 23 [O. Efimova and P. J. Hore, Role of exchange and dipolar interactions in the radical pair model of the avian magnetic compass, Biophys. J. **94**, 1565 \(2008\).](#)
- 24 [H. Wakaura and T. Tanimae, Active-site geometry, not the shared flavin cofactor, sets the weak-field magnetosensitivity of neurotransmitter-metabolising flavoenzymes \(companion manuscript, 2026\).](#)
- 25 [W. Maass, T. Natschläger, and H. Markram, Real-time computing without stable states: A new framework for neural computation based on perturbations, Neural Comput. **14**, 2531 \(2002\).](#)
- 26 [H. Jaeger and H. Haas, Harnessing nonlinearity: Predicting chaotic systems and saving energy in wireless communication, Science **304**, 78 \(2004\).](#)
- 27 [K. Fujii and K. Nakajima, Harnessing disordered-ensemble quantum dynamics for machine learning, Phys. Rev. Applied **8**, 024030 \(2017\).](#)
- 28 [K. Nakajima et al., Boosting computational power through spatial multiplexing in quantum reservoir computing, Phys. Rev. Applied **11**, 034021 \(2019\).](#)
- 29 [R. Martínez-Peña et al., Dynamical phase transitions in quantum reservoir computing, Phys. Rev. Lett. **127**, 100502 \(2021\).](#)
- 30 [P. Mujal, R. Martínez-Peña, J. Nokkala, J. García-Bení, G. L. Giorgi, M. C. Soriano, and R. Zambrini, Opportunities in quantum reservoir computing and extreme learning machines, Adv. Quantum Technol. **4**, 2100027 \(2021\).](#)
- 31 [R. Haberkorn, Density matrix description of spin-selective radical pair reactions, Mol. Phys. **32**, 1491 \(1976\).](#)
- 32 [H. Jaeger, Short term memory in echo state networks, GMD Report 152, German National Research Center for Information Technology \(2002\).](#)
- 33 [J. Dambre, D. Verstraeten, B. Schrauwen, and S. Massar, Information processing capacity of dynamical systems, Sci. Rep. **2**, 514 \(2012\).](#)
- 34 [D. E. Korenchan, J. Lu, M. H. Levitt, and A. Jerschow, \$^{31}\text{P}\$ nuclear spin singlet lifetimes in a system with switchable magnetic inequivalence: experiment and simulation, Phys. Chem. Chem. Phys. **23**, 19465 \(2021\).](#)
- 35 [N. Tjandra, S. E. Feller, R. W. Pastor, and A. Bax, Rotational diffusion anisotropy of human ubiquitin from \$^{15}\text{N}\$ NMR relaxation, J. Am. Chem. Soc. **117**, 12562 \(1995\).](#)
- 36 [S. Y. Wong, I. A. Solov'yov, P. J. Hore, and D. R. Kattinig, Nuclear polarization effects in cryptochrome-based magnetoreception, J. Chem. Phys. **154**, 035102 \(2021\).](#)
- 37 [T. C. Player and P. J. Hore, Posner qubits: spin dynamics of entangled \$\text{Ca}_9\(\text{PO}_4\)_6\$ molecules and their role in neural processing, J. R. Soc. Interface **15**, 20180494 \(2018\).](#)
- 38 [Y. A. Goddard, J.-P. Korb, and R. G. Bryant, Water molecule contributions to proton spin-lattice relaxation in rotationally immobilized proteins, J. Magn. Reson. **199**, 68 \(2009\).](#)
- 39 [E. P. Sunde and B. Halle, Mechanism of \$^1\text{H}\$ - \$^{14}\text{N}\$ cross-relaxation in immobilized proteins, J. Magn. Reson. **203**, 257 \(2010\).](#)
- 40 [W. Bogner et al., Assessment of \$^{31}\text{P}\$ relaxation times in the human calf muscle: a comparison between 3 T and 7 T in vivo, Magn. Reson. Med. **62**, 574 \(2009\).](#)
- 41 [S. S. Thamarath, J. Heberle, P. J. Hore, T. Kottke, and J. Matysik, Solid-state photo-CIDNP effect observed in phototropin LOV1-C57S by \$^{13}\text{C}\$ magic-angle spinning NMR spectroscopy, J. Am. Chem. Soc. **132**, 15542 \(2010\).](#)
- 42 [M. R. Wasielewski, Energy, charge, and spin transport in molecules and self-assembled nanostructures inspired by photosynthesis, J. Org. Chem. **71**, 5051 \(2006\).](#)