

## Feedback — Module 3 Quiz

[Help](#)

You submitted this quiz on **Thu 10 Apr 2014 7:44 PM PDT**. You got a score of **11.00** out of **11.00**.

### Question 1

In a new population, a sample of 9 men yielded a sample average brain volume of 1,100cc and a standard deviation of 30cc. What is a 95% Student's T confidence interval for the mean brain volume in this new population?

Your Answer	Score	Explanation
<input checked="" type="radio"/> About 1,077 to 1,123 cc	✓ 1.00	
<input type="radio"/> About 1,080 to 1,120 cc		
<input type="radio"/> About 1,131 to 1,169 cc		
<input type="radio"/> About 1,092 to 1,108		
Total	1.00 / 1.00	

### Question 2

A diet pill is given to 9 subjects over six weeks. The average difference in weight (follow up - baseline) is -2 pounds. What would the standard deviation have to be for the 95% T confidence interval to lie entirely below 0?

Your Answer	Score	Explanation
<input type="radio"/> Around 2.8 pounds or less		

☐ Around 2.1 pounds or less

☐ Around 2.4 pounds or less

☐ Around 3.0 pounds or less

☐ Around 2.0 pounds or less.

☐ Around 2.2 pounds or less

☐ Around 2.3 pounds or less

☐ Around 2.5 pounds or less

☐ Around 2.9 pounds or less

☐ Around 2.7 pounds or less

☒ Around 2.6 pounds or less ✓ 1.00

Total 1.00 / 1.00

## Question 3

Refer to the previous question. The interval would up being  $[-3.5, -0.5]$  pounds. What can be said about the population mean weight loss at 95% confidence?

**Your Answer**

**Score**

**Explanation**

☐ We can not rule out the possibility of no mean weight loss at 95% confidence.

☒ There is support at 95% confidence of mean weight loss. ✓ 1.00

☐ We can not rule out the possibility of mean weight gain at 95% confidence.

☐ There is support of mean weight gain at 95% confidence.

Total 1.00 /

1.00

## Question 4

In an effort to improve efficiency, hospital administrators are evaluating a new triage system for their emergency room. In an validation study of the system, 5 patients were tracked in a mock ER under both the new and old triage system. Their waiting times were recorded. Would it be better to use an independent group or paired T confidence interval in this setting?

Your Answer	Score	Explanation
<input checked="" type="radio"/> A paired interval	✓ 1.00	
<input type="radio"/> Independent group interval		
Total	1.00 / 1.00	

## Question 5

Refer to the setting of the previous question. To further test the system, administrators selected 20 nights and randomly assigned the new triage system to be used on 10 nights and the standard system on the remaining 10 nights. They calculated the nightly median waiting time (MWT) to see a physician. The average MWT for the new system was 3 hours with a variance of 0.60 while the average MWT for the old system was 5 hours with a variance of 0.68. Give a 95% confidence interval estimate for the differences of the mean MWT associated with the new system. Assume a constant variance.

Your Answer	Score	Explanation
<input type="radio"/> A mean decrease of between 3.68 and 0.32 hours.		
<input checked="" type="radio"/> A mean decrease of between 2.75 and 1.25 hours.	✓ 1.00	
<input type="radio"/> A mean decrease of between 2.60 and 1.40 hours.		
Total	1.00 / 1.00	

## Question 6

Suppose that you create a 95% T confidence interval. You then create a 90% interval using the same data. What can be said about the 90% interval with respect to the 95% interval?

Your Answer	Score	Explanation
<input type="radio"/> The interval will be the same width.		
<input checked="" type="radio"/> The interval will be narrower.	✓ 1.00	
<input type="radio"/> The interval will be wider.		
<input type="radio"/> The interval will be centered at a different number.		
Total	1.00 / 1.00	

## Question 7

Let distribution 1 be  $N(\mu_1, \sigma_1^2)$  and distribution 2 be  $N(\mu_2, \sigma_2^2)$ . Let  $x_{1,\alpha}$  and  $x_{2,\alpha}$  be the  $\alpha^{th}$  quantile from the two distributions, respectively. How are the two mathematically related?

Your Answer	Score	Explanation
<input checked="" type="radio"/> They are related as a line $x_{2,\alpha} = \mu_2 - \frac{\sigma_2}{\sigma_1} \mu_1 + \frac{\sigma_2}{\sigma_1} x_{1,\alpha}$	✓ 1.00	
<input type="radio"/> They are related via a parabola $x_{1,\alpha} = x_{2,\alpha}^2$		
<input type="radio"/> They are related by the identity line $x_{1,\alpha} = x_{2,\alpha}$		
<input type="radio"/> There is no closed form relationship.		
Total	1.00 / 1.00	

## Question 8

Consider data points  $x_1, \dots, x_n$ . Imagine a probability mass function, so that a random variable  $X$  from this distribution has  $P(X = x_i) = p(x_i) = \frac{1}{n}$ . This is the so-called bootstrap distribution. What is the mean of the bootstrap distribution?

Your Answer	Score	Explanation
<input type="radio"/> The sample median of the data		
<input type="radio"/> The sample variance of the data		
<input type="radio"/> It equals 1		
<input checked="" type="radio"/> The sample mean of the data	✓ 1.00	
Total	1.00 / 1.00	

## Question 9

Suppose we were to simulate a large number of standard normal random variables and a large number of exponential random variables. What would a plot of the exponential quantiles (horizontal axis) versus the standard normal quantiles (vertical axis) look like? Let  $\Phi$  be the standard normal distribution function.

Your Answer	Score	Explanation
<input checked="" type="radio"/> It will look like the function $f(x) = \Phi^{-1}(1 - e^{-x})$ for $x > 0$ .	✓ 1.00	
<input type="radio"/> It will look like a straight line		
<input type="radio"/> It will look like the function $f(x) = -\Phi^{-1}(-\log(1 - x))$ for $0 \leq x \leq 1$ .		
<input type="radio"/> It will look like the function $f(x) = x$ for $0 < x$		
Total	1.00 / 1.00	

## Question 10

Let  $F(x)$  be a distribution function. Notice that  $G(x) = F(a^2x + b)$  is also a distribution function for  $a \neq 0$ . If you were to take large samples from  $F$  and  $G$ , what must the QQ plot look like without knowing the specific values of  $a$  and  $b$ ?

Your Answer	Score	Explanation
<input type="radio"/> The function $H(x) = F(a^2 F^{-1}(x) + b)$ for $0 \leq x \leq 1$ .		
<input type="radio"/> An identity line.		
<input checked="" type="radio"/> A line.	✓ 1.00	
<input type="radio"/> A parabola.		
Total	1.00 / 1.00	

## Question 11

Let your data be the two points  $\{1, 3\}$ . What is the bootstrap distribution of the sample mean?

Your Answer	Score	Explanation
<input type="radio"/> The distribution puts probability 1/2 on the numbers 1 and 3 both.		
<input type="radio"/> The distribution puts probabilities 1/4 and 3/4 on 1 and 3, respectively.		
<input checked="" type="radio"/> The distribution puts probabilities 1/4, 1/2 and 1/4 on the numbers 1, 2 and 3, respectively	✓ 1.00	
Total	1.00 / 1.00	

