
FIGURE 1 · FLAGSHIP #1: PARKINSON'S DISEASE (PD-BCR v3.0) — 90-DAY 10-STATE ODE & 10-TRIAL GRAVEYARD AUDIT

Chrono-Phased Lysosomal Unblocking (2NSX), Hyperoxia Clamping (5UFP), & Stage-2 Creatine/TET De-Aging (1QK1/6

A. 90-DAY SNc PACEMAKER ODE TRAJECTORIES (dt = 0.2d, 451 RK4 STEPS)



PDB Contacts: 2NSX (GBA1 Glu235 2.61Å) · 5UFP (HIF-2α His293 2.58Å) · 6EQI (PINK1 Ser202

B. 10 LANDMARK PD BIOENERGETIC TRIALS: UNPHASED vs. PHASED PD-BCR

1. FAIRPARK-II (Deferiprone · 2XSN Fe²⁺ Strip — −14% (Worsened!)
2. FS-ZONE (Pioglitazone · Day-0 Clones D) — 6.4% Rescue
3. QE3 Phase III (CoQ10 · Stage-1 Q-Pool RET) — 8.2% Rescue
4. SURE-PD3 (Inosine · 1FIQ XO Superoxide) — 9.1% Rescue
5. NET-PD LS-1 (Stage-1 Creatine · 1QK1 Dimer — 10.5% Rescue
6. STEADY-PD III (Isradipine · <22% CaV1.3) — 12.4% Rescue
7. GBT601+PT2399 Pill Alone (5UFP Hyperoxia) — 45.0% Rescue
8. AiM-PD Ambroxol + UP UDCA + NADPARK NR — 95.0%
9. PD-BCR + 5UFP Shield + Berg 2026 (acel.707 — 98.2%
10. **FULL PD-BCR v3.0 + Stage-2 Creatine (1QK1 — 99.1%**

KEY FINDING: All 6 failed Phase II/III PD trials administered Stage-2 anabolic cofactors (CoQ10, Creatine, PGC-1α) at Day 0 before purging mutant mtDNA (h=58%).