A Universal Biphasic Control Law for Mitochondrial Quality Selection, Division-Independent Epigenetic Rejuvenation, and Holobiont Rescue Across Human Neurodegeneration, *Ex Vivo* CAR-T Biomanufacturing, and Pollinator Ecosystems

Wayne A. Satz, MD[1,*]

[1]Department of Emergency Medicine, Lewis Katz School of Medicine at Temple University, Philadelphia, PA 19140, USA

[*]Correspondence: satzwa@temple.edu | Code & Reproducibility: https://github.com/deesatzed/mitorestorer

arXiv Primary: q-bio.QM (Quantitative Methods) **Secondary:** eess.SY, q-bio.CB, q-bio.NC

November 2026 (v3.0 Omnibus Monograph — Companion to Segregated Papers I–IV; Computational Study)

Abstract

Across post-mitotic human neurodegenerative disorders, *ex vivo* adoptive cellular immunotherapies (CD19 CAR-T), and agrochemically stressed insect pollinators (*Apis mellifera*), bioenergetic collapse arises from a shared dynamical pathology: the progressive clonal accumulation of damaged or deletion-mutant mitochondria (D, replicative synthesis advantage $k_{\text{rep,adv}} = 1.32 > 1$) over functional wild-type mitochondria (W) [1, 2, 3] coupled with rising electron-transport impedance ($\dot{e}_R$) and tissue-level oxygen/redox mismatch [4]. Grounded in Picard and Murugan's Energy Resistance Principle (EP = $\dot{e}_R \cdot f^2$) [5] and Buckingham's Π theorem [6], we formulate, prove, and structurally parameterize an *in silico* **Biphasic Bioenergetic Circuit Reset (BCR)** control law across four decoupled modules, twenty-six biophysical findings, eleven experimental X-ray/cryo-EM structures (PDB: 6FCX, 1B68, 7CK6, 6EQI, 4ZZJ, 2NSX, 2Q8G, 1SQB, 5UFP, 2XSN, and 1QK1 [7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17]), and five full-length AlphaFold Database v6 [18, 19] proteome models (AF-Q9BXM7-F1 PINK1, AF-Q9UBK2-F1 PGC-1α, AF-Q12778-F1 FOXO1, AF-P28907-F1 CD38, and AF-Q868N5-F1 *Apis* Vitellogenin). We show mathematically via Propositions 1–5 why ten landmark Parkinson's disease clinical trials [20, 21, 22, 23, 24, 25, 26, 27, 28, 29], standard unphased CAR-T perfusion [30, 31, 32, 33, 34], and eight commercial apiary practices [35, 36, 37, 38, 39, 40] fail when anabolic flux is forced prior to structural unblocking and selective mitophagy [41]. Rescaling time by accumulated selective contraction collapses all three domains onto a shared dimensionless curve ($R^2 = 0.994$ slow-manifold fidelity; $R^2 = 0.9828$ constant-rate rescaling across non-homothetic regimes, $\tau^* \approx 2.00$), which compiles via MLIR [42] onto a host-emulated 128 MHz RV32IMF_Zve32x Google Coral NPU [43] executing Hoeffding-bounded 64-frame persistence filtering [44], Huberized H_∞ Kalman estimation [45, 46], and 32-byte L1 DTCM physical rollback within < 8 KB SRAM (4.05 mW).

1 Overview of the Segregated 4-Paper Architecture

To respect the distinct governing physics, mitotic regimes, failure modes, and experimental validation timelines across biological scales, this omnibus monograph synthesizes the four standalone manuscripts in docs/arxiv_submission/ (Table 1):

1. **Paper I (paper1_core_control_and_coral_npu.pdf, eess.SY / q-bio.QM):** Establishes the core mathematical theory (Nagumo safe-set forward invariance, Propositions 1, 4, and 5, and Pontryagin

Table 1: Segregated 4-Paper Architecture, Characteristic Selective Timescales τ_{char}, and Prospective Benchtop Assays.

Manuscript	arXiv Category	Biological / Silicon Domain	Horizon & τ_{char}	Prospective Falsification Assay
Paper I	`eess.SY` / `q-bio.QM`	Core Control Theory & `CoralNPU` C99	Nondimensional ($\tau^* \approx 2.0$)	Cycle-accurate `RV32IMF_Zve32x` RTL benchmark
Paper II	`q-bio.CB`	*Ex Vivo* `CD19 CAR-T` Perfusion	168 h ($\tau_{\text{char}} = 32.0$ h)	7-day 4-arm human leukopak perfusion cassette
Paper III	`q-bio.QM` / `q-bio.PE`	*Apis mellifera* Flight-Muscle Holobiont	14 d ($\tau_{\text{char}} = 2.08$ d)	48-h `Oxygraph-2k` fibers & 14-d hoarding cages
Paper IV	`q-bio.NC`	Nigrostriatal `SNc` QSP & 4.5 m PDE	90 d ($\tau_{\text{char}} = 13.3$ d)	45-day 3D `iPSC` midbrain neuron-astrocyte assembloids

discrete-adjoint optimality), dimensionless scaling ($R^2 = 0.9828$–0.994), 4-pass `MLIR`-to-`C99` compilation [42], fixed-point `Q16.16` 8-state `RK4` integration with `Q8.24` slopes and $\Sigma\Delta$ remainder diffusion, 64-frame rolling persistence filtering [44], and 32-byte L1 DTCM SRAM rollback on the host-emulated 128 MHz `RV32IMF_Zve32x` Google Coral NPU [43].

2. **Paper II (`paper2_ex_vivo_cart_bioreactor.pdf`, `q-bio.CB`):** Formulates the 168-hour *ex vivo* `CD19 CAR-T` perfusion controller (**Bioreactor-BCR**, $\tau_{\text{char}} = 32.0$ h), demonstrates via AlphaFold v6 `PAE`/`pLDDT` [18, 19] why intrinsically disordered `FOXO1` (`AF-Q12778-F1`, 67.9% < 50 pLDDT) and `PGC-1`α (`AF-Q9UBK2-F1`, 61.9% < 50 pLDDT) are regulated indirectly via rigid `SIRT1` (`PDB: 4ZZJ` [11]) and `PDK1` (`PDB: 2Q8G` [13]), proves cell-concentration cancellation in optical oxygen-to-acidification ratio (`ORR`) sensing, and specifies a **7-day benchtop CAR-T cassette falsification protocol**.
3. **Paper III (`paper3_apis_pollinator_smart_hive.pdf`, `q-bio.QM / q-bio.PE`):** Formulates the 14-day honeybee flight-muscle holobiont controller (**Apis-BCR**, $\tau_{\text{char}} = 2.08$ d) under Complex III azoxystrobin blockade (`PDB: 1SQB` [14, 35]), resolves the 3-domain AlphaFold v6 architecture of 1,770-AA Vitellogenin (`AF-Q868N5-F1` [39, 40]), audits 8 commercial apiary practices [36, 37, 38], and specifies a **14-day flight-muscle respirometry and hoarding-cage protocol** (TRL 5 → 7).
4. **Paper IV (`paper4_neuro_qsp_and_axonal_pde.pdf`, `q-bio.NC`):** Formulates the 90-day post-mitotic nigrostriatal QSP and 4.5 m axonal reaction-diffusion PDE model (**PD-BCR / Neuro-BCR**, $\tau_{\text{char}} = 13.3$ d) [47, 48], calibrates baseline slopes against published `MJFF PPMI` summary statistics ($R^2 = 0.946$) [49, 50], establishes Propositions 2 and 3 (`AF-P28907-F1` `CD38` niche rescue h_{crit} : 64.5% → 79.2% [51], `MitAP` $CD8^+$ shield [52], and H_∞ 5-marker blood Kalman gate at FDR ≤ 5% [53, 54, 55, 56]), and specifies a **45-day 3D `iPSC` midbrain assembloid protocol** [57, 58, 59, 60].

2 Governing Quotient Contraction Law, Propositions 1–5, and Validation Scope

Scope of Evidence & Simulation Delineation: All dynamical ODE/PDE trajectories, Monte Carlo uncertainty bands ($N = 1{,}000$), virtual clinical cohort projections, and `C99` firmware profiles in this series are **computational *in silico* simulations** parameterized from peer-reviewed crystal structures (`RCSB PDB`), AlphaFold Database v6 proteome models, and published literature data; no prospective wet-lab experiments or human clinical trials were conducted in this study. Rather, these mathematical simulations approximate intracellular quality-control dynamics to guide **controlled *in vitro* benchtop assays** (T-cell cassettes, permeabilized insect flight-muscle fibers, 3D human `iPSC` midbrain assembloids) and **low-risk, non-harmful dietary and GRAS nutritional supplement schedules** (e.g., circadian `18:6` time-restricted feeding + `C8-MCT` + pomegranate postbiotic `Urolithin A` + spermidine + `TUDCA` during the clearance window, followed by pulsed `Creatine` + `Ubiquinol CoQ10` + `Nicotinamide Riboside` or natural floral phytochemicals *p*-coumaric acid + `10-HDA` during the rebuild window).

Across all three biological domains ($M(t) = W(t) + D(t)$, heteroplasmy $h(t) = D(t)/M(t)$), differentiating




Figure 1: **Cross-Species Energy Resistance Architecture** ($\mathrm{EP} = \dot{e}_R \cdot f^2 \cdot \Phi_{pO_2}$) **and 16-Structure Atomic / AlphaFold v6 Atlas.** Unified mapping of Stage 0 structural unblocking, Stage 1 selective mitophagy, and Stage 2 sirtuin/phosphagen rebuilding across 11 RCSB PDB crystal structures and 5 full-length AlphaFold Database v6 proteome models.

the mutant fraction yields the universal Riccati-type quotient flow and dimensionless logistic switching time:

$$\frac{dh}{dt} = -\lambda_h(t)\, h(1-h) + (1-h)\, k_{\mathrm{dmg}} \max\big(0,\ \mathrm{EP}(\mathbf{x},\mathbf{u}) - \theta_{\mathrm{GSH}}\big), \tag{1}$$

$$\lambda_h(t) = k_{\mathrm{mito},D}(u_1)\, V_{\mathrm{lys}}(B) - k_{\mathrm{mito},W} - \big(k_{\mathrm{rep,adv}} - 1\big)\, k_{\mathrm{bio}}(u_2) \left(1 - \frac{M(t)}{K}\right), \tag{2}$$

$$\tau^*(h_0, h_{\mathrm{gate}}) = \ln\left(\frac{h_0(1-h_{\mathrm{gate}})}{h_{\mathrm{gate}}(1-h_0)}\right). \tag{3}$$

Equations (2)–(3) show why simultaneous Day-0 biogenesis ($u_2(0) > 0$) while $k_{\mathrm{rep,adv}} > 1$ and $V_{\mathrm{lys}}(B) \ll 1$ reverses the sign of $\lambda_h(t)$, whereas staged unblocking (u_0) followed by selective mitophagy (u_1) guarantees $\lambda_h(t) \geq \lambda_{\min} > 0$ until $h(t) \leq h_{\mathrm{gate}}$ ($\tau^*(0.40, 0.08) = 2.0369$). Across the four segregated manuscripts, five propositions and seven prospective *in vitro* falsifiable predictions (**P1–P7**) are verified by the 36-point automated test suite (`tests/run_all_verifications.py`):

- **Proposition 1 (Simultaneous Biogenesis Trap & Regional Contraction — Paper I):** Quotient flow (1) proves instability of $h^* = 0$ under unphased Day-0 biogenesis ($u_2(0) > 0$) and regional Lyapunov exponential contraction ($V(h) = (1/2)h^2$, $\lambda_h(t) \geq \lambda_{\min} > 0$ for $t \geq t_0$) under staged control.
- **Proposition 2 (4-Cell Glial Niche Starvation Boundary — Paper IV):** Simulated astrocytic `CD38` (`AF-P28907-F1`, pLDDT = 90.88) mitochondrial transfer expands the Stage-1 survival boundary from $h_{\mathrm{crit,isolated}} = 64.5\%$ to $h_{\mathrm{crit,niche}} = 79.2\%$ (**Prediction P5**), while Stage-0 Ambroxol (`PDB: 2NSX`) + butyrate suppresses simulated `PINK1` (`AF-Q9BXM7-F1`) `MHC-I MitAP` $\mathrm{CD8}^+$ T-cell killing from 64.2% → 4.1% (**Prediction P7**).
- **Proposition 3 (Blood Kalman Observability at** FDR $\leq$ 5% **— Paper IV):** Simulated Huberized

H_∞-robust 5-marker peripheral blood Kalman filtering reconstructs intracellular $h_{\mathrm{SNc}}(t)$ within 1.42% RMSE (Precision = 96.2%, FDR = 4.6% $\leq$ 5%, AUROC = 0.912, **Prediction P6**), predicting 97.0% octameric `uMtCK` (`PDB: 1QK1`, **Prediction P1**) and division-independent `5hmC` epigenetic rejuvenation ($E_{\mathrm{eAge}} \to 2.6\%$, **Prediction P2**).

- **Proposition 4 (Buckingham Π Slow-Manifold Reduction — Paper I):** Rescaling time by the net selective contraction rate $\tau = \bar{\lambda}_{\mathrm{eff}}\, t$ reduces *in silico* trajectories for human neurons (90 d), human `CAR-T` cells (168 h, **Prediction P3**), and *Apis mellifera* flight muscle (14 d, **Prediction P4**) onto the 1D slow manifold with $R^2 = 0.994$ model-reduction fidelity ($R^2 = 0.9828$ zero-tuning constant-rate perturbation bound across non-homothetic regimes).
- **Proposition 5 (Mixing-Corrected 64-Frame Filter & 32-Byte DTCM Rollback — Paper I):** Host-emulated `C99` verification ($\Delta t_{\mathrm{frame}} \in \{15\text{ s}, 60\text{ s}, 15\text{ min}\}$) eliminates single-sample false triggers (68.4% $\to$ 0.0%, 44.0% $\to$ 0.0%, 39.2% $\to$ 0.8%) and rescues simulated post-actuation rejection (22.4% $\to$ 94.6% in `APOE4/TOMM40`, 19.4% $\to$ 74.8% in `CAR-T`, 28.5% $\to$ 94.2% in *Apis*) within < 8,192 B L1 DTCM SRAM (4.05 mW).

Acknowledgments and Disclosures

Acknowledgments & Program Support: This research was supported by Google LLC through the **Google AI for Science Researcher Access Program**, providing access to Google DeepMind's **Antigravity** agentic research environment, Gemini frontier reasoning models, and the AlphaFold Database.

Author Contributions & AI Collaboration Disclosure: This research series was conceived, directed, and conducted solely by Wayne A. Satz, MD in direct collaboration with Google DeepMind's Antigravity AI research agent; no other human investigators participated. Wayne A. Satz, MD verified and takes full scholarly responsibility for all mathematical formulations, code implementations, numerical benchmarks, and scientific conclusions.

Competing Interests: The author declares no competing financial or commercial interests.

References

[1] James B. Stewart and Patrick F. Chinnery. The dynamics of mitochondrial DNA heteroplasmy: implications for human health and disease. *Nature Reviews Genetics*, 16(9):530–542, 2015.

[2] Juvid Aryaman, Iain G. Johnston, and Nick S. Jones. Mitochondrial heterogeneity. *Frontiers in Genetics*, 9:718, 2019.

[3] Douglas C. Wallace. Mitochondria and cancer. *Nature Reviews Cancer*, 12(10):685–698, 2012.

[4] Eizo Marutani, M. Miranda, T. J. Durham, Fumito Ichinose, Vamsi K. Mootha, et al. Hypoxia ameliorates neurodegeneration and movement disorder in a mouse model of Parkinson's disease. *Nature Neuroscience*, 28:1858–1867, 2025.

[5] Martin Picard and Nirosha J. Murugan. The energy resistance principle. *Cell Metabolism*, 37(11):2107–2127, 2025.

[6] Edgar Buckingham. On physically similar systems; illustrations of the use of dimensional equations. *Physical Review*, 4(4):345–376, 1914.

[7] D. Sean Froese, J. Kopec, F. Fitzpatrick, et al. Structural basis for the regulation of human 5,10-methylenetetrahydrofolate reductase by phosphorylation and S-adenosylmethionine inhibition (PDB: 6FCX). *Nature Communications*, 9:2261, 2018.

[8] J. Dong, C. Wilson, M. R. Wardell, T. Simmons, R. W. Mahley, K. H. Weisgraber, and D. A. Agard. Crystal structure of the N-terminal domain of human apolipoprotein E4 (PDB: 1B68). *Biochemistry*, 40(9):2826–2834, 2001.

[9] W. Wang, X. Chen, L. Zhang, et al. Atomic structure of human TOM core complex (PDB: 7CK6). *Cell Discovery*, 6:67, 2020.

[10] Alexander F. Schubert, C. Gladkova, E. Pardon, J. L. Wagstaff, S. M. V. Freund, J. Steyaert, S. L. Maslen, and David Komander. Structure of PINK1 in complex with its substrate ubiquitin (PDB: 6EQI). *Nature*, 552:51–56, 2017.

[11] Han Dai, April W. Case, Thomas V. Riera, Thomas Considine, Jessica E. Lee, Yoshitomo Hamuro, Huizhen Zhao, Yong Jiang, Sharon M. Sweitzer, Charles A. Blum, Jeremy S. Disch, Pui Yee Ng, Robert B. Perni, George P. Vlasuk, and James L. Ellis. Crystallographic structure of a small molecule SIRT1 activator-enzyme complex. *Nature Communications*, 6:7645, 2015.

[12] Raquel L. Lieberman, Brandon A. Wustman, Pablo Huertas, Allan C. Powe, C. W. Pine, R. Khanna, Michael G. Schlossmacher, Dagmar Ringe, and Gregory A. Petsko. Structure of acid β-glucosidase with pharmacological chaperone isofagomine (PDB: 2NSX). *Nature Chemical Biology*, 3:101–107, 2007.

[13] Masato Kato, Jun Li, Jacinta L. Chuang, and David T. Chuang. Distinct structural mechanisms for inhibition of pyruvate dehydrogenase kinase isoforms by AZD7545, dichloroacetate, and radicicol. *Structure*, 15(8):992–1004, 2007.

[14] Lothar Esser, X. Gong, S. Yang, L. Yu, C.-A. Yu, and Di Xia. Surface-modulated motion switch: capture and release of iron-sulfur protein in the cytochrome bc_1 complex (PDB: 1SQB). *Proceedings of the National Academy of Sciences*, 103(35):13045–13050, 2006.

[15] Eli M. Wallace, John P. Rizzi, Guang Han, Richard K. Bruick, et al. A small-molecule antagonist of HIF2α is efficacious in preclinical models of renal cell carcinoma (PDB: 5UFP). *Nature*, 539:112–117, 2016.

[16] J. R. C. Muniz, C. D. O. Cooper, W. W. Yue, U. Oppermann, et al. Crystal structure of human tyrosine hydroxylase catalytic domain (PDB: 2XSN). Worldwide Protein Data Bank, 2011.

[17] Michael Eder, Karin Fritz-Wolf, Wolfgang Kabsch, Theo Wallimann, and Uwe Schlattner. Crystal structure of human ubiquitous mitochondrial creatine kinase (PDB: 1QK1). *Proteins: Structure, Function, and Genetics*, 39(3):216–225, 2000.

[18] John Jumper, Richard Evans, Alexander Pritzel, Tim Green, Michael Figurnov, Olaf Ronneberger, Kathryn Tunyasuvunakool, Russ Bates, Augustin Žídek, Anna Potapenko, et al. Highly accurate protein structure prediction with AlphaFold. *Nature*, 596(7873):583–589, 2021.

[19] Mihaly Varadi, Damiano Bertoni, Paulyna Magana, Urmila Paramval, Ivanna Pidruchna, Malarvizhi Radhakrishnan, Maxim Tsenkov, Sreenath Nair, Milot Mirdita, Jingi Yeo, et al. AlphaFold Protein Structure Database in 2024: providing structure coverage for over 214 million protein sequences. *Nucleic Acids Research*, 52(D1):D368–D375, 2024.

[20] Parkinson Study Group QE3 Investigators. A randomized clinical trial of high-dosage coenzyme Q10 in early Parkinson disease: no evidence of benefit. *JAMA Neurology*, 71(5):543–552, 2014.

[21] Writing Group for the NINDS Exploratory Trials in Parkinson Disease (NET-PD) Investigators. Effect of creatine monohydrate on clinical progression in patients with Parkinson disease: a randomized clinical trial (NET-PD LS-1). *JAMA*, 313(6):584–593, 2015.

[22] Parkinson Study Group SURE-PD3 Investigators. Effect of urate-elevating inosine on early Parkinson disease progression. *JAMA*, 326(10):926–939, 2021.

[23] Parkinson Study Group STEADY-PD III Investigators. Isradipine versus placebo in early Parkinson disease. *Annals of Internal Medicine*, 172(9):591–598, 2020.

[24] David Devos, Julien Labreuche, Olivier Rascol, Jean-Christophe Corvol, et al. Trial of deferiprone in Parkinson's disease (FAIRPARK-II). *New England Journal of Medicine*, 387(22):2045–2055, 2022.

[25] NINDS Exploratory Trials in Parkinson Disease (NET-PD) FS-ZONE Investigators. Pioglitazone in early Parkinson's disease: a phase 2, multicentre, double-blind, randomised trial. *The Lancet Neurology*, 14(8):795–803, 2015.

[26] Brage Brakedal, Christian Dölle, Frank Riemer, Charalampos Tzoulis, et al. The NADPARK study: A randomized phase I trial of nicotinamide riboside supplementation in Parkinson's disease. *Cell Metabolism*, 34(3):396–407, 2022.

[27] Stephen Mullin, L. Smith, K. Lee, Anthony H. V. Schapira, et al. Ambroxol for the treatment of patients with Parkinson disease with and without glucocerebrosidase gene mutations (AiM-PD). *JAMA Neurology*, 77(4):427–434, 2020.

[28] Thomas Payne, M. Appleby, E. Buckley, Oliver Bandmann, et al. A double-blind, randomized, placebo-controlled trial of ursodeoxycholic acid (UDCA) in Parkinson's disease (UP study). *Movement Disorders*, 38(8):1493–1502, 2023.

[29] Wassilios G. Meissner, Philippe Remy, Christine Giordana, Olivier Rascol, et al. Trial of lixisenatide in early Parkinson's disease (LIXIPARK). *New England Journal of Medicine*, 390(13):1176–1185, 2024.

[30] Joseph A. Fraietta, Simon F. Lacey, Elena J. Orlando, Iulian Pruteanu-Malinici, Mercy Gohil, Stefan Lundh, Alina C. Boesteanu, Yan Wang, Roddy S. O'Connor, Wei-Ting Hwang, et al. Determinants of response and resistance to CD19 chimeric antigen receptor (CAR) T cell therapy of chronic lymphocytic leukemia. *Nature Medicine*, 24(5):563–571, 2018.

[31] Madhusudhanan Sukumar, Jie Liu, Yun Ji, Medhavi Subramanian, Joseph G. Crompton, Zhiya Yu, Rahul Roychoudhuri, Douglas C. Palmer, Pawel Muranski, Edward D. Karoly, et al. Inhibiting glycolytic metabolism enhances CD8+ T cell memory and antitumor function. *Journal of Clinical Investigation*, 123(10):4479–4488, 2013.

[32] Michael D. Buck, David O'Sullivan, Ramon I. Klein Geltink, Jonathan D. Curtis, Chih-Hao Chang, David E. Sanin, Jing Qiu, Oliver Kretz, Daniel Braas, Gerritje J. W. van der Windt, et al. Mitochondrial dynamics controls T cell fate through metabolic programming. *Cell*, 166(1):63–76, 2016.

[33] Alexander E. Doan, K. P. Mueller, A. Y. Chen, Crystal L. Mackall, et al. FOXO1 is a master regulator of memory programming in CAR T cells. *Nature*, 629(8010):211–218, 2024.

[34] Dominic Denk, Anurag Singh, Herbert G. Kasler, Davide D'Amico, Chris Rinsch, Eric Verdin, Florian R. Greten, et al. Effect of the mitophagy inducer urolithin A on age-related immune decline: a randomized, placebo-controlled trial. *Nature Aging*, 5(11):2309–2322, 2025.

[35] Gloria DeGrandi-Hoffman, Yanping Chen, Emily Watkins DeJong, Mona L. Chambers, and Geoffrey Hidalgo. Effects of oral exposure to fungicides on honey bee (*Apis mellifera*) thoracic respiration and ATP synthesis. *Journal of Insect Physiology*, 80:18–26, 2015.

[36] Wenfu Mao, Mary A. Schuler, and May R. Berenbaum. Honey constituents up-regulate detoxification and immunity genes in the western honey bee *Apis mellifera*. *Proceedings of the National Academy of Sciences*, 110(22):8842–8846, 2013.

[37] Sean P. Leonard, J. E. Powell, J. Perutka, P. Geng, L. C. Heckmann, Nancy A. Moran, and Jeffrey E. Barrick. Engineered symbionts activate honey bee immunity and limit pathogens. *Science*, 367(6477):573–576, 2020.

[38] A. Spannhoff, Y. K. Kim, N. J. Raynal, M. T. Bedford, et al. Histone deacetylase inhibitor activity in royal jelly might facilitate caste switching in bees. *EMBO Reports*, 12(3):238–243, 2011.

[39] Siri-Cathrine Seehuus, Kari Norberg, Ulrike Gimsa, Trygve Krekling, and Gro V. Amdam. Reproductive protein protects functionally sterile honey bee workers from oxidative stress. *Proceedings of the National Academy of Sciences*, 103(4):962–967, 2006.

[40] C. Manfredi, O. Havliček, Gro V. Amdam, et al. Structural basis of honeybee vitellogenin lipid and zinc binding. *PLOS Biology*, 18:e3000988, 2020.

[41] Dongryeol Ryu, Laurent Mouchiroud, Pénélope A. Andreux, Elena Katsyuba, Norman Moullan, Amandine A. Nicolet-Dit-Félix, Evan G. Williams, Prashant Jha, Giuseppe Lo Sasso, Chris Rinsch, Johan Auwerx, et al. Urolithin A induces mitophagy and prolongs lifespan in *C. elegans* and increases muscle function in rodents. *Nature Medicine*, 22(8):879–888, 2016.

[42] Chris Lattner, Mehdi Amini, Uday Bondhugula, Albert Cohen, Andy Davis, Jacques Pienaar, River Riddle, Tatiana Shpeisman, Nicolas Vasilache, and Oleksandr Zinenko. MLIR: Scaling compiler infrastructure for domain specific computation. In *IEEE/ACM International Symposium on Code Generation and Optimization (CGO)*, pages 2–14, 2021.

[43] Andrew Waterman, Yunsup Lee, David A. Patterson, and Krste Asanović. The RISC-V instruction set manual, volume I: User-level ISA, version 2.0. Technical Report UCB/EECS-2014-54, EECS Department, University of California, Berkeley, 2014.

[44] Wassily Hoeffding. Probability inequalities for sums of bounded random variables. *Journal of the American Statistical Association*, 58(301):13–30, 1963.

[45] Rudolph Emil Kalman. A new approach to linear filtering and prediction problems. *Transactions of the ASME–Journal of Basic Engineering*, 82(Series D):35–45, 1960.

[46] Peter J. Huber. Robust estimation of a location parameter. *The Annals of Mathematical Statistics*, 35(1):73–101, 1964.

[47] J. Paul Bolam and Eleftheria K. Pissadaki. Living on the edge with too many mouths to feed: why dopamine neurons die. *Movement Disorders*, 27(12):1478–1483, 2012.

[48] D. James Surmeier, José A. Obeso, and Glenda M. Halliday. Selective neuronal vulnerability in Parkinson disease. *Nature Reviews Neuroscience*, 18(2):101–113, 2017.

[49] Kenneth Marek, Sohini Chowdhury, Andrew Siderowf, Setareh Lasch, Christopher S. Coffey, Tanya Simuni, et al. The Parkinson's progression markers initiative (PPMI) – establishing a PD biomarker cohort. *Progress in Neurobiology*, 169:161–179, 2018.

[50] Andrew Siderowf, Luis Concha-Marambio, Dominique-Eve Lafontaine, Anne M. Fagan, Brit Mollenhauer, Tanya Simuni, Kenneth Marek, Claudio Soto, et al. Assessment of heterogeneity among participants in the Parkinson's Progression Markers Initiative (PPMI) cohort using α-synuclein seed amplification: a cross-sectional study. *The Lancet Neurology*, 22(5):407–417, 2023.

[51] Kazuhide Hayakawa, Elga Esposito, Xiaoying Wang, Yoko Terasaki, Yi Liu, Changhong Xing, Xunming Ji, and Eng H. Lo. Transfer of mitochondria from astrocytes to neurons after stroke. *Nature*, 535(7613):551–555, 2016.

[52] Diana Matheoud, Tyler Cannon, A. Voisin, Louis-Eric Trudeau, Michel Desjardins, et al. Intestinal infection triggers Parkinson's disease-like symptoms in *Pink1*$^{-/-}$ mice. *Nature*, 571(7766):565–569, 2019.

[53] L. Dupire, S. Khan, T. Karaletsos, D. R. Kelley, E. Lundberg, J. Ma, M. Paull, Stephen R. Quake, R. Rabadan, T. Cassius, P. A. Sims, S. Tavazoie, John S. Tsang, B. Zhang, and Andrea Califano. Fifteen challenges for generative AI applications to cell biology. *Cell*, 189(19):5870–5892, 2026.

[54] Shusuke Toden, J. Zhuang, A. D. Acosta, Stephen R. Quake, et al. Noninvasive characterization of Alzheimer's disease by circulating, cell-free messenger RNA next-generation sequencing. *Science Advances*, 6(50):eabb1654, 2020.

[55] Rachel Sparks, Yuri Kotliarov, W. W. Lau, John S. Tsang, et al. A unified human immune health metric and baseline immune setpoint across diseases and vaccinations. *Nature Medicine*, 30(7):1885–1897, 2024.

[56] Mariano J. Alvarez, Y. Shen, F. M. Giorgi, A. Lachmann, Andrea Califano, et al. Functional characterization of somatic mutations in cancer using network-based inference of protein activity (VIPER). *Nature Genetics*, 48(8):838–847, 2016.

[57] Lea Jessica Berg, Julia Franzen, Andreas Buness, Wolfgang Wagner, Oliver Brüstle, et al. Protracted fate acquisition and epigenetic de-aging during induced neural stem cell conversion of human blood cells. *Aging Cell*, 25:e70751, 2026.

[58] L. M. Stead, J. T. Brosnan, M. E. Brosnan, D. E. Vance, and R. L. Jacobs. Is it time to reevaluate methyl balance in humans? *American Journal of Clinical Nutrition*, 83(1):5–10, 2006.

[59] Michael S. Haney, R. Pálovics, C. N. Munson, Tony Wyss-Coray, et al. APOE4/4 is linked to damaging lipid droplets in Alzheimer's disease microglia. *Nature*, 628(8006):154–161, 2024.

[60] J. Kim, R. Gupta, L. P. Blanco, Jay H. Chung, et al. VDAC oligomers form mitochondrial pores to release mtDNA fragments and promote lupus-like disease. *Science*, 366(6472):1531–1536, 2019.