

# Fundamentals of Optimization

## Exercise 2

### Remarks

- All questions that are available in the STACK quiz are duly marked. Please solve those using STACK.
- We have added marks for each question. Please note that those are purely for illustrative purposes. The exercise set will not be marked.

## STACK Problems

### 1 Basic Solutions and Basic Feasible Solutions (3 marks)

#### STACK question

Consider the following polyhedron

$$\mathcal{P} = \{x \in \mathbb{R}^3 : -4x_1 + x_2 - x_3 \leq -4, -2x_1 + x_2 - x_3 \leq -2, -4x_2 + x_3 \leq 4, 4x_2 + x_3 = 0, x_3 \leq 5, x_3 \geq 0\}.$$

Decide, for each of the points  $\hat{x}$  given below, whether  $\hat{x}$  is *infeasible and not a basic solution*, *feasible but not a basic feasible solution*, *a basic solution but infeasible*, or *a basic feasible solution*.

$$(1.1) \quad \hat{x} = [0, 0, 0]^T.$$

$$(1.2) \quad \hat{x} = [1, 0, 0]^T.$$

$$(1.3) \quad \hat{x} = [1, -4/3, -4/3]^T.$$

$$(1.4) \quad \hat{x} = [3/8, -1/2, 2]^T.$$

$$(1.5) \quad \hat{x} = [11/16, -1/4, 1]^T.$$

### 2 Graphical Method (2 marks)

#### STACK question

Consider the following polyhedron:

$$\mathcal{P} = \{[x_1, x_2]^T \in \mathbb{R}^2 : -x_1 + x_2 \leq 2, x_1 - 2x_2 \geq -6, x_1 + 2x_2 \geq 1, x_2 \geq 0\}.$$

Using the graphical method, determine, for each of the following objective functions, the optimal value denoted by  $z^*$  (use **+inf** for  $+\infty$  and **-inf** for  $-\infty$ ), and whether the set of optimal solutions, denoted by  $\mathcal{P}^*$ , is either *empty*, a *single vertex*, a *line segment*, a *half line*, or  $\mathcal{P}^* = \mathcal{P}$ .

$$(2.1) \quad \min\{3x_1 : x \in \mathcal{P}\}.$$

$$(2.2) \quad \max\{-2x_1 - 4x_2 : x \in \mathcal{P}\}.$$

$$(2.3) \quad \min\{0 : x \in \mathcal{P}\}.$$

$$(2.4) \quad \min\{2x_1 - 4x_2 : x \in \mathcal{P}\}.$$

## Open Ended Problems

### 3 Polyhedra in Standard Form (1 mark)

(3.1) Convert the following general linear programming problem into standard form:

$$\begin{array}{llllllll} \max & -x_1 & + & 3x_2 & - & 5x_3 & + & x_4 \\ \text{s.t.} & & & & & & & \\ & -x_1 & + & 3x_2 & - & 2x_3 & - & 3x_4 & \geq & 6 \\ & 3x_1 & - & x_2 & + & 5x_3 & - & 2x_4 & = & -3 \\ & -x_1 & - & 3x_2 & + & 2x_3 & - & x_4 & \leq & 2 \\ & x_1 \leq 0 & , & x_2 \geq 0 & , & & & x_4 \geq 0. \end{array}$$

*Remark* It is irrelevant whether the problem is actually feasible or not.

### 4 Polytopes vs Polyhedra (2 marks)

(4.1) Let  $\mathcal{P} \subseteq \mathbb{R}^n$  be a nonempty polyhedron. Prove the following result:

$\mathcal{P} \subseteq \mathbb{R}^n$  is a polytope if and only if, for every  $c \in \mathbb{R}^n$ , each of the two linear programming problems given by

$$(P1) \quad \min\{c^T x : x \in \mathcal{P}\} \quad \text{and} \quad (P2) \quad \max\{c^T x : x \in \mathcal{P}\}$$

has a finite optimal value.

### 5 Existence of Vertices in Polyhedra (2 marks)

(5.1) Either prove the following result or give a counterexample:

Let  $\mathcal{P} \subseteq \mathbb{R}^n$  be a nonempty polyhedron. If  $\mathcal{P}$  has no vertices, then there exists a vector  $c \in \mathbb{R}^n$  such that each of the two linear programming problems given by

$$(P1) \quad \min\{c^T x : x \in \mathcal{P}\} \quad \text{and} \quad (P2) \quad \max\{c^T x : x \in \mathcal{P}\}$$

is unbounded.

(5.2) Either prove the following result or give a counterexample:

Let  $\mathcal{P} \subseteq \mathbb{R}^n$  be a nonempty polyhedron. If there exists a vector  $c \in \mathbb{R}^n$  such that each of the two linear programming problems given by

$$(P1) \quad \min\{c^T x : x \in \mathcal{P}\} \quad \text{and} \quad (P2) \quad \max\{c^T x : x \in \mathcal{P}\}$$

is unbounded, then  $\mathcal{P}$  has no vertices.