

Side channel protections for CSIDH

Luca De Feo

IBM Research Zürich

October 16, 2019, PHISIC, Gardanne

based on joint work with

D. Cervantes-Vázquez, M. Chenu, J.J. Chi-Domínguez, F. Rodríguez-Henríquez, B. Smith

Slides online at <https://defeo.lu/docet>



Why isogenies?


Six families still in NIST post-quantum competition:

Lattices	9 encryption	3 signature
Codes	7 encryption	
Multivariate		4 signature
Isogenies	1 encryption	
Hash-based		1 signature
MPC		1 signature

Why isogenies?

Six families still in NIST post-quantum competition:


Lattices	9 encryption	3 signature
Codes	7 encryption	
Multivariate		4 signature
Isogenies	1 encryption	1 signature
Hash-based		1 signature
MPC		



Why isogenies?

Six families still in NIST post-quantum competition:

Lattices	9 encryption	3 signature
Codes	7 encryption	
Multivariate		4 signature
Isogenies	1 encryption	
Hash-based		1 signature
MPC		1 signature



Iso-what?!

Keywords

- An **isogeny** is a **map** between two **elliptic curves**;

Iso-what?!

Keywords

- An **isogeny** is a **map** between two **elliptic curves**;
- It is a **group morphism**:

$$\phi(P + Q) = \phi(P + Q);$$

Iso-what?!

Keywords

- An **isogeny** is a **map** between two **elliptic curves**;
- It is a **group morphism**:

$$\phi(P + Q) = \phi(P) + \phi(Q);$$

- It is an **algebraic map**:

$$\phi(x, y) = \left(\frac{g(x)}{h(x)}, y \left(\frac{g(x)}{h(x)} \right)' \right);$$

Iso-what?!

Keywords

- An **isogeny** is a **map** between two **elliptic curves**;
- It is a **group morphism**:

$$\phi(P + Q) = \phi(P) + \phi(Q);$$

- It is an **algebraic map**:

$$\phi(x, y) = \left(\frac{g(x)}{h(x)}, y \left(\frac{g(x)}{h(x)} \right)' \right);$$

- It is entirely determined by its **kernel** (i.e., by a single **point**);

Iso-what?!

Keywords

- An **isogeny** is a **map** between two **elliptic curves**;
- It is a **group morphism**:

$$\phi(P + Q) = \phi(P) + \phi(Q);$$


- It is an **algebraic map**:

$$\phi(x, y) = \left(\frac{g(x)}{h(x)}, y \left(\frac{g(x)}{h(x)} \right)' \right);$$


- It is entirely determined by its **kernel** (i.e., by a single **point**);
- Isogeny **degree** = size of the kernel = order of kernel generator \approx size of the polynomials;

Isogenies: an example over \mathbb{F}_{11}

$$E : y^2 = x^3 + x$$



$$E' : y^2 = x^3 - 4x$$




$$\phi(x, y) = \left(\frac{x^2 + 1}{x}, \quad y \frac{x^2 - 1}{x^2} \right)$$

Isogenies: an example over \mathbb{F}_{11}

$$E : y^2 = x^3 + x$$


$$E' : y^2 = x^3 - 4x$$




$$\phi(x, y) = \left(\frac{x^2 + 1}{x}, \quad y \frac{x^2 - 1}{x^2} \right)$$

- Kernel generator in red.
- This is a degree 2 map.
- Analogous to $x \mapsto x^2$ in \mathbb{F}_q^* .


Isogeny graphs




Isogeny graphs




Isogeny graphs




Isogeny graphs




Isogeny graphs




Isogeny graphs




Isogeny graphs




Isogeny graphs




Isogeny graphs




Isogeny graphs




Isogeny graphs




Isogeny graphs




Isogeny graphs




Isogeny graphs




Isogeny graphs




Isogeny graphs




Isogeny graphs



Isogeny graphs



Isogeny graphs




Isogeny graphs



Isogeny graphs

$$j = \overset{\bullet}{1728}$$

Isogeny graphs





Isogeny graphs



The beauty and the beast

(credit: Lorenz Panny)

Components of particular isogeny graphs look like this:





Which of these is good for crypto?

The beauty and the beast

(credit: Lorenz Panny)

Components of particular isogeny graphs look like this:




Which of these is good for crypto? Both.

The beauty and the beast

(credit: Lorenz Panny)


At this time, there are two distinct families of systems:



\mathbb{F}_p

CSIDH [pron.: sea-side]

<https://csidh.isogeny.org>



\mathbb{F}_{p^2}

SIDH

<https://sike.org>


CSIDH vs SIDH

	CSIDH	SIDH
Speed (on x64 arch., NIST 1)	~ 70ms	~ 6ms
Public key size (NIST 1)	64B	346B
Key compression		
↳ speed		~ 11ms
↳ size		209B
Submitted to NIST	no	yes
TRL	4	6
Best classical attack	$p^{1/4}$	$p^{1/4} (p^{3/8})$
Best quantum attack	$\tilde{O}\left(3^{\sqrt{\log_3 p}}\right)$	$p^{1/6} (p^{3/8})$
Key size scales	quadratically	linearly
CPA security	yes	yes
CCA security	yes	Fujisaki-Okamoto
Constant time	it's complicated	yes
Non-interactive key exchange	yes	no
Signatures	short but (slow do not scale)	big and slow

CSIDH vs SIDH


	CSIDH	SIDH
Speed (on x64 arch., NIST 1)	~ 70ms	~ 6ms
Public key size (NIST 1)	64B	346B
Key compression		
↳ speed		~ 11ms
↳ size		209B
Submitted to NIST	no	yes
TRL	4	6
Best classical attack	$p^{1/4}$	$p^{1/4} (p^{3/8})$
Best quantum attack	$\tilde{O}\left(3^{\sqrt{\log_3 p}}\right)$	$p^{1/6} (p^{3/8})$
Key size scales	quadratically	linearly
CPA security	yes	yes
CCA security	yes	Fujisaki-Okamoto
Constant time	it's complicated	yes
Non-interactive key exchange	yes	no
Signatures	short but (slow do not scale)	big and slow

The CSIDH graph



Vertices are
supersingular elliptic curves
over \mathbb{F}_p .

The CSIDH graph




Vertices are
supersingular elliptic curves
over \mathbb{F}_p .

Edges are isogenies
of bounded prime degree.

— degree 3

The CSIDH graph




Vertices are
supersingular elliptic curves
over \mathbb{F}_p .

Edges are isogenies
of bounded prime degree.

— degree 3

— degree 5

The CSIDH graph



Vertices are
supersingular elliptic curves
over \mathbb{F}_p .


Edges are isogenies
of bounded prime degree.

— degree 3

— degree 5

— degree 7


CSIDH key exchange



Public parameters:

- A supersingular curve E_0/\mathbb{F}_p ;
- A set of small prime degree isogenies.


CSIDH key exchange



Public parameters:

- A supersingular curve E_0/\mathbb{F}_p ;
- A set of small prime degree isogenies.
- ① **Alice** takes a **secret** random walk $\phi_A : E_0 \rightarrow E_A$ of length $O(\log p)$;


CSIDH key exchange



Public parameters:

- A supersingular curve E_0/\mathbb{F}_p ;
 - A set of small prime degree isogenies.
- 1 **Alice** takes a **secret** random walk $\phi_A : E_0 \rightarrow E_A$ of length $O(\log p)$;
 - 2 **Bob** does the same;


CSIDH key exchange



Public parameters:

- A supersingular curve E_0/\mathbb{F}_p ;
 - A set of small prime degree isogenies.
- 1 **Alice** takes a **secret** random walk $\phi_A : E_0 \rightarrow E_A$ of length $O(\log p)$;
 - 2 **Bob** does the same;
 - 3 They publish E_A and E_B ;


CSIDH key exchange



Public parameters:

- A supersingular curve E_0/\mathbb{F}_p ;
 - A set of small prime degree isogenies.
- 1 **Alice** takes a **secret** random walk $\phi_A : E_0 \rightarrow E_A$ of length $O(\log p)$;
 - 2 **Bob** does the same;
 - 3 They publish E_A and E_B ;
 - 4 **Alice** repeats her secret walk ϕ_A starting from E_B .

CSIDH key exchange



Public parameters:

- A supersingular curve E_0/\mathbb{F}_p ;
 - A set of small prime degree isogenies.
- 1 **Alice** takes a **secret** random walk $\phi_A : E_0 \rightarrow E_A$ of length $O(\log p)$;
 - 2 **Bob** does the same;
 - 3 They publish E_A and E_B ;
 - 4 **Alice** repeats her secret walk ϕ_A starting from E_B .
 - 5 **Bob** repeats his secret walk ϕ_B starting from E_A .

CSIDH data flow

Your secret: a vector of number of **isogeny steps** for each degree

$(5, 1, -4, \dots)$




Your public key: (the j -invariant of) a supersingular elliptic curve

$j =$ 0x23baf75419531a44f3b97cc9d8291a275047fcdae0c9a0c0ebb993964f821f2
0c11058a4200ff38c4a85e208345300033b0d3119ff4a7c1be0acd62a622002a9

Isogeny evaluation

Repeat:


- Take a **random point** $P \in E(\mathbb{F}_p)$;
- Set $Q = [c]P$,
where c is an **appropriate cofactor**, so
that $N = \#\langle Q \rangle$ contains only **useful**
prime factors;
- Advance by the **degree N** isogeny of
kernel $\langle Q \rangle$.




Isogeny evaluation

Repeat:

- Take a **random point** $P \in E(\mathbb{F}_p)$;
- Set $Q = [c]P$,
where c is an **appropriate cofactor**, so
that $N = \#\langle Q \rangle$ contains only **useful**
prime factors;
- Advance by the **degree N** isogeny of
kernel $\langle Q \rangle$.




$$\#\langle P \rangle = 3 \cdot 5 \cdot 7 \cdot 11$$



Isogeny evaluation


Repeat:

- Take a **random point** $P \in E(\mathbb{F}_p)$;
- Set $Q = [c]P$,
where c is an **appropriate cofactor**, so
that $N = \#\langle Q \rangle$ contains only **useful**
prime factors;
- Advance by the **degree N** isogeny of
kernel $\langle Q \rangle$.



$$\#\langle P \rangle = 3 \cdot 5 \cdot 7 \cdot 11$$



$$Q = P$$



Isogeny evaluation

Repeat:


- Take a **random point** $P \in E(\mathbb{F}_p)$;
- Set $Q = [c]P$,
where c is an **appropriate cofactor**, so
that $N = \#\langle Q \rangle$ contains only **useful**
prime factors;
- Advance by the **degree N** isogeny of
kernel $\langle Q \rangle$.



Isogeny evaluation


Repeat:

- Take a **random point** $P \in E(\mathbb{F}_p)$;
- Set $Q = [c]P$,
where c is an **appropriate cofactor**, so
that $N = \#\langle Q \rangle$ contains only **useful**
prime factors;
- Advance by the **degree N** isogeny of
kernel $\langle Q \rangle$.



$$\#\langle P \rangle = 3 \cdot 5 \cdot 7 \cdot 11$$


$$Q = [11]P$$



Isogeny evaluation


Repeat:

- Take a **random point** $P \in E(\mathbb{F}_p)$;
- Set $Q = [c]P$,
where c is an **appropriate cofactor**, so
that $N = \#\langle Q \rangle$ contains only **useful**
prime factors;
- Advance by the **degree N** isogeny of
kernel $\langle Q \rangle$.



$$\#\langle P \rangle = 3 \cdot 5 \cdot 7 \cdot 11$$


$$Q = [11]P$$



Isogeny evaluation

Repeat:


- Take a **random point** $P \in E(\mathbb{F}_p)$;
- Set $Q = [c]P$,
where c is an **appropriate cofactor**, so
that $N = \#\langle Q \rangle$ contains only **useful**
prime factors;
- Advance by the **degree N** isogeny of
kernel $\langle Q \rangle$.



Isogeny evaluation


Repeat:

- Take a **random point** $P \in E(\mathbb{F}_p)$;
- Set $Q = [c]P$,
where c is an **appropriate cofactor**, so
that $N = \#\langle Q \rangle$ contains only **useful**
prime factors;
- Advance by the **degree N** isogeny of
kernel $\langle Q \rangle$.



$$\#\langle P \rangle = 3 \cdot 7 \cdot 11$$


$$Q = [3 \cdot 11]P$$



Isogeny evaluation


Repeat:

- Take a **random point** $P \in E(\mathbb{F}_p)$;
- Set $Q = [c]P$,
where c is an **appropriate cofactor**, so
that $N = \#\langle Q \rangle$ contains only **useful**
prime factors;
- Advance by the **degree N isogeny** of
kernel $\langle Q \rangle$.



$$\#\langle P \rangle = 3 \cdot 7 \cdot 11$$


$$Q = [3 \cdot 11]P$$



Isogeny evaluation

Repeat:


- Take a **random point** $P \in E(\mathbb{F}_p)$;
- Set $Q = [c]P$,
where c is an **appropriate cofactor**, so
that $N = \#\langle Q \rangle$ contains only **useful**
prime factors;
- Advance by the **degree N** isogeny of
kernel $\langle Q \rangle$.



Isogeny evaluation


Repeat:

- Take a **random point** $P \in E(\mathbb{F}_p)$;
- Set $Q = [c]P$,
where c is an **appropriate cofactor**, so
that $N = \#\langle Q \rangle$ contains only **useful**
prime factors;
- Advance by the **degree N isogeny** of
kernel $\langle Q \rangle$.



$$\#\langle P \rangle = 3 \cdot 5 \cdot 7 \cdot 11$$


$$Q = [3 \cdot 7 \cdot 11]P$$



Isogeny evaluation


Repeat:

- Take a **random point** $P \in E(\mathbb{F}_p)$;
- Set $Q = [c]P$,
where c is an **appropriate cofactor**, so
that $N = \#\langle Q \rangle$ contains only **useful**
prime factors;
- Advance by the **degree N** isogeny of
kernel $\langle Q \rangle$.



$$\#\langle P \rangle = 3 \cdot 5 \cdot 7 \cdot 11$$



$$Q = [3 \cdot 7 \cdot 11]P$$



Isogeny evaluation

Repeat:


- Take a **random point** $P \in E(\mathbb{F}_p)$;
- Set $Q = [c]P$,
where c is an **appropriate cofactor**, so
that $N = \#\langle Q \rangle$ contains only **useful**
prime factors;
- Advance by the **degree N** isogeny of
kernel $\langle Q \rangle$.



Isogeny evaluation


Repeat:

- Take a **random point** $P \in E(\mathbb{F}_p)$;
- Set $Q = [c]P$,
where c is an **appropriate cofactor**, so
that $N = \#\langle Q \rangle$ contains only **useful**
prime factors;
- Advance by the **degree N isogeny** of
kernel $\langle Q \rangle$.



$$\#\langle P \rangle = 3 \cdot 5 \cdot 7 \cdot 11$$



$$Q = [3 \cdot 7 \cdot 11]P$$



Isogeny evaluation

Repeat:


- Take a **random point** $P \in E(\mathbb{F}_p)$;
- Set $Q = [c]P$,
where c is an **appropriate cofactor**, so
that $N = \#\langle Q \rangle$ contains only **useful**
prime factors;
- Advance by the **degree N** isogeny of
kernel $\langle Q \rangle$.



Isogeny evaluation


Repeat:

- Take a **random point** $P \in E(\mathbb{F}_p)$;
- Set $Q = [c]P$,
where c is an **appropriate cofactor**, so
that $N = \#\langle Q \rangle$ contains only **useful**
prime factors;
- Advance by the **degree N** isogeny of
kernel $\langle Q \rangle$.



$$\#\langle P \rangle = 3 \cdot 5 \cdot 7 \cdot 11$$


$$Q = [3 \cdot 7 \cdot 11]P$$



Isogeny evaluation

Repeat:


- Take a **random point** $P \in E(\mathbb{F}_p)$;
- Set $Q = [c]P$,
where c is an **appropriate cofactor**, so
that $N = \#\langle Q \rangle$ contains only **useful**
prime factors;
- Advance by the **degree N** isogeny of
kernel $\langle Q \rangle$.



Isogeny evaluation

Repeat:

- Take a **random point** $P \in E(\mathbb{F}_p)$;
- Set $Q = [c]P$,
where c is an **appropriate cofactor**, so
that $N = \#\langle Q \rangle$ contains only **useful**
prime factors;
- Advance by the **degree N** isogeny of
kernel $\langle Q \rangle$.



In seek of constant time

Two obstacles for constant time:

- 1 Some random points P may lack some factors;
- 2 Number of isogeny evaluations dependent on secret key.

In seek of constant time

Two obstacles for constant time:

- ① ~~Some random points P may lack some factors;~~ Unrelated to secret key if truly random.
- ② Number of isogeny evaluations dependent on secret key.

In seek of constant time

Two obstacles for constant time:

- 1 Some random points P may lack some factors; Unrelated to secret key if truly random.
- 2 Number of isogeny evaluations dependent on secret key.

Meyer, Campos, Reith 2018; Onuki, Aikawa, Yamazaki, Takagi 2019

- “Dummy” isogenies:
 - ▶ Always do exactly the same number of isogeny evaluations per prime degree,
 - ▶ discard computations in excess;
- $4\times$ slowdown (MCR) / $2.5\times$ slowdown (OAYT).
- Protected against SPA...

In seek of constant time

Two obstacles for constant time:


- 1 Some random points P may lack some factors; Unrelated to secret key if truly random.
- 2 Number of isogeny evaluations dependent on secret key.

Meyer, Campos, Reith 2018; Onuki, Aikawa, Yamazaki, Takagi 2019


- “Dummy” isogenies:
 - ▶ Always do exactly the same number of isogeny evaluations per prime degree,
 - ▶ discard computations in excess;
- $4\times$ slowdown (MCR) / $2.5\times$ slowdown (OAYT).
- Protected against SPA...but **very easy to attack by fault!**



Isogeny evaluation with dummies




Isogeny evaluation with dummies




$$\#\langle P \rangle = 3 \cdot 5 \cdot 7 \cdot 11$$




Isogeny evaluation with dummies




$$\#\langle P \rangle = 3 \cdot 5 \cdot 7 \cdot 11$$




Isogeny evaluation with dummies




$$\#\langle P \rangle = 3 \cdot 7 \cdot 11$$




Isogeny evaluation with dummies




$$\#\langle P \rangle = 3 \cdot 5 \cdot 7 \cdot 11$$




Isogeny evaluation with dummies




$$\#\langle P \rangle = 3 \cdot 5 \cdot 7 \cdot 11$$




Isogeny evaluation with dummies



$$\#\langle P \rangle = 3 \cdot 5 \cdot 7 \cdot 11$$



Isogeny evaluation with dummies



$$\#\langle P \rangle = 3 \cdot 5 \cdot 7 \cdot 11$$



Our work (Latincrypt 2019)

- Fixed a leak related to the sampling of random points.

Our work (Latincrypt 2019)

- Fixed a leak related to the sampling of random points.
- Speed-up both MCR and OAYT constant time implementations:
 - ▶ Fully Twisted Edwards implementation;
 - ▶ Use of Shortest Differential Addition Chains;

Our work (Latincrypt 2019)

- Fixed a leak related to the sampling of random points.
- Speed-up both MCR and OAYT constant time implementations:
 - ▶ Fully Twisted Edwards implementation;
 - ▶ Use of Shortest Differential Addition Chains;
- Protection against fault attack at the cost of a $2\times$ slowdown:
 - ▶ Got rid of “dummy isogenies”.

Our work (Latincrypt 2019)


- Fixed a leak related to the sampling of random points.
- Speed-up both MCR and OAYT constant time implementations:
 - ▶ Fully Twisted Edwards implementation;
 - ▶ Use of Shortest Differential Addition Chains;
- Protection against fault attack at the cost of a $2\times$ slowdown:
 - ▶ Got rid of “dummy isogenies”.
- Initiated study of fully constant time variant (very expensive, though).

Avoiding dummies

We change the format of the secret key:

Original: vectors with coefficients in $[-B, B]$.

Modified: vectors with **odd**¹ coefficients in $[-B, B]$.



¹Or **even**, all the same.

Avoiding dummies

We change the format of the secret key:

Original: vectors with coefficients in $[-B, B]$.

Modified: vectors with **odd**¹ coefficients in $[-B, B]$.

- Translate vector to **sum of ± 1 vectors**;
- Each vector costs exactly **one isogeny evaluation** per degree.

The diagram illustrates the decomposition of a vector with coefficients $[3, 5, -3, 1]$ into a sum of five vectors, each with coefficients in $\{-1, 1\}$. The original vector is represented by four colored bars: blue (3), red (5), green (-3), and orange (1). The red bar is the tallest, with a dotted line indicating its height. The decomposition is shown as an equation: the original vector equals the sum of five vectors. Each of these five vectors has a height of 1, represented by a dotted line. The coefficients are distributed as follows: the first three vectors have blue, red, and orange bars; the fourth has red and orange bars; and the fifth has red and green bars.

¹Or **even**, all the same.


Summary

- Repeat with me: I need isogeny-based crypto!
- CSIDH is the new Diffie–Hellman:
Very short keys, easy key validation, ...
- Implementing isogeny-based crypto efficiently is challenging, even more so with side-channel protections.



Thank you

<https://defeo.lu/>

 @luca_defeo