

The Millennium Problems

Seven Walls

Diego Rincón

Independent

Abstract

The Clay Mathematics Institute designated seven problems in 2000, each carrying a prize of one million dollars. Each is a Wall [1]: a place where two grammars press against each other and no crossing has been found. One has been solved. Six remain. This document states each problem precisely and names its Wall.

1 The Riemann Hypothesis

Conjecture 1 (Riemann, 1859). *The Riemann zeta function*

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}, \quad \operatorname{Re}(s) > 1,$$

extends to a meromorphic function on $\mathbb{C}$ with a simple pole at $s = 1$. Its non-trivial zeros — the zeros not at negative even integers — all satisfy

$$\operatorname{Re}(s) = \frac{1}{2}.$$

Remark 2 (The Wall). The Wall is between the spectral grammar (the zeros) and the arithmetic grammar (the primes). The explicit formula holds them in balance [2]. The missing step: self-adjointness of the operator D on $L^2(\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}^*, d^*x)$ [3]. Self-adjointness forces real spectrum. Real spectrum forces $\operatorname{Re}(\rho) = \frac{1}{2}$. The Wall is archimedean: the L^2 estimate at ∞ is what no proof has yet crossed.

2 The Birch and Swinnerton-Dyer Conjecture

Conjecture 3 (Birch–Swinnerton-Dyer, 1965). *Let E be an elliptic curve over $\mathbb{Q}$. The rank of the Mordell–Weil group $E(\mathbb{Q})$ equals the order of vanishing of the L -function $L(E, s)$ at $s = 1$:*

$$\operatorname{rank} E(\mathbb{Q}) = \operatorname{ord}_{s=1} L(E, s).$$

Moreover, the leading coefficient of the Taylor expansion of $L(E, s)$ at $s = 1$ is given by an explicit formula involving the period, the regulator, the Tate–Shafarevich group, and the Tamagawa numbers of E .

Remark 4 (The Wall). The Wall is between geometry ($E(\mathbb{Q})$, the rational points) and analysis ($L(E, s)$, the L -function). The geometric rank is a number counted by algebra. The analytic order of vanishing is a number counted by complex analysis. That they are equal is a resonance [4]: two grammars reading the same thing. Known: the conjecture holds when $\text{ord}_{s=1} L(E, s) \leq 1$ (Coates–Wiles, Gross–Zagier, Kolyvagin). Unknown: the general case.

3 The Hodge Conjecture

Conjecture 5 (Hodge, 1950). *Let X be a smooth projective variety over $\mathbb{C}$. Every Hodge class — every class in $H^{2k}(X, \mathbb{Q}) \cap H^{k,k}(X)$ — is a rational linear combination of classes of algebraic cycles.*

Remark 6 (The Wall). The Wall is between topology ($H^{2k}(X, \mathbb{Q})$) and algebra (algebraic cycles). A Hodge class is a cohomology class that looks algebraic from the analytic side. The conjecture says it is algebraic. The motive [5] is the proper design [6] from which both sides derive: if the standard conjectures hold, the Hodge conjecture follows. The Wall is the gap between the motivic grammar and the algebraic cycle grammar.

4 The P vs. NP Problem

Problem 7 (Cook, 1971). *Let P be the class of decision problems solvable in polynomial time by a deterministic Turing machine. Let NP be the class of decision problems whose solutions can be verified in polynomial time. Is $P = NP$?*

Remark 8 (The Wall). The Wall is between finding and checking [7]. Checking a proof is easy (NP). Finding a proof may be hard (P -hard). If $P = NP$: finding is as easy as checking. Every problem whose solution can be recognized can be solved just as fast. The Wall would not exist. If $P \neq NP$: the Wall is real. The universal Turing machine cannot short-circuit the difference between generation and verification. Almost everyone believes $P \neq NP$. No one has proved it.

5 Yang–Mills Existence and Mass Gap

Problem 9 (Jaffe–Witten, 2000). *For any compact simple gauge group G , there exists a quantum Yang–Mills theory on $\mathbb{R}^4$ satisfying the Wightman axioms, and there is a constant $\Delta > 0$ (the mass gap) such that the energy of every excited state exceeds the energy of the vacuum by at least Δ .*

Remark 10 (The Wall). The Wall is between the classical gauge theory (well-defined, rich in geometry) and the quantum theory (requiring a rigorous measure on the space of gauge connections). The Yang–Mills functional $\mathcal{A} \mapsto \int_{\mathbb{R}^4} |F_{\mathcal{A}}|^2$ is classical. Its quantization — a path integral over all connections — requires a mathematically rigorous construction that does not yet exist in four dimensions. The mass gap is the spectral gap of the quantum Hamiltonian: a gap in the nothing [8] between the vacuum and the first state.

6 Navier–Stokes Existence and Smoothness

Problem 11 (Fefferman, 2000). *In three space dimensions and time, given smooth initial data $u_0 : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ with $\nabla \cdot u_0 = 0$, does there exist a smooth solution $u : \mathbb{R}^3 \times [0, \infty) \rightarrow \mathbb{R}^3$ to the Navier–Stokes equations*

$$\partial_t u + (u \cdot \nabla)u = -\nabla p + \nu \Delta u, \quad \nabla \cdot u = 0,$$

for all time? Or can solutions blow up in finite time?

Remark 12 (The Wall). The Wall is between the local and the global: local smooth solutions exist (Leray, 1934); whether they extend to all time is unknown. The nonlinear term $(u \cdot \nabla)u$ can concentrate energy into smaller and smaller scales. Whether this concentration reaches a singularity or is smoothed by the viscosity ν is the question. Global regularity in 2D is proved (energy estimates close). In 3D, the energy estimates fail to close: the dimension is exactly critical for the nonlinearity. The Wall is at the critical dimension.

7 The Poincaré Conjecture

Conjecture 13 (Poincaré, 1904 — **Solved, Perelman 2003**). *Every simply connected, closed 3-manifold is homeomorphic to the 3-sphere S^3 .*

Remark 14 (The crossing). Perelman proved this in 2002–2003 using Hamilton’s Ricci flow with surgery: a geometric flow that deforms the metric of the manifold toward constant curvature. Singularities are removed by surgery. The flow terminates at a round sphere. Perelman declined the Clay prize. The Wall was crossed. The crossing used the geometry of the manifold itself as the proof: the manifold deformed into its own evidence.

8 The Six Remaining Walls

Remark 15 (What is left). Six Walls remain. Each has been approached. None has been crossed.

Problem	The Wall
Riemann Hypothesis	Spectral vs. arithmetic; archimedean estimate
BSD	Geometric rank vs. analytic order of vanishing
Hodge	Topology vs. algebraic cycles; motivic grammar
P vs. NP	Finding vs. checking; generation vs. verification
Yang–Mills	Classical gauge theory vs. quantum measure
Navier–Stokes	Local smoothness vs. global existence; critical dimension

Each Wall is a place where two grammars press against each other and no crossing has been found. The crossing, when it comes, will not be constructed. It will be found — already there, already vibrating [9], waiting for the grammar that can hear it.

References

- [1] Rincón, D. (2026). The Arithmetic Wall. Preprint.
- [2] Rincón, D. (2026). Week 09: The Resonance Object. Preprint.
- [3] Rincón, D. (2026). Week 10: Self-Adjointness. Preprint.
- [4] Rincón, D. (2026). Resonance: When Two Grammars Read the Same Thing. Preprint.
- [5] Rincón, D. (2026). Motives, Standard Conjectures, and the Unification of L -functions. Preprint.
- [6] Rincón, D. (2026). Design: The One That Precludes. Preprint.
- [7] Rincón, D. (2026). Turing Completeness and the Wall. Preprint.
- [8] Rincón, D. (2026). Nothing: The First Object. Preprint.
- [9] Rincón, D. (2026). The String. Preprint.