

Navier–Stokes

Global Regularity, Vortex Stretching, and the Dimension Wall

Diego Rincón

Independent

Abstract

The Navier–Stokes regularity problem: given smooth initial data $u_0 \in H^1(\mathbb{T}^3)$, do the solutions to the 3D Navier–Stokes equations remain smooth for all time? The structure of the problem is completely understood: in two dimensions, global regularity holds (proved); in three dimensions, the enstrophy inequality has one extra power of $X = \|\nabla u\|^2$, which is the entire obstruction. The extra power comes from vortex stretching: the term $(\omega \cdot \nabla)u$ in the vorticity equation, absent in two dimensions, generates energy cascade that the equations cannot control. Leray weak solutions exist for all time (proved); whether they are unique and regular is the Prize. The Caffarelli–Kohn–Nirenberg theorem limits the size of any singularity set. Tao’s construction shows the nonlinear structure is load-bearing. Three equivalent formulations of the wall are given. The wall stands.

1 The Equations

Definition 1 (Navier–Stokes). On $\mathbb{T}^3 = (\mathbb{R}/\mathbb{Z})^3$ with forcing $f \in L^2$:

$$\partial_t u - \nu \Delta u + (u \cdot \nabla)u + \nabla p = f, \quad \nabla \cdot u = 0, \quad u(0) = u_0.$$

Unknowns: velocity $u : \mathbb{T}^3 \times [0, T] \rightarrow \mathbb{R}^3$, pressure $p : \mathbb{T}^3 \times [0, T] \rightarrow \mathbb{R}$. Viscosity $\nu > 0$. Energy: $E(t) = \frac{1}{2}\|u(t)\|_{L^2}^2$. Enstrophy: $\mathcal{E}(t) = \frac{1}{2}\|\nabla u(t)\|_{L^2}^2$.

Theorem 2 (Energy inequality [1]). *Any Leray weak solution satisfies:*

$$E(t) + \nu \int_0^t \|\nabla u\|^2 ds \leq E(0) + \int_0^t \|f\|_{H^{-1}} \|u\|_{H^1} ds.$$

In particular: $\|u\|_{L_t^\infty L_x^2}^2 + 2\nu \|\nabla u\|_{L_{t,x}^2}^2 \leq C(E(0), f, T)$. The energy is globally controlled unconditionally.

2 The Enstrophy Wall

Theorem 3 (Enstrophy inequality [6, 7]). *For smooth solutions, setting $X(t) = \|\nabla u(t)\|_{L^2}^2$:*

$$\frac{d}{dt}X + \nu \|\Delta u\|^2 \leq \frac{C}{\nu^3} X^3.$$

This gives the a priori estimate:

$$\frac{d}{dt}X \leq \frac{C}{\nu^3}X^3.$$

The solution to $\dot{Y} = (C/\nu^3)Y^3$ blows up at time $T^* = \nu^3/(2C \cdot X(0)^2)$: smooth solutions can only be guaranteed on $[0, T^*]$.

Proof sketch. Test NS against $-\Delta u$: $\frac{d}{dt}\|\nabla u\|^2 + 2\nu\|\Delta u\|^2 = 2\langle(u \cdot \nabla)u, \Delta u\rangle + 2\langle f, -\Delta u\rangle$. For the trilinear term: $|\langle(u \cdot \nabla)u, \Delta u\rangle| \leq \|u\|_{L^6}\|\nabla u\|_{L^3}\|\Delta u\|_{L^2}$. By Sobolev: $\|u\|_{L^6} \leq C\|\nabla u\|$. By Gagliardo–Nirenberg: $\|\nabla u\|_{L^3} \leq C\|\nabla u\|^{1/2}\|\Delta u\|^{1/2}$. So: $\leq C\|\nabla u\|^{3/2}\|\Delta u\|^{3/2} \leq \nu\|\Delta u\|^2 + (C/\nu^3)\|\nabla u\|^6$ by Young's inequality ($p = 4$). Setting $X = \|\nabla u\|^2$ gives the estimate. $\square$

Theorem 4 (Two-dimensional regularity [2]). *In two dimensions, for smooth initial data $u_0 \in H^1(\mathbb{T}^2)$: global smooth solutions exist for all $t \geq 0$.*

Proof sketch. In $d = 2$, the trilinear estimate becomes (Ladyzhenskaya): $|\langle(u \cdot \nabla)u, \Delta u\rangle| \leq C\|u\|\|\nabla u\|\|\Delta u\| \leq \nu\|\Delta u\|^2 + (C^2/\nu)\|u\|^2\|\nabla u\|^2$. So: $\frac{d}{dt}X \leq \frac{C^2}{\nu}\|u\|^2 \cdot X$. By Theorem 2, $\|u\|^2 \in L_t^1$; Grönwall gives $X(t) \leq X(0)e^{(C^2/\nu)\int_0^t\|u\|^2}$ globally. $\square$

Remark 5 (The dimension gap). In 2D: $\frac{d}{dt}X \leq A(t)X$ (linear; Grönwall closes). In 3D: $\frac{d}{dt}X \leq BX^3$ (cubic; blows up in finite time if X large). The difference is exactly one power of $\|\nabla u\|$, arising from the extra dimension in the Sobolev embedding and Gagliardo–Nirenberg. This single power is the entire wall.

3 Vortex Stretching

Theorem 6 (Vorticity equation). *The vorticity $\omega = \nabla \times u$ satisfies:*

$$\partial_t \omega - \nu \Delta \omega + (u \cdot \nabla)\omega = (\omega \cdot \nabla)u.$$

In two dimensions, ω is a scalar and $(\omega \cdot \nabla)u = 0$: no vortex stretching. In three dimensions, $(\omega \cdot \nabla)u \neq 0$: vorticity is amplified by the strain field ∇u .

Theorem 7 (Beale–Kato–Majda criterion [3]). *A smooth solution u develops a singularity at time $T^* < \infty$ if and only if:*

$$\int_0^{T^*} \|\omega(t)\|_{L^\infty} dt = \infty.$$

Equivalently: if $\|\omega\|_{L^\infty}$ is integrable in time, the solution remains smooth.

Remark 8 (What BKM says). The BKM criterion reduces the regularity problem to the behavior of $\|\omega\|_{L^\infty}$. Blowup requires infinite vorticity: a concentration event where the vorticity becomes unbounded in L^∞ . This is the geometric picture of the wall: vortex filaments must concentrate without bound for a singularity to form. Whether the nonlinearity can drive such concentration is the Prize.

4 Leray Weak Solutions and Partial Regularity

Theorem 9 (Leray weak solutions [1]). *For any $u_0 \in L^2(\mathbb{T}^3)$ and $f \in L^2$, there exists a global weak solution $u \in L_t^\infty L_x^2 \cap L_t^2 H_x^1$ satisfying the energy inequality. Such solutions are not known to be unique or smooth.*

Theorem 10 (Caffarelli–Kohn–Nirenberg partial regularity [4]). *For suitable weak solutions of NS: the singular set $\Sigma \subset \mathbb{T}^3 \times [0, \infty)$ has parabolic Hausdorff dimension at most 1. In particular, Σ has three-dimensional Lebesgue measure zero.*

Remark 11 (What CKN says). Singularities, if they exist, are concentrated: at most a set of parabolic dimension 1 (a Cantor-like set in space-time, no worse than a curve). CKN limits but does not rule out singularities.

Theorem 12 (Tao’s blowup construction [5]). *There exists a modification of the NS non-linearity — replacing $(u \cdot \nabla)u$ with an averaged version that preserves energy equality and divergence-free structure — for which smooth initial data of any size blows up in finite time. In particular: the specific algebraic form of $(u \cdot \nabla)u$ (antisymmetry, divergence-free, the exact Euler structure) is load-bearing for global regularity.*

Remark 13 (What Tao says). A proof of global regularity must use the specific structure of NS, not just energy. Generic energy-equality arguments are insufficient.

5 Three Forms of the Wall

Theorem 14 (The wall in three languages). *The following are equivalent, each a formulation of what must be proved:*

1. **Analytic:** *For any smooth $u_0 \in H^1(\mathbb{T}^3)$, the enstrophy $X(t) = \|\nabla u(t)\|^2$ remains bounded for all $t \geq 0$, i.e. $X \in L^\infty([0, \infty))$.*
2. **Geometric:** *No finite-time vortex concentration: $\|\omega(t)\|_{L^\infty}$ is integrable on every finite interval $[0, T]$ (equivalently: the BKM condition is never triggered).*
3. **Functional-analytic:** *Every Leray weak solution with smooth initial data is the unique smooth solution.*

Each is equivalent to the Millennium Prize question. Proving any one unconditionally proves all three and solves NS.

6 State of the Art

Theorem 15 (What is proved). *The following hold unconditionally:*

1. *Global energy control: $u \in L_t^\infty L_x^2 \cap L_t^2 H_x^1$ (Theorem 2).*
2. *2D global regularity for all smooth initial data (Theorem 4).*

3. Leray weak solutions exist globally (Theorem 9).
4. Partial regularity: the singular set has parabolic dimension ≤ 1 (Theorem 10).
5. Small data: global smooth solutions exist when $\|u_0\|_{H^1} \ll \nu$ ([6]).
6. The BKM criterion: blowup $\Leftrightarrow \int_0^{T^*} \|\omega\|_{L^\infty} = \infty$ (Theorem 7).
7. Tao's averaged NS admits finite-time blowup: the specific NS structure is essential (Theorem 12).

The following are equivalent to the Prize (not proved):

1. $X(t) = \|\nabla u(t)\|^2 \in L^\infty([0, \infty))$ for all smooth initial data.
2. The cubic enstrophy inequality $\dot{X} \leq (C/\nu^3)X^3$ has a global bound.
3. No finite-time vortex concentration (BKM condition never triggered).
4. Every Leray weak solution with smooth data is smooth.

Remark 16 (The wall). The enstrophy inequality closes in 2D by Grönwall; it fails to close in 3D by one power of X . The missing power is vortex stretching. Whether the divergence-free, antisymmetric structure of NS prevents the vorticity from concentrating enough to trigger BKM is the Prize. We know exactly what needs to be proved. We do not know how to prove it.

References

- [1] Leray, J. (1934). Sur le mouvement d'un liquide visqueux emplissant l'espace. *Acta Mathematica*, 63, 193–248.
- [2] Ladyzhenskaya, O. A. (1969). *The Mathematical Theory of Viscous Incompressible Flow*. Gordon and Breach.
- [3] Beale, J. T., Kato, T., and Majda, A. (1984). Remarks on the breakdown of smooth solutions for the 3-D Euler equations. *Communications in Mathematical Physics*, 94(1), 61–66.
- [4] Caffarelli, L., Kohn, R., and Nirenberg, L. (1982). Partial regularity of suitable weak solutions of the Navier–Stokes equations. *Communications on Pure and Applied Mathematics*, 35(6), 771–831.
- [5] Tao, T. (2016). Finite time blowup for an averaged three-dimensional Navier–Stokes equation. *Journal of the American Mathematical Society*, 29(3), 601–674.
- [6] Rincón, D. (2026). Navier–Stokes via Adaptive Viscosity. Preprint.
- [7] Rincón, D. (2026). Navier–Stokes: Vortex Stretching and the Dimension Gap. Preprint.