

Zeros of L -Functions: Depth, Rank, and the Arithmetic at the Critical Point

Diego Rincón

Independent

Abstract

Every major open problem in arithmetic geometry is a statement about a zero of an L -function. The Riemann Hypothesis asks where the zeros of $\zeta(s)$ lie. BSD asks for the order of vanishing of $L(E, s)$ at $s = 1$. The Bloch–Kato conjecture identifies the order of vanishing of $L(M, s)$ at $s = 0$ with the dimension of a Selmer group. Beilinson’s conjectures identify the leading coefficient with a regulator. In each case, the zero is not absence: it is depth. The order of vanishing is a measure of arithmetic complexity — the rank of the Mordell–Weil group, the dimension of the Selmer group, the rank of the motivic cohomology. We collect these instances into a single picture: the L -function as a probe of arithmetic, its zeros as the wells where arithmetic lives, and the leading coefficient as the volume of the well. The functional equation, the Euler product, and the critical strip are the geometry of the probe. The open conjectures are the question of how deep each well is.

1 The Structure of an L -Function

Definition 1 (L -function). An L -function is a Dirichlet series $L(s) = \sum_{n=1}^{\infty} a_n n^{-s}$, convergent in some right half-plane $\operatorname{Re}(s) > \sigma_0$, satisfying:

1. (Euler product) $L(s) = \prod_p L_p(s)^{-1}$, where $L_p(s) = P_p(p^{-s})$ for a polynomial P_p of bounded degree.
2. (Analytic continuation) $L(s)$ extends to a meromorphic function on $\mathbb{C}$, holomorphic except possibly at $s = 0$ and $s = 1$.
3. (Functional equation) There exists a completed L -function $\Lambda(s) = N^{s/2} \cdot \Gamma_{\infty}(s) \cdot L(s)$ satisfying $\Lambda(s) = \varepsilon \cdot \Lambda(k - s)$ for some integer k (the *weight*), conductor N , and root number $\varepsilon \in \{+1, -1\}$.

The *critical strip* is $0 < \operatorname{Re}(s) < k$. The *central point* is $s = k/2$.

Remark 2 (The zero as a well). A zero of L at $s = s_0$ of order r means: $L(s) = (s - s_0)^r \cdot c + O((s - s_0)^{r+1})$, $c \neq 0$. The order r is the *depth* of the zero. The leading coefficient $c = L^{(r)}(s_0)/r!$ is the *volume*: the arithmetic content stored at depth r . A zero of order 0 (a non-vanishing value) has depth 0 — no well. A zero of order $r \geq 1$ is a well of depth r . The open conjectures assert that the depth equals the rank of a specific arithmetic object.

2 The Riemann Hypothesis: Location of the Wells

Conjecture 3 (Riemann Hypothesis [1]). *All non-trivial zeros of $\zeta(s)$ (zeros in the critical strip $0 < \operatorname{Re}(s) < 1$) lie on the critical line $\operatorname{Re}(s) = 1/2$.*

Remark 4 (What the location means). The functional equation $\zeta(s) = \zeta(1-s)$ (up to gamma factors) reflects zeros across $\operatorname{Re}(s) = 1/2$. A zero off the critical line would come in a quadruple $\rho, 1-\rho, \bar{\rho}, 1-\bar{\rho}$ (by the functional equation and the reflection principle $\overline{\zeta(s)} = \zeta(\bar{s})$). The Riemann Hypothesis asserts that no such quadruple exists: all wells are on the central axis. The Weil conjectures [2] prove the analogue over $\mathbb{F}_q$: every zero of $Z(X, q^{-s})$ lies on $\operatorname{Re}(s) = i/2$ for some integer i (purity). The Riemann Hypothesis is the case $X = \operatorname{Spec} \mathbb{Z}$.

Theorem 5 (Distribution of zeros: GUE statistics). *Montgomery (1973) and Odlyzko (1987) found numerically and conjectured that the pair correlation of the imaginary parts of the Riemann zeros follows the distribution of eigenvalues of random matrices in the Gaussian Unitary Ensemble (GUE). This is the statistical signature of a self-adjoint operator — the Hilbert–Pólya operator H whose spectrum is the set of zeros. The wells of ζ are distributed as if they were energy levels of a quantum system. The quantum system has not been found.*

3 BSD: Depth Equals Rank

Conjecture 6 (BSD, rank part [3]). *For an elliptic curve $E/\mathbb{Q}$, the L -function $L(E, s)$ has a zero at $s = 1$ of order exactly $\operatorname{rank} E(\mathbb{Q})$:*

$$\operatorname{ord}_{s=1} L(E, s) = \operatorname{rank} E(\mathbb{Q}).$$

Remark 7 (The well of BSD). The zero of $L(E, s)$ at $s = 1$ is the well of BSD. Its depth (order of vanishing) is the rank — the number of independent rational points. A curve of rank 0 has no well: $L(E, 1) \neq 0$. A curve of rank 1 has a well of depth 1: one rational generator, one unit of arithmetic depth. A curve of rank r has a well of depth r . The volume of the well (the leading coefficient) encodes: the regulator, the Tate-Shafarevich group, the Tamagawa numbers, and the periods — the full arithmetic of E .

Theorem 8 (Known cases [6]). *1. $\operatorname{ord}_{s=1} L(E, s) = 0$ implies $\operatorname{rank} E(\mathbb{Q}) = 0$ (Kolyvagin–Kato).*

2. $\operatorname{ord}_{s=1} L(E, s) = 1$ implies $\operatorname{rank} E(\mathbb{Q}) = 1$ (Gross–Zagier, Kolyvagin).

3. $\operatorname{ord}_{s=1} L(E, s) \geq 2$: rank ≥ 2 is unproved in general. The well has depth, but we cannot yet reach the bottom.

4 Bloch–Kato: Depth Equals Selmer Dimension

Conjecture 9 (Bloch–Kato, rank part [4]). *For a pure motive M of weight -1 :*

$$\operatorname{ord}_{s=0} L(M, s) = \dim H_f^1(G_{\mathbb{Q}}, V) - \dim H^0(G_{\mathbb{Q}}, V) - \dim H^0(G_{\mathbb{Q}}, V^*(1)).$$

The depth of the well at $s = 0$ equals the net dimension of the Bloch–Kato Selmer group.

Proposition 10 (Unification of depth). *BSD is the case $M = h^1(E)(1)$, $s = 1$ shifted to $s = 0$ by Tate twist. Beilinson’s conjecture for $h^{2k-1}(X)(k)$ is the general case. The Dedekind zeta function $\zeta_k(s)$ at $s = 0$ gives $\text{ord}_{s=0}\zeta_k = -r_1 - r_2 + 1$ (Dirichlet’s unit theorem), the simplest case: depth equals rank of the unit group. Every case is the same statement: depth of the well equals rank of the relevant arithmetic group.*

5 The Leading Coefficient: Volume of the Well

Definition 11 (Leading coefficient). For $L(M, s)$ with a zero of order r at $s = s_0$:

$$L^*(M, s_0) = \lim_{s \rightarrow s_0} (s - s_0)^{-r} L(M, s).$$

This is the *volume of the well*: the arithmetic content at depth r .

Conjecture 12 (Leading coefficient conjectures). 1. (BSD full formula) $L^*(E, 1) = \Omega_E \cdot R(E) \cdot \#\text{III}(E) \cdot \prod_\ell c_\ell / (\#E(\mathbb{Q})_{\text{tors}})^2$.

2. (Bloch–Kato) $L^*(M, 0) = \tau(M) \cdot R(M) \cdot \Omega(M)$, $\tau(M) \in \mathbb{Q}^\times$ ([4]).

3. (Beilinson) *The regulator $R(M)$ is the determinant of the period–cycle pairing on motivic cohomology* ([5]).

In each case: volume = regulator $\times$ period $\times$ (arithmetic correction). The arithmetic correction (Tate-Shafarevich group, Tamagawa numbers) is the discrete part; the regulator is the continuous part; the period bridges them.

6 The Unified Picture

Theorem 13 (All open problems as well-zero statements).

<i>Problem</i>	<i>L-function</i>	<i>Zero at</i>	<i>Depth =</i>
<i>Riemann Hypothesis</i>	$\zeta(s)$	$\text{Re}(s) = 1/2$	<i>(location, not depth)</i>
<i>BSD</i>	$L(E, s)$	$s = 1$	$\text{rank } E(\mathbb{Q})$
<i>Bloch–Kato</i>	$L(M, s)$	$s = 0$	$\dim H_f^1$
<i>Beilinson</i>	$L(h^{2k-1}(X), s)$	$s = 0, \dots, 1 - k$	$\dim H_{\mathcal{M}}^{2k-1}$
<i>Tate conjecture</i>	$L(h^{2r}(X), s)$	$s = r$	$\dim \text{NS}^r$

Every conjecture asserts that a certain depth (order of vanishing) equals a certain rank (of an arithmetic or motivic group). The leading coefficient conjecture asserts that the volume of the well equals a product of geometric and arithmetic invariants. The Arithmetic Wall [7] is the obstruction to reading the depth from the well: locally at each prime, the depth is accessible (via Frobenius eigenvalues); globally over $\mathbb{Q}$, the reading requires what does not yet exist.

Remark 14 (The Riemann Hypothesis as location, not depth). The Riemann Hypothesis is different from the other entries in the table. It does not ask for the depth of a zero but for the location of all zeros. In the well metaphor: it asks not how deep each well is, but

that all wells lie on a single axis. This is a statement about the symmetry of the probe, not the arithmetic being probed. The proof requires constructing the Hilbert–Pólya operator — the instrument that makes the symmetry visible. All other conjectures in the table are statements about the wells. RH is a statement about the ground they are dug into.

References

- [1] Riemann, B. (1859). Über die Anzahl der Primzahlen unter einer gegebenen Größe. *Monatsber. Königl. Preuss. Akad. Wiss. Berlin*, 671–680.
- [2] Rincón, D. (2026). The Weil Conjectures as Proof of Concept for the Riemann Hypothesis. Preprint.
- [3] Rincón, D. (2026). BSD via Arithmetic Gan–Gross–Prasad. Preprint.
- [4] Rincón, D. (2026). The Bloch–Kato Conjecture. Preprint.
- [5] Rincón, D. (2026). Motives, Standard Conjectures, and the Unification of L -functions. Preprint.
- [6] Rincón, D. (2026). Iwasawa Theory: The p -adic Machine for BSD and Beyond. Preprint.
- [7] Rincón, D. (2026). The Arithmetic Wall. Preprint.