

# Probability and Computing, 2nd Edition

Solutions to Chapter 4: Chernoff and Hoeffding Bounds

Hahndeul Kim

July 2025

## 4.1

Let the number of games that Alice wins be  $X$ , where  $X \sim B(n, 0.6)$ . Alice will lose the tournament with probability  $\Pr(X \leq \frac{n-1}{2})$ . Now, let  $\delta$  s. t.  $(1 - \delta) \times \frac{3n}{5} = \frac{n-1}{2}$  to obtain the tightest bound.  
 $\Pr(X \leq \frac{n-1}{2}) = \Pr(X \leq (1 - \delta)\mathbf{E}[X]) \leq \exp(-\frac{3n}{5} \cdot \delta^2 \cdot \frac{1}{2})$   
 $= \exp(-\frac{1}{10}(\frac{1}{12}n + \frac{5}{6} + \frac{25}{12n})) \leq \exp(-\frac{1}{8})$  (AM-GM inequality).

## 4.2

With Markov's inequality,  $\Pr(X \geq n/4) \leq (n/6)/(n/4) = 2/3$ .

With Chebyshev's inequality,  $\Pr(X \geq n/4) \leq \Pr(|X - n/6| \geq n/12) \leq \frac{\mathbf{Var}[X]}{(n/12)^2}$   
 $= \frac{144}{n^2} \times (n \cdot \frac{1}{6} \cdot \frac{5}{6}) = 20/n$ .

To use Chernoff bounds, let  $\delta = 1/2$ . Then  $\Pr(X \geq n/4) = \Pr(X \geq (1+\delta)\mathbf{E}[X])$   
 $\leq \left(\frac{e^{0.5}}{1.5^{1.5}}\right)^{n/6} = \left(\frac{e}{1.5^3}\right)^{n/12}$ .

## 4.3

(a) Let  $X \sim B(n, p)$ . Then  $M_X(t) = \mathbf{E}[e^{tX}] = \sum_{i=0}^n e^{it} \Pr(X = i)$

$$= \sum_{i=0}^n e^{it} \binom{n}{i} p^i (1-p)^{n-i} = \sum_{i=0}^n \binom{n}{i} (pe^t)^i (1-p)^{n-i} = (pe^t + 1 - p)^n.$$

(b)  $M_{X+Y}(t) = \mathbf{E}[e^{t(X+Y)}] = \mathbf{E}[e^{tX} e^{tY}] = \mathbf{E}[e^{tX}] \mathbf{E}[e^{tY}] = (pe^t + 1 - p)^{m+n}$ .

(c) Since moment generating function uniquely determines the distribution,  $X + Y \sim B(m + n, p)$ .

## 4.4

Let the total number of heads be  $X$ , where  $X \sim B(100, \frac{1}{2})$ . Then we find  $\Pr(X \geq 55) \approx 0.1841$ .

From Chernoff bound, we find that  $\Pr(X \geq (1 + \frac{1}{10})50) \leq \exp(-\frac{50}{3} \cdot \frac{1}{10^2}) = \exp(-\frac{1}{6}) \approx 0.8465$ .

For  $Y \sim B(1000, \frac{1}{2})$ ,  $\Pr(Y \geq 550) \approx 0.0009$ .

From Chernoff bound, we find that  $\Pr(Y \geq (1 + \frac{1}{10})500) \leq \exp(-\frac{500}{3} \cdot \frac{1}{10^2}) = \exp(-\frac{5}{3}) \approx 0.1889$ .

## 4.5

Let  $Y = NX$ , so that we aim to satisfy  $\Pr(|Y - Np| > N\epsilon p) \leq \delta$ . Consider that  $\Pr(Y > Np(1 + \epsilon)) < \exp(-Np \cdot \frac{\epsilon^2}{3})$ , and  $\Pr(Y < Np(1 - \epsilon)) < \exp(-Np \cdot \frac{\epsilon^2}{2})$ . Thus, we aim to satisfy  $\exp(-Np \cdot \frac{\epsilon^2}{3}) + \exp(-Np \cdot \frac{\epsilon^2}{2}) \leq 2 \exp(-Np \cdot \frac{\epsilon^2}{3}) \leq \delta$ .

$\therefore N \geq \frac{3}{p\epsilon^2} \ln \frac{2}{\delta}$ . With  $\epsilon = 0.1$ ,  $\delta = 0.05$  and  $0.2 \leq p \leq 0.8$ ,  $N \geq 1500 \ln 40 \approx 5533$ .

## 4.6

(a) Let  $X \sim B(1000000, 0.02)$ . Then  $\Pr(X \geq 40000) \leq e^{-20000/3}$ .  
(b) Set  $X$  and  $Y$  as given and choose  $k, l$  such that  $l \leq k - 10000$  so that bounding  $\Pr((X > k) \cap (Y < l))$  suffices. As examples, we choose  $k = 15300$  and  $l = 4900$  here. Since  $X \sim B(510000, 0.02)$ ,  $Y \sim B(490000, 0.02)$  and  $X \perp\!\!\!\perp Y$ ,  $\Pr((X > k) \cap (Y < l)) = \Pr(X > k) \Pr(Y < l) \leq e^{-10200/12} \times e^{-9800/8} = e^{-2025}$ .

## 4.7

Recall that  $M_X(t) = \prod_{i=1}^n (p_i e^t + (1 - p_i)) = \prod_{i=1}^n (1 + p_i(e^t - 1)) \leq \prod_{i=1}^n e^{p_i(e^t - 1)}$

$= e^{\mu(e^t - 1)}$  holds when  $X$  is the sum of Poisson trials ( $\Pr(X_i = 1) = p_i$ ).

Let  $t = \ln(1 + \delta)$  and follow the derivation of Chernoff bounds.

$$\Pr(X \geq (1 + \delta)\mu_H) \leq \frac{\mathbf{E}[e^{tX}]}{e^{t(1+\delta)\mu_H}} \leq \frac{e^{\mu(e^t - 1)}}{e^{t(1+\delta)\mu_H}} \leq \left( \frac{e^{e^t - 1}}{e^{t(1+\delta)}} \right)^{\mu_H} = \left( \frac{e^\delta}{(1+\delta)^{(1+\delta)}} \right)^{\mu_H}.$$

Similarly, let  $t = \ln(1 - \delta)$  and prove the latter inequality.

$$\Pr(X \leq (1 - \delta)\mu_L) \leq \frac{\mathbf{E}[e^{tX}]}{e^{t(1-\delta)\mu_L}} \leq \frac{e^{\mu(e^t - 1)}}{e^{t(1-\delta)\mu_L}} \leq \left( \frac{e^{e^t - 1}}{e^{t(1-\delta)}} \right)^{\mu_L} = \left( \frac{e^{-\delta}}{(1-\delta)^{(1-\delta)}} \right)^{\mu_L}. \blacksquare$$

## 4.8

For any permutation  $\pi$  produced with the given approach,  $\Pr(f = \pi) = \prod_{i=1}^n \frac{1}{k+1-i}$

holds. Since the number of possible permutations is  $\frac{k!}{(k-n)!} = \frac{1}{\Pr(f=\pi)}$ , the given approach produces a permutation chosen uniformly at random from all permutations.

Now, let  $X_j$  be the number of black box calls to determine  $f(j)$ . Then  $X_j \sim \text{Geom}(\frac{k+1-j}{k})$  holds. Thus,  $\mathbf{E}[\sum_{i=1}^n X_i] = \sum_{i=1}^n \mathbf{E}[X_i] = \sum_{i=1}^n \frac{k}{k+1-i}$ .

When  $k = n$ ,  $\mathbf{E}[\sum_{i=1}^n X_i] = \sum_{i=1}^n \frac{n}{i} = nH(n) \approx n \ln n$ .

Similarly, when  $k = 2n$ ,  $\mathbf{E}[\sum_{i=1}^n X_i] = \sum_{i=1}^n \frac{2n}{n+i} = 2n(H(2n) - H(n)) \approx 2n \ln 2$ . In

this case,  $\frac{2n+1-j}{2n} \geq \frac{2n+1-n}{2n} \geq \frac{1}{2}$ .

Now, to derive the desired Chernoff bound, we first compute the moment generating function of  $X = \sum_{i=1}^n X_j$ . Let  $p_i = \frac{2n+1-i}{2n}$ . Since  $X_i$  are independent,

$$\mathbf{E}[e^{tX}] = \prod_{i=1}^n \mathbf{E}[e^{tX_i}] = \prod_{i=1}^n \left( \prod_{j=1}^{\infty} (e^{tj} p_i (1 - p_i)^{j-1}) \right) = \prod_{i=1}^n \left( \frac{p_i}{1-p_i} \prod_{j=1}^{\infty} (e^t (1 - p_i))^j \right).$$

Suppose that we choose  $t$  s. t.  $0 < t < \ln 2$  when deriving the Chernoff bound.

Then  $\mathbf{E}[e^{tX}] = \prod_{i=1}^n \frac{p_i e^t}{1 - e^t(1-p_i)}$ . Since  $t > 0$ ,  $\frac{\partial}{\partial p_i} \left( \frac{p_i e^t}{1 - e^t(1-p_i)} \right) = \frac{1 - e^t}{(1 - e^t(1-p_i))^2} < 0$ .

This leads to  $\mathbf{E}[e^{tX}] = \prod_{i=1}^n \frac{p_i e^t}{1 - e^t(1-p_i)} \leq \left( \frac{\frac{1}{2} e^t}{1 - \frac{1}{2} e^t} \right)^n$ .

Now derive the desired Chernoff bound with  $\Pr(X \geq 4n) \leq \frac{\mathbf{E}[e^{tX}]}{e^{4nt}} \leq \left( \frac{1}{(2 - e^t)e^{3t}} \right)^n$ .

Since the function  $(2 - e^t)e^{3t}$  has its maximum at  $t = \ln \frac{3}{2}$  and  $0 < \ln \frac{3}{2} < \ln 2$ , we choose  $t = \ln \frac{3}{2}$  for the tightest possible bound.

The desired bound would be  $\Pr(X \geq 4n) \leq \left( \frac{1}{(2 - e^t)e^{3t}} \right)^n \Big|_{t=\ln \frac{3}{2}} = \left( \frac{16}{27} \right)^n$ .

## 4.9

(a) By Chebyshev's inequality,  $\Pr\left[\left|\sum_{i=1}^t X_i - \mathbf{E}[X]\right| \geq \epsilon \mathbf{E}[X]\right] \leq \frac{\mathbf{Var}[X]}{t(\epsilon \mathbf{E}[X])^2} = \frac{r^2}{t\epsilon^2}$ .

Thus, setting  $t$  to satisfy  $\frac{r^2}{t\epsilon^2} \leq \delta$  suffices. This leads to  $t \geq \frac{r^2}{\epsilon^2 \delta}$ , which proves the claim.

(b) Set  $\delta = 1 - 3/4 = 1/4$ . Then we get  $t \geq \frac{4r^2}{\epsilon^2}$ , which proves the claim.

(c) Let  $Y_i$  be indicator variables that are 1 if  $|X_i - \mathbf{E}[X]| \geq \epsilon \mathbf{E}[X]$ . Then let the median of  $Y_i$ s be  $m$ , and bound the probability  $\Pr(|m - \mathbf{E}[X]| \geq \epsilon \mathbf{E}[X])$ .

Note that  $\mathbf{E}[\sum_{i=1}^t Y_i] \leq t/4$  by definition, and  $|m - \mathbf{E}[X]| \geq \epsilon \mathbf{E}[X]$  holds only

if  $\sum_{i=1}^t Y_i \geq t/2$ . Then,  $\Pr(|m - \mathbf{E}[X]| \geq \epsilon \mathbf{E}[X]) \leq \Pr\left(\sum_{i=1}^t Y_i \geq t/2\right)$ . Let

$Y = \sum_{i=1}^t Y_i$ . Then  $\Pr(Y \geq t/2) = \Pr\left(Y \geq (1 + (\frac{t}{2\mathbf{E}[Y]} - 1))\mathbf{E}[Y]\right)$

$\leq \left(\frac{e^\delta}{(1+\delta)^{1+\delta}}\right)^{\mathbf{E}[Y]} = \left(\frac{2e}{t}\right)^{t/2} \times e^{-\mathbf{E}[Y]} \mathbf{E}[Y]^{t/2}$ .

Since  $\frac{\partial}{\partial \mathbf{E}[Y]} \left( \left(\frac{2e}{t}\right)^{t/2} e^{-\mathbf{E}[Y]} \mathbf{E}[Y]^{t/2} \right) = \left(\frac{2e}{t}\right)^{t/2} e^{-\mathbf{E}[Y]} \mathbf{E}[Y]^{t/2-1} (t/2 - \mathbf{E}[Y]) > 0$ ,

substitute  $t/4$  for  $\mathbf{E}[Y]$  to derive our bound. Thus,  $\Pr(Y \geq t/2) \leq \left(\frac{e}{4}\right)^{t/4}$ .

Here we need  $t$  that satisfies  $(\frac{e}{4})^{t/4} \leq \delta$ , which leads to  $t \geq \frac{4}{\ln \frac{4}{e}} \ln \frac{1}{\delta}$ . Therefore, together with 4.9.(b), we only need  $O(\log(1/\delta))$  estimates constructed from  $O(r^2 \log(1/\delta)/\epsilon^2)$  samples.

## 4.10