

Convergence of SMACOF

Jan de Leeuw

July 16, 2026

TBD

Table of contents

1	Introduction	1
2	Notation	3
3	SMACOF	4
4	Global Convergence of SMACOF	5
5	Local Convergence of SMACOF	7
6	Modified SMACOF	9
7	Coefficient Space	10
8	Software	11
9	Examples	12
9.1	Ekman	12
9.2	De Gruijter	14
10	Conclusion	17
	References	18

1 Introduction

The (Euclidean, metric, least squares, p-dimensional) multidimensional scaling (MDS) problem is the minimization of the real valued loss function ¹

$$\sigma(X) := \frac{1}{2} \sum_{1 \leq i < j \leq n} w_{ij} (\delta_{ij} - d_{ij}(X))^2, \quad (1)$$

over *configurations* $X \in \mathcal{X}$, where *configuration space* $\mathcal{X}$ is a subset of $\mathbb{R}^{n \times p}$, the space of all $n \times p$ matrices, with $1 < p < n - 1$. If $\mathcal{X}$ is all of $\mathbb{R}^{n \times p}$ the MDS problem is *unconstrained*, if $\mathcal{X}$ is a proper subset it is *constrained*.

In (1) the *dissimilarities* Δ and *weights* W are the *data* of the problem. Both Δ and W are symmetric, non-negative, and hollow² matrices Δ of order n . The $d_{ij}(X)$ are Euclidean distances between rows x_i and x_j of X . Thus

$$d_{ij}(X) := \sqrt{(x_i - x_j)'(x_i - x_j)}. \quad (2)$$

We call $\sigma(X)$ the *stress* of configuration X (Kruskal (1964a), Kruskal (1964b)). Kruskal does not use weights in his definition and calls our (1) *raw stress*. Since he is only interested in non-metric MDS his definition of stress (or *normalized stress*) adds the sum of squared distances as a denominator in (1).

In this paper we assume that the dimensionality p satisfies $1 < p < n - 1$. Both $p = 1$ and $p = n - 1$ are excluded, because these boundary cases of MDS have special properties which makes them fundamentally different from MDS for the intermediate values of p (De Leeuw (2023), De Leeuw et al. (2016)). We also assume, without loss of generality, that W is *irreducible*, which means in this context that the MDS problem does not separate into a number of smaller MDS problems that can be solved independently (De Leeuw (1977)).

¹The symbol $:=$ is used for definitions.

²A hollow matrix has a zero diagonal.

2 Notation

First some convenient notation, introduced in De Leeuw (1977). A *unit vector* e_i has n elements, with element i equal to $+1$, and all other elements zero.

$$A_{ij} := (e_i - e_j)(e_i - e_j)', \quad (3)$$

which means elements (i, i) and (j, j) are equal to $+1$, while (i, j) and (j, i) are -1 . All other elements are zero. Then

$$d_{ij}^2(X) = \text{tr } X' A_{ij} X, \quad (4)$$

We also define the matrix

$$V := \sum_{1 \leq i < j \leq n} w_{ij} A_{ij}, \quad (5)$$

and the matrix-valued function B with

$$B(X) := \sum_{1 \leq i < j \leq n} w_{ij} r_{ij}(X) A_{ij}, \quad (6)$$

where

$$r_{ij}(X) := \begin{cases} \frac{\delta_{ij}}{d_{ij}(X)} & \text{if } d_{ij}(X) > 0, \\ 0 & \text{if } d_{ij}(X) = 0. \end{cases} \quad (7)$$

If we assume, again without any real loss of generality, that

$$\frac{1}{2} \sum_{1 \leq i < j \leq n} w_{ij} \delta_{ij}^2 = 1, \quad (8)$$

then

$$\sigma(X) = 1 - \text{tr } X' B(X) X + \frac{1}{2} \text{tr } X' V X. \quad (9)$$

Matrix V is symmetric with non-positive off-diagonal elements. It is doubly-centered (rows and columns add up to zero) and thus weakly diagonally dominant. It follows that it is positive semi-definite (see Varga (1962), section 1.5). Because of irreducibility it has rank $n - 1$, and the vectors in its null space are all proportional to e , the vector with all elements equal to $+1$. For all X the matrix $B(X)$ is also symmetric, positive semi-definite, and doubly-centered.

3 SMACOF

Define the *Guttman Transform* of configuration X as

$$\Gamma(X) := V^+B(X)X \quad (10)$$

The *SMACOF algorithm* for unconstrained MDS is, with iteration counter $k \geq 1$,

$$X^{(k)} = \Gamma(X^{(k-1)}) = \Gamma^k(X^0), \quad (11)$$

where $\Gamma^0 = I$ and $\Gamma^k(X) := \Gamma(\Gamma^{k-1}(X))$.

De Leeuw (1977) shows that the SMACOF iterations (32) tend (in a specific sense described later in this paper) to a fixed point of the Guttman transform, i.e. to an X with $\Gamma(X) = X$. If stress is differentiable at X then the fixed points of the Guttman transform are stationary points, where the derivative of stress vanishes. De Leeuw (1984) shows that at a local minimum we have $d_{ij}(X) > 0$ for all (i, j) with $w_{ij}\delta_{ij} > 0$, i.e. stress is differentiable at local minima. The equation $V^+B(X)X = X$ says that the columns of X are eigenvectors of $V^+B(X)$, with eigenvalues equal to one. De Leeuw (2014) shows that if $V^+B(X)X = X$ and all eigenvalues of $V^+B(X)$ are less than or equal to one then stress has its global minimum at X .

The algorithm (without weights) was proposed by Guttman (1968) by differentiating stress and setting the derivatives equal to zero. This derivation ignores the problem that stress is not differentiable if one or more distances are zero, and it also does not say anything about convergence of the algorithm. In De Leeuw (1977) and De Leeuw and Heiser (1977) a new derivation of the algorithm was given, using a majorization argument based on the Cauchy-Schwarz inequality. This make it possible to avoid problems with differentiability, and it leads to a simple convergence proof.

4 Global Convergence of SMACOF

Following De Leeuw (1977) we define

$$\rho(X) = \sum_{1 \leq i < j \leq n} w_{ij} \delta_{ij} d_{ij}(X) = \text{tr } X' B(X) X, \quad (12)$$

$$\eta^2(X) = \sum_{1 \leq i < j \leq n} w_{ij} d_{ij}^2(X) = \text{tr } X' V X, \quad (13)$$

and

$$\lambda(X) = \frac{\rho(X)}{\eta(X)}. \quad (14)$$

The main result in De Leeuw (1977) is:

1. The SMACOF iterates $X^{(k)}$ are in the compact set $\eta(X) \leq 1$,
2. $\{\rho(X^k)\}$, $\{\eta(X^k)\}$ and $\{\lambda(X^k)\}$ are increasing sequences, with limits ρ_∞, η_∞ and λ_∞ , where $\eta_\infty = \lambda_\infty = \sqrt{\rho_\infty}$.
3. $\{\sigma(X^k)\}$ is a decreasing sequence converging to $\sigma_\infty = 1 - \eta_\infty^2$.
4. At any accumulation point X_∞ of $\{\sigma(X^k)\}$ we have $\sigma(X_\infty) = \sigma_\infty$ and $X_\infty = \Gamma(X_\infty)$.
5. $\eta(X^{(k+1)} - X^{(k)}) \rightarrow 0$ and thus either $\{\sigma(X^k)\}$ converges or the set of accumulation points of $\{\sigma(X^k)\}$ is a continuum.

So, in words, the sequence of stress values converges monotonically. The sequence of configurations is asymptotically regular and has one or more accumulation points. Each accumulation point is a fixed point of the Guttman transform. All accumulation points have the same function value, which is also the limit of the stress sequence.

But it is important to emphasize that De Leeuw (1977) did not show that the configuration sequence $\{X^{(k)}\}$ actually a Cauchy sequence, and converges to a configuration X_∞ . This is basically because of the rotational invariance of the MDS problem. If K is a rotation matrix, i.e. $K'K = KK' = I$, then $d_{ij}(XK) = d_{ij}(X)$ for all i and j , and thus $\sigma(XK) = \sigma(X)$. If X is a fixed point of the Guttman transform, then so is XK . Consequently there are no isolated fixed points, each fixed point is part of a nonlinear continuum of fixed points.

It is also of interest that De Leeuw (1984) showed that if X is a local minimizer of stress then $d_{ij}(X) > 0$ for all i and j such that $w_{ij} \delta_{ij} > 0$. Thus, if weights and dissimilarities are positive, stress is differentiable at a local minimum. De Leeuw (1993) shows that stress has only a single local maximum at $X = 0$, and consequently the only possible stationary points are local minima and saddle points. It is shown by De Leeuw (2019) that all fixed points of the Guttman transform which are not of full column rank are saddle points. Thus if X is a local minimizer of MDS, then adding q zero columns to X produces a saddle point $[X \mid 0]$

for $(p+q)$ MDS. At saddle points there are always directions in which stress can be decreased, and consequently the SMACOF algorithm with enough iterations will eventually get close to a continuum of local minima.

5 Local Convergence of SMACOF

De Leeuw (1988) studies the local convergence of SMACOF, i.e. the *rate of convergence* of the SMACOF iterations. There is, however, a problem not adequately addressed in that article, which has quite a bit of hand-waiving, and some incorrect statements. If there is no convergence to a single configuration then the usual rate of convergence is not defined either.

We know the sequence $\{\eta(X^{(k+1)} - X^{(k)})\}$ tends to zero. We can measure its rate of convergence by the *ratio factor*

$$q_k = \frac{\eta(X^{(k+1)} - X^{(k)})}{\eta(X^{(k)} - X^{(k-1)})}, \quad (15)$$

and the *root factor*

$$r_k = \eta(X^{(k+1)} - X^{(k)})^{1/k}. \quad (16)$$

There is no guarantee that either $\{q_k\}$ or $\{r_k\}$ converges, but

$$\liminf_{k \rightarrow \infty} q_k \leq \liminf_{k \rightarrow \infty} r_k \leq \limsup_{k \rightarrow \infty} r_k \leq \limsup_{k \rightarrow \infty} q_k \quad (17)$$

Thus if $\lim_{k \rightarrow \infty} q_k$ exists then $\lim_{k \rightarrow \infty} r_k$ exist and the two limit values are equal. See Rudin (1976), p. 68, theorem 3.37.

The rate of convergence of the SMACOF iterations at a fixed point X is computed in De Leeuw (1988) as the spectral norm $\kappa(X) = \|\mathcal{D}\Gamma(X)\|_\infty$, i.e. as the modulus of the largest eigenvalue of the derivative of the Guttman transform. There are good reasons to compute this quantity. From Ostrowski's Theorem (Ostrowski (1973), theorem 22.1, p. 151, Ortega and Rheinboldt (1970), theorem 10.1.3, p. 300) we know that if $\kappa(X) < 1$, then X is a *point of attraction* of the SMACOF iteration. If we start close enough then the iterations will converge to X . We also know (Ostrowski (1973), theorem 22.2, p. 152) that if $\kappa(X) > 1$ then X is a *point of repulsion*, which means the iterations will diverge when started in a certain solid angle with X at its apex.

If we define, with Ortega and Rheinboldt (1970), chapter 9, for a sequence $\{X^{(k)}\}$ converging to X , the ratio and root convergence factors

$$Q_1(\{X^{(k)}\}) = \limsup_{k \rightarrow \infty} \frac{\eta(X^{(k+1)} - X)}{\eta(X^{(k)} - X)}, \quad (18)$$

and

$$R_1(\{X^{(k)}\}) = \limsup_{k \rightarrow \infty} \eta(X^{(k)} - X)^{1/k}, \quad (19)$$

then $\kappa(X) < 1$ implies $R_1(X^{(k)}) = \kappa(X)$ (Ortega and Rheinboldt (1970), theorem 10.1.4, p. 301). Also $R_1(X^{(k)})$ is independent of the norm we have chosen, which is η in the SMACOF

case (Ortega and Rheinboldt (1970), theorem 9.9.2, p. 288). Because $Q_1(X^{(k)})$ depends on the norm we can only assert that $Q_1(X^{(k)}) \geq R_1(X^{(k)}) = \kappa(X)$, although it is guaranteed in the SMACOF case that some norm exists for which there is equality (Ortega and Rheinboldt (1970), Notes and Remarks NR 10.1-5, p. 306).

There is a problem, however, with applying the Ostrowski and Ortega-Rheinboldt results in our case. Because of the invariance of distances under rotation we have $\kappa(X) = 1$, no matter what X is. And, basically for the same reason, we have not shown that the SMACOF iterations converge to a fixed point at all. De Leeuw (1988) on page 171 acknowledges the problem. In fact, he proposes a way around the problem in the proof of theorem 3 on page 175, but the proof is not very convincing, although convincing enough to get past the reviewers. I will try to be more specific and precise.

Observe that

$$\mathcal{D}\Gamma(X) = V^+ \mathcal{D}^2 \rho(X), (\#eq : dgamma) \quad (20)$$

and

$$\mathcal{D}^2 \sigma(X) = V - \mathcal{D}^2 \rho(X) = V(I - \mathcal{D}\Gamma(X)). (\#eq : d2sigma) \quad (21)$$

Because ρ is convex $\mathcal{D}^2 \rho(X)$ is symmetric and positive semi-definite. The eigenvalues of $\mathcal{D}\Gamma(X)$ are the generalized eigenvalues of the positive semi-definite matrix pair $(\mathcal{D}^2 \rho(X), V)$.

An explicit formula for the derivatives of the Guttman transform, or equivalently for the second derivatives of stress, was first given by De Leeuw (1988). We write $\mathcal{D}\Gamma(X)[Y]$ for the derivative evaluated at Y . Partial derivatives can be obtained by choosing $Y = e_i e'_s$. Of course we assume here that $d_{ij}(X) > 0$ for all $i < j$ with $w_{ij} \delta_{ij} > 0$.

$$\mathcal{D}\Gamma(X)[Y] = V^+ \{B(X)Y - H(X, Y)X\}, (\#eq : dgamma) \quad (22)$$

with

$$H(X, Y) = \sum_{1 \leq i < j \leq n} w_{ij} \frac{\delta_{ij}}{d_{ij}(X)} \frac{\text{tr } X' A_{ij} Y}{d_{ij}^2(X)} A_{ij}. (\#eq : hdef) \quad (23)$$

These formulas, together with equations @ref(eq:dgamma) and @ref(eq:d2sigma), prove the main results in De Leeuw (1988).

1. All eigenvalues of $\mathcal{D}\Gamma(X)$ are non-negative.
2. **(Homogeneity)** X is an eigenvector of $\mathcal{D}\Gamma(X)$ with eigenvalue zero.
3. **(Translation)** $\mathcal{D}\Gamma(X)$ has at least p additional eigenvalues equal to zero, corresponding with eigenvectors of the form ee'_s , where e has all elements equal to one and e_s has one of its elements equal to one and the others equal to zero.
4. **(Rotation)** If X is a fixed point of the Guttman transform then $\mathcal{D}\Gamma(X)$ has at least $\frac{1}{2}p(p-1)$ eigenvalues equal to one, corresponding with eigenvectors of the form XA , with A anti-symmetric.

5. At a local minimum point X of stress all eigenvalues of $\mathcal{D}\Gamma(X)$ are less than or equal to one. At a strict local minimum they are strictly less than one.
6. At a saddle point X of stress the largest eigenvalue of $\mathcal{D}\Gamma(X)$ is larger than one.

6 Modified SMACOF

To avoid the problems caused by a continuum of fixed points De Leeuw (1988) proposes to rotate each SMACOF iterate to a unique position, for example to principal components. For any X with singular value decomposition $X = K\Lambda L'$ we define $\Pi(X) = K\Lambda = XL$. Note that if one or more singular values are equal then Π is not unique and becomes a point-to-set map. We can rotate within each of the spaces corresponding to multiple singular values, and thus we have not completely eliminated indeterminacy due to rotation. In the sequel we simply assume that the p singular values of X are different.

Now consider the iterations

$$\tilde{X}^{(k+1)} = \Pi(\Gamma(\tilde{X}^{(k)})) = (\Pi\Gamma)^k(\tilde{X}^0). \quad (24)$$

Let's call this modified SMACOF, or mSMACOF for short.

It looks initially as if mSMACOF require a great deal of extra computation, but this is actually not the case. If M is a rotation matrix we have $\Pi(XM) = \Pi(X)$ and $\Gamma(XM) = \Gamma(X)M$. This implies that $\Pi\Gamma\Pi(X) = \Pi\Gamma(X)$ for all X . As a result the sequences $\{X^{(k)}\}$ and $\{\tilde{X}^{(k)}\}$ have a simple relationship. If we start the $\{\tilde{X}^{(k)}\}$ sequence with X^0 then for each k we have $\sigma(\tilde{X}^{(k)}) = \sigma(X^{(k)})$ and $\tilde{X}^{(k)} = \Pi(X^{(k)})$. Thus the two sequences of stress values are exactly the same for SMACOF and mSMACOF and the two configurations, and consequently also the accumulation points of the two sequences of configurations, just differ by a rotation. To get $\tilde{X}^{(k)}$ we can just compute the SMACOF iterate $X^{(k)}$ and rotate it to principal components.

The derivative of $\Pi\Gamma$ is, by the chain rule,

$$\mathcal{D}\Pi\Gamma(X) = \mathcal{D}\Pi(\Gamma(X))\mathcal{D}\Gamma(X). \quad (25)$$

After some computation we find

$$\mathcal{D}\Pi(X)[Y] = YL + K\Lambda M, \quad (26)$$

where $X = K\Lambda L'$ and M is the anti-symmetric matrix with off-diagonal elements

$$m_{ij} = -\frac{\lambda_i u_{ij} + \lambda_j u_{ji}}{\lambda_i^2 - \lambda_j^2} \quad (27)$$

with $U = K'YL$.

If X satisfies $\Pi(X) = X$ then $L = I$ and thus $\mathcal{D}\Pi(X)[Y] = Y + XM$ and $U = K'Y$. We proceed to compute the eigenvectors and eigenvalues of the Jacobian at an X with $\Pi(X) = X$.

1. If $Y = XA$ with A antisymmetric then $M = -L'AL$ and $\mathcal{D}\Pi(X)[Y] = 0$. This corresponds with $\frac{1}{2}p(p-1)$ zero eigenvalues
2. If $Y = K\Lambda^{-1}A$ with A antisymmetric then $M = 0$. This gives $\frac{1}{2}p(p-1)$ eigenvectors with eigenvalue one.
3. If $Y = K\Lambda^{-1}D$ with D diagonal then $M = 0$. This gives another p eigenvectors with eigenvalue one.
4. If $Y = K_{\perp}S$ with $K_{\perp}$ a basis for the orthogonal complement of X then $M = 0$ and we have another $p(n-p)$ eigenvalues equal to one.

This is a complete set of eigenvectors. It turns out all eigenvalues are equal to one, except for $\frac{1}{2}p(p-1)$ zero eigenvalues. It follows that fixed points of $\Pi\Gamma$ are of the form $\tilde{X} = XL$, with X a fixed point of Γ and L the rotation of X to principal components). Thus $\Gamma(\tilde{X}) = \Gamma(X)L = XL = \tilde{X}$, and $\tilde{X}$ is a fixed point of both Γ and $\Pi\Gamma$. The eigenvalues of $\mathcal{D}\Pi\Gamma(\tilde{X})$ are the same as the eigenvalues of $\mathcal{D}\Gamma(X)$, except for the $\frac{1}{2}p(p-1)$ eigenvalues equal to one, corresponding with XA for antisymmetric A , which are replaced by zero eigenvalues. The Ostrowski and Ortega-Rheinboldt results now apply directly to $\Pi\Gamma$. If there are no zero distances, if the singular values of X are different, and if the largest eigenvalue of $\mathcal{D}\Pi\Gamma(X)$ is less than one, then X is a point of attraction, the SMACOF iterations converge to $\tilde{X}$ if started sufficiently close to it (in the *attraction ball* of X), and $R_1(X)$ is equal to the largest eigenvalue.

7 Coefficient Space

Also define

$$\{H_{ij}\}_{st} := \text{tr } Y_s' A_{ij} Y_t$$

It is of some interest to look at the special case in which $Y_s = e_k e_p'$ and $Y_t = e_l e_q'$. Then

$$\{H_{ij}\}_{st} = \text{tr } e_p e_k' (e_i - e_j)(e_i - e_j)' e_l e_q' = \delta^{pq}(\delta^{ik} - \delta^{jk})(\delta^{il} - \delta^{jl})$$

It is of some interest to look at the special case in which $Y_s = e_k e_p'$ and $Y_t = e_l e_q'$. Then

$$\{H_{ij}\}_{st} = \text{tr } e_p e_k' (e_i - e_j)(e_i - e_j)' e_l e_q' = \delta^{pq}(\delta^{ik} - \delta^{jk})(\delta^{il} - \delta^{jl})$$

and thus

$$v_{st} = \delta^{pq} \sum_{1 \leq i < j \leq n} w_{ij} (\delta^{ik} - \delta^{jk}) (\delta^{il} - \delta^{jl})$$

For the notation in coefficient space we need the square, symmetric, positive semi-definite matrices A_{ij} of order m with elements

$$\{A_{ij}\}_{st} := \text{tr } Y'_s A_{ij} Y_t. \quad (28)$$

The squared distance equals

$$d_{ij}^2(\alpha) := \alpha' A_{ij} \alpha. \quad (29)$$

Note that we continue to overload symbols such as A_{ij} and d . The same is true for V from (5), which we define in coefficient space by using definition (28), and for $\tilde{B}(\alpha)$ from (6). In coefficient space

$$\sigma(\alpha) := 1 - \alpha' B(\alpha) \alpha + \frac{1}{2} \alpha' V \alpha. \quad (30)$$

and the MDS problem is to minimize (30) over $\alpha \in \mathbb{R}^m$.

In coefficient space the Guttman Transform is

$$\Gamma(\alpha) := V^+ B(\alpha) \alpha \quad (31)$$

The *SMACOF algorithm* in configuration space is

$$\alpha^{(k)} = \Gamma(\alpha^{(k-1)}) = \Gamma^k(\alpha^0). \quad (32)$$

8 Software

The appendix has an ad-hoc version of the `smacof` function. Its arguments are

```
args(smacof)
```

```
function (delta, w, p = 2, xold = torgerson(delta, p), pca = FALSE,
        verbose = FALSE, eps = 1e-15, itmax = 10000)
NULL
```

The function expects both dissimilarities and weights to be of class `dist`. The default initial configuration uses classical MDS (Torgerson (1958)). We iterate until either the distance $\eta(X^{(k)} - X^{(k-1)})$ between successive configurations is less than `eps`, which has the ridiculously small default value of 10^{-15} , or until the maximum number of iterations `itmax` is reached, which defaults to the ridiculously large default value of 10000. If `pca` is `TRUE` all iterates are rotated to principal components. If `verbose` is `TRUE` we print, for each iteration, the iteration number k , the stress $\sigma(X^{(k)})$, the change $\eta(X^{(k)} - X^{(k-1)})$, and the root and ratio convergence factors.

The second file `derivative.R` computes the Jacobian of the iteration functions at the limit (i.e. at the point where SMACOF stops). There are actually six functions in the file. The three functions `dGammaA`, `dPiA` and `dPiGammaA` use the analytical expressions we have derived earlier for $\mathcal{D}\Gamma(X)$, $\mathcal{D}\Pi(X)$ and $\mathcal{D}\Pi\Gamma(X)$. The corresponding functions `dGammaN`, `dPiN` and `dPiGammaN` numerically compute the Jacobian using the `jacobian` function from the `numDeriv` package (Gilbert and Varadhan (2019)). They are basically used to check the formulas.

9 Examples

9.1 Ekman

Time for a numerical example. We will use the color similarity data of Ekman (1954), taken from the SMACOF package (De Leeuw and Mair (2009)). We transform Ekman's similarities s_{ij} to dissimilarities δ_{ij} using $\delta_{ij} = (1 - s_{ij})^3$, and we use unit weights in W .

SMACOF requires 52 iterations for convergence. The minimum of stress is 0.0110248119, the root factor is equal to 0.5085800483 and the ratio factor is 0.5329246268. The eigenvalues of $\mathcal{D}\Gamma(X)$ are

```
0.999999999999999
0.538510668196408
0.532498554224166
0.529669334191043
0.525541097754551
0.519602718218807
0.516533687841054
0.513727908691608
0.510295310409166
0.506357695934493
```

```

0.495649713446001
0.481518780073304
0.469548625536451
0.445038589020916
0.410479252654484
0.394284315478173
0.357670750114585
0.348395413045201
0.330341477528788
0.313520384427863
0.287598015841956
0.280399104121623
0.267278474596759
0.247631495462991
0.216576009083047
0.000000000000000
0.000000000000000
0.000000000000000

```

As an aside, the Ekman example (with this particular transformation of the similarities) is special, because the two-dimensional solution is actually the global minimum over all configurations (i.e. the global minimum of MDS for all $1 \leq p \leq n$). As shown in De Leeuw (2014) this follows from the fact that the two unit eigenvalues of $V^+B(X)$ are actually its two largest eigenvalues, and consequently X is the solution of the convex full-dimensional nMDS relaxation of the MDS problem (De Leeuw et al. (2016)). The eigenvalues of $V^+B(X)$ are

```

1.000000000000000
1.000000000000000
0.923497086367287
0.907901212970889
0.862936584878895
0.852692003103344
0.829803620857357
0.814556167662381
0.793238576338502
0.791651722456649
0.786442678118770
0.747679475705025
0.728268247434340
0.000000000000001

```

We now set `pca=TRUE` and run `mSMACOF`. It requires 51 iterations for convergence. The minimum of stress is 0.0110248119, and the root factor is equal to 0.5080118475 and the ratio factor is 0.5718672565. The eigenvalues of $\mathcal{D}\Pi\Gamma(X)$ at the fixed point X are

```
0.538510668196407
0.532498554224166
0.529669334191045
0.525541097754552
0.519602718218807
0.516533687841055
0.513727908691609
0.510295310409168
0.506357695934493
0.495649713446002
0.481518780073306
0.469548625536451
0.445038589020917
0.410479252654486
0.394284315478174
0.357670750114585
0.348395413045201
0.330341477528788
0.313520384427864
0.287598015841957
0.280399104121623
0.267278474596758
0.247631495462991
0.216576009083047
0.000000000000000
0.000000000000000
0.000000000000000
0.000000000000000
```

They are equal to the eigenvalues of $\mathcal{D}\Gamma(X)$ and the root convergence factor $R_1(\{X^{(k)}\})$ is 0.5385106682.

9.2 De Gruijter

The second example are dissimilarities between nine Dutch political parties, collected a long time ago by De Gruijter ([1967](#)).

	KVP	PvdA	VVD	ARP	CHU	CPN	PSP	BP
PvdA	2.63							
VVD	2.27	3.72						
ARP	1.60	2.64	2.46					
CHU	1.80	3.22	1.97	0.20				
CPN	4.54	2.12	5.13	4.84	4.80			
PSP	3.73	1.59	4.55	3.73	4.08	1.08		
BP	4.18	4.22	3.90	4.28	3.96	3.34	3.88	
D66	3.17	2.47	1.67	3.13	3.04	4.42	3.36	4.36

We analyze the data with SMACOF using three dimensions.

SMACOF requires 779 iterations for convergence. The minimum of stress is 0.003442194, the root factor is equal to 0.9565946372, and the ratio factor is 0.9684304863. The eigenvalues of $\mathcal{D}\Gamma(X)$ are

```

1.0000000000000001
1.0000000000000000
1.0000000000000000
0.965505429805660
0.940592046981168
0.919047686446446
0.863993920924431
0.822263696226349
0.810020971414307
0.771947328045071
0.728539213663602
0.712621684571535
0.659318624263222
0.638485365688918
0.624552464148482
0.616085432872672
0.557480497708779
0.501954186928526
0.458630667850014
0.450598367846591
0.355243400669833
0.309186479174062
0.247708397109091
0.0000000000000002
0.0000000000000002

```

0.0000000000000000
0.0000000000000000

In this case there is no guarantee that we have the three-dimensional global minimum. The unit eigenvalues of the matrix $V^+B(X)$ are not the three largest ones.

1.079524009371955
1.032606649163671
1.0000000000000000
1.0000000000000000
1.0000000000000000
0.986706272372898
0.971839080877692
0.906211919383164
0.0000000000000001

mSMACOF in three dimensions requires 778 iterations for convergence. The minimum of stress is 0.003442194, the root factor is equal to 0.9565709099, and the ratio factor is 0.9714170285. The eigenvalues of $\mathcal{D}\Pi(X)$ are

0.965505429805661
0.940592046981168
0.919047686446446
0.863993920924433
0.822263696226351
0.810020971414307
0.771947328045072
0.728539213663603
0.712621684571534
0.659318624263222
0.638485365688918
0.624552464148483
0.616085432872671
0.557480497708778
0.501954186928526
0.458630667850015
0.450598367846592
0.355243400669831
0.309186479174061

0.247708397109091
0.0000000000000001
0.0000000000000001
0.0000000000000001
0.0000000000000000
0.0000000000000000
0.0000000000000000
0.0000000000000000
0.0000000000000000

Again, as expected, the eigenvalues for SMACOF and mSMACOF are the same and the root convergence factor is the spectral norm of the mSMACOF derivative, which is 0.9655054298.

10 Conclusion

From a practical point of view there is no need to ever use modified SMACOF. We only use the Guttman transform, and compute the largest eigenvalue of its derivative at the solution X , ignoring the $\frac{1}{2}p(p-1)$ trivial unit eigenvalues. That largest eigenvalue, if it is strictly smaller than one, is the root convergence factor. In that case SMACOF can be said to converge to $\Pi(X)$, which is the SMACOF solution rotated to principal components.

References

- De Gruijter, Dato N. M. 1967. *The Cognitive Structure of Dutch Political Parties in 1966*. Report Nos. E019-67. Psychological Institute, University of Leiden.
- De Leeuw, Jan. 1977. “Applications of Convex Analysis to Multidimensional Scaling.” In *Recent Developments in Statistics*, edited by J. R. Barra, F. Brodeau, G. Romier, and B. Van Cutsem. North Holland Publishing Company.
- De Leeuw, Jan. 1984. “Differentiability of Kruskal’s Stress at a Local Minimum.” *Psychometrika* 49: 111–13.
- De Leeuw, Jan. 1988. “Convergence of the Majorization Method for Multidimensional Scaling.” *Journal of Classification* 5: 163–80.
- De Leeuw, Jan. 1993. *Fitting Distances by Least Squares*. Preprint Series No. 130. UCLA Department of Statistics. <https://jansweb.netlify.app/publication/deleeuw-r-93-c/deleeuw-r-93-c.pdf>.
- De Leeuw, Jan. 2014. “Bounding, and Sometimes Finding, the Global Minimum in Multidimensional Scaling.” UCLA Department of Statistics. <https://jansweb.netlify.app/publication/deleeuw-u-14-b/deleeuw-u-14-b.pdf>.
- De Leeuw, Jan. 2019. “Fitting Distances by Least Squares.” <https://jansweb.netlify.app/publication/deleeuw-r-93-c/deleeuw-r-93-c.pdf>.
- De Leeuw, Jan. 2023. “Least Squares Unidimensional Scaling.” <https://doi.org/10.13140/RG.2.2.27920.17927>.
- De Leeuw, Jan, Patrick J. F. Groenen, and Patrick Mair. 2016. “Full-Dimensional Scaling.” <https://jansweb.netlify.app/publication/deleeuw-groenen-mair-e-16-e/deleeuw-groenen-mair-e-16-e.pdf>.
- De Leeuw, Jan, and Willem J. Heiser. 1977. “Convergence of Correction Matrix Algorithms for Multidimensional Scaling.” Chap. 32 in *Geometric Representations of Relational Data*, edited by J. C. Lingoes. Mathesis Press.
- De Leeuw, Jan, and Patrick Mair. 2009. “Multidimensional Scaling Using Majorization: SMACOF in R.” *Journal of Statistical Software* 31 (3): 1–30. <https://www.jstatsoft.org/article/view/v031i03>.

- Ekman, Gosta. 1954. "Dimensions of Color Vision." *Journal of Psychology* 38: 467–74.
- Gilbert, Paul, and Ravi Varadhan. 2019. *numDeriv: Accurate Numerical Derivatives*. <https://CRAN.R-project.org/package=numDeriv>.
- Guttman, Louis. 1968. "A General Nonmetric Technique for Fitting the Smallest Coordinate Space for a Configuration of Points." *Psychometrika* 33: 469–506.
- Kruskal, Joseph B. 1964a. "Multidimensional Scaling by Optimizing Goodness of Fit to a Nonmetric Hypothesis." *Psychometrika* 29: 1–27.
- Kruskal, Joseph B. 1964b. "Nonmetric Multidimensional Scaling: a Numerical Method." *Psychometrika* 29: 115–29.
- Ortega, James M., and Werner C. Rheinboldt. 1970. *Iterative Solution of Nonlinear Equations in Several Variables*. Academic Press.
- Ostrowski, Alexander M. 1973. *Solution of Equations in Euclidean and Banach Spaces*. Third Edition of Solution of Equations and Systems of Equations. Academic Press.
- Rudin, Walter. 1976. *Principles of Mathematical Analysis*. Third Edition. McGraw-Hill.
- Torgerson, Warren S. 1958. *Theory and Methods of Scaling*. Wiley.
- Varga, Richard S. 1962. *Matrix Iterative Analysis*. Prentice Hall.