

Voronoi Homogeneity Analysis of Categorical Data

Jan de Leeuw

May 10, 2026

TBD

Table of contents

1	Introduction	3
1.1	Homogeneity Analysis	3
1.2	Voronoi Analysis	5
2	Algorithm	7
2.1	Monotone Regression	8
2.2	Majorization	9
3	Examples	10
	References	11

Note: This is a working manuscript which will be expanded/updated frequently. All suggestions for improvement are welcome. All Rmd, tex, html, pdf, R, and C files are in the public domain. Attribution will be appreciated, but is not required. The files can be found at <https://github.com/deleeuw/voronoi>

1 Introduction

1.1 Homogeneity Analysis

Homogeneity Analysis, a.k.a. *Multiple Correspondence Analysis*, can be introduced in many different ways. We refer to Guttman (1941), Tenenhaus and Young (1985), Bekker and De Leeuw (1988), Gifi (1990), Greenacre and Blasius (2006), Le Roux and Rouanet (2010), and De Leeuw (2023) for a bouquet of discussions and interpretations of the equations and algorithms defining the technique.

The data are measurements of n objects on m categorical variables. Variable j has k_j categories, which are the different unique values the variable assumes for the n objects. We also say that a category has a number of objects that “belong to” the category.

Here is a small example with $n = 10$ and $m = 3$, taken from Gifi (1990), page 65.

	[, 1]	[, 2]	[, 3]
[1,]	"a"	"p"	"u"
[2,]	"b"	"q"	"v"
[3,]	"a"	"r"	"v"
[4,]	"a"	"p"	"u"
[5,]	"b"	"p"	"v"
[6,]	"c"	"p"	"v"
[7,]	"a"	"p"	"u"
[8,]	"a"	"p"	"v"
[9,]	"c"	"p"	"v"
[10,]	"a"	"p"	"v"

Suppose the objects are mapped into n points of Euclidean space $\mathbb{R}^p$. For each variable we can construct an additional k_j points in the same space by computing for each category the centroids of all object-points that belong to it. Thus each variable defines its own plot. We then draw lines from the n object points to the category point they belong to and we obtain k_j “stars”. Homogeneity Analysis maps the objects into space in such a way that the stars are as small as possible, where the size of the star is measured as the sum of squares of the distances between each category points and the object points of the oibjects that belong to the category (summed over categories and over variables). In order to avoid the trivial solution in which all objects and categories are mapped into the same point homogeneity analysis requires that the *configuration* of object points (an $n \times p$ matrix) is column-centered and orthonormal.


There are a huge number of variations (De Leeuw (2004)) possible on the same theme. Instead of the centroids we could use the Weber points to define our stars. Instead of Euclidean distance

we could use any other distance on $\mathbb{R}^p$. Instead of the stars we could use the convex hulls or the minimum spanning trees, or any other way to visualize and quantify the compactness of the point clouds of each category. Any one of these choices defines a technique that could be called homogeneity analysis, but in this paper we mean the classical choice that uses stars, squared Euclidean distance, and orthonormal object points.

The key mathematical object in homogeneity analysis is the *indicator matrix*, which is a classical way to code categorical data¹. For a variable j with k_j categories it is an $n \times k_j$ binary matrix G_j in which each row i contains a single element equal to one, indicating into which category the object i maps for variable j . The cross product $D_j := G_j' G_j$ is diagonal, with the *marginal frequencies* of the categories on the diagonal. For two different variables the crossproduct $C_{j\ell} = G_j' G_\ell$ is the $k_j \times k_\ell$ *cross table* or *contingency table* of the two variables. Here are the three indicator matrices of our small example, next to each other.

	a	b	c		p	q	r		u	v
[1,]	1	0	0	1	0	0	1	1	0	1
[2,]	0	1	0	0	1	0	0	0	1	1
[3,]	1	0	0	0	0	1	0	0	1	1
[4,]	1	0	0	1	0	0	0	1	0	1
[5,]	0	1	0	1	0	0	0	0	1	1
[6,]	0	0	1	1	0	0	0	0	1	1
[7,]	1	0	0	1	0	0	0	1	0	1
[8,]	1	0	0	1	0	0	0	0	1	1
[9,]	0	0	1	1	0	0	0	0	1	1
[10,]	1	0	0	1	0	0	0	0	1	1

A Homogeneity Analysis in two dimensions produces the following three star plots. Objects are numbered and in BLUE, categories have bigger labels and are in RED. The analysis minimizes the sum of squares of the length of all black lines in the three plots, under the condition that the matrix of object scores is column-centered and orthonormal. We can think of the results as a form of *graph drawing*, where the graph is the union of the m bipartite graphs of the relation “belonging to” (Michailidis and De Leeuw (2001)).



¹Since the indicator matrix is the core of the Gifi system we avoid the irreverent alternative “dummy coding”.

We now deviate from the usual Gifi and French approaches (Husson et al. (2016)) and define homogeneity analysis as a form of non-metric multidimensional scaling, using the *stress* loss function (Kruskal (1964a), Kruskal (1964b)). Define²

$$\sigma(X, Y_1, \dots, Y_m, \Delta_1, \dots, \Delta_m) := \sum_{j=1}^m \sum_{i=1}^n \sum_{\ell=1}^{k_j} (\delta_{i\ell}^j - d(x_i, y_\ell^j))^2, \quad (1)$$

with $d(., .)$ Euclidean distance. Homogeneity analysis minimizes this loss over all configurations X with $X'X = I$, over all Y_j , and over all Δ_j that satisfy $\delta_{i\ell}^j = 0$ whenever $g_{i\ell}^j = 1$. All other elements of Δ are only constrained to be non-negative.

If

$$\sigma_\star(X, Y_1, \dots, Y_m) := \min_{\Delta_1, \dots, \Delta_m} \sigma(X, Y_1, \dots, Y_m, \Delta_1, \dots, \Delta_m), \quad (2)$$

then

$$\sigma_\star(X, Y_1, \dots, Y_m) = \sum_{j=1}^m \sum_{i=1}^n \sum_{\ell=1}^{k_j} g_{i\ell}^j d^2(x_i, y_\ell^j). \quad (3)$$

Next define

$$\sigma_{\star\star}(X) := \min_{Y_1, \dots, Y_m} \sigma_\star(X; Y_1, \dots, Y_m). \quad (4)$$

The minimum over the Y_j is attained at the centroids $\hat{Y}_j := D_j^{-1} G_j' X$, and if we substitute these optimal $\hat{Y}_j$ in (3) we see that $\sigma_{\star\star}(X)$ is indeed equal to the sum of squares of all stars. Minimizing $\sigma_{\star\star}(X)$ over $X'X = I$, and thus minimizing (1) over X , the Y_j , and the Δ_j , defines homogeneity analysis. Computing the solution turns out to be a straightforward eigenvalue-eigenvector problem. The details are in the publications we referenced at the beginning of this section.

1.2 Voronoi Analysis

Although using loss function (1) makes homogeneity analysis a form of non-metric multidimensional scaling there are some important differences with more familiar applications of minimizing stress. There are no local minima in homogeneity analysis, because it turns out to be a symmetric eigenvalue problem. Also, and perhaps most importantly, we do not really expect the minimum loss function value to be zero, or even close to zero. The loss functions measures deviations from a very unrealistic model. Zero loss would imply all object points coincide with all category points they belong to. This can only be realized, except in trivial cases, if all object and category points map into a single point, which is prohibited by the orthonormality constraint on the object scores. Thus we get a perfectly determinate and stable

²The symbol $:=$ is used for definitions

solution, at low computational cost, but by imposing unrealistic constraints and a rather arbitrary normalization. We can get around this seemingly unfortunate situation by admitting we are not fitting or testing a “model”, but we are making drawings of the bipartite graphs that constitute the data.

The lack of a realistic Model makes it interesting to generalize loss function (1) to

$$\sigma(X, Y_1, \dots, Y_m, \Delta_1, \dots, \Delta_m) := \sum_{j=1}^m \sum_{i=1}^n \sum_{\ell=1}^{k_j} (\delta_{i\ell}^j - d(x_i, y_\ell^j))^2, \quad (5)$$

with the constraints that if $g_{i\ell}^j = 1$ then merely $\delta_{i\ell}^j \leq \delta_{i\nu}^j$ for all $\nu = 1, \dots, k_j$. This is significantly weaker than the homogeneity analysis constraints, which require the $\delta_{i\ell}^j$ to be zero. If object i belongs to category ℓ of variable j we do not require the object point to coincide with the category point, but we merely require that the object point is at least as close to category point ℓ as to any of the other $k_j - 1$ category points of that variable.

As a consequence, the plots corresponding with loss function (5) are also different. As in homogeneity analysis there is a plot for each variable j with both n object points and k_j category points. The k_j category points define k_j *Voronoi regions* that partition the space. Voronoi region $\mathcal{V}_\ell^j$ is the region of space where the points are at least as close to category point y_ℓ^j as to any of the other category points of variable j . And loss (4) is zero (for a variable) if all object points are in the “correct” Voronoi region, i.e. in the Voronoi region of the category point corresponding with the category they belong to.

One (very inefficient, but conceptually easy) way of constructing Voronoi regions is to draw the perpendicular bisectors of the (hypothetical) lines connecting all pairs of category points of a variable. The convex polyhedral regions thus formed, some bounded and some unbounded, define the Voronoi regions. The literature on properties and computation on Voronoi regions is large, and we refer to the textbooks of Okabe et al. (2000) and Aurenhammer et al. (2013).

Since the category point itself is obviously in its own Voronoi region and since Voronoi regions are convex it follows that if we make stars by connecting category points with object points as in homogeneity analysis then we actually require each star to be in its Voronoi region. It is no longer true that a category point is the centroid of points of the objects belonging to the category. The category point can even be outside the convex hull of “its” object points.

Voronoi Homogeneity Analysis of categorical data, as defined here, minimizes loss function (5) over X , over the Y_j and over the Δ_j satisfying the ordinal constraints. It is consequently another form of non-metric (ordinal) multidimensional scaling. To be more precise, we have m linked non-metric multidimensional unfolding problems, which are linked because they share the same X . The data are binary, and we use Kruskal’s primary approach to ties (De Leeuw (1977b)).

The constraint $X'X = I$ from homogeneity analysis is no longer needed, but some form of normalization is still necessary. We still need to exclude the trivial solution in which all points and regions collapse into a single point and all Δ_j are zero. But, more urgently, we have to deal with the trivial solutions familiar from multidimensional unfolding (Heiser (1981), De Leeuw (1983), Busing (2010)). For example, we can collapse all category points into a single point and put all object points on a sphere with the category point at its center. Then all distances between category and object points are the same and we can choose all dissimilarities equal to that distance value and obtain stress zero. It does not help to require $X'X = I$, because we can easily construct orthogonal configurations with all points on the sphere. Stronger normalization constraints are required.

In this paper we use the constraint

$$\sum_{\ell=1}^{k_j} (\delta_{i\ell}^j - \bar{\delta}_{ij})^2 = c_j, \quad (6)$$

with $\bar{\delta}_{ij}$ the mean of the $\delta_{i\ell}^j$ and with c_j constants. For instance we could choose $c_j = 1$ or $c_j = k_j$ for all j . In addition, of course, the $\delta_{i\ell}^j$ must satisfy the ordinal constraints. Constraint (6) explicitly excludes the possibility that the $\delta_{i\ell}^j$ for given i and j are equal for all ℓ .

2 Algorithm

We use an *Alternating Least Squares* or *ALS* algorithm, similar to many other non-metric multidimensional scaling methods, to minimize loss (5). For context, see Borg and Groenen (2005). We alternate minimization over Δ for given X and Y with minimization over X and Y for given Δ . Using superscript (μ) for the iteration index we can write

$$\Delta^{(\mu)} = \underset{\Delta}{\operatorname{argmin}} \sigma(X^{(\mu)}, Y^{(\mu)}, \Delta), \quad (7)$$

$$(X^{(\mu+1)}, Y^{(\mu+1)}) = \underset{X, Y}{\operatorname{argmin}} \sigma(X, Y, \Delta^{(\mu)}). \quad (8)$$

The first substep of iteration μ is a form of monotone regression for a very simple partial order, in which one of k_j elements must be smaller than the other elements. The second substep is a metric multidimensional scaling step. It cannot be carried out completely, because it would need an infinite iterative process by itself. Thus the second substep consist of a finite number of *inner iterations* within the second substep of an *outer iteration*. For the inner iterations we use standard smacof majorization steps (De Leeuw (1977a)).

2.1 Monotone Regression

The nm monotone regressions are applied to each row i of each variable j . Forgetting about all these indices for a moment we need to minimize the sum of squares $\|\delta_\ell - d_\ell\|^2$ over δ , where one of the elements, say δ_ν , must be smaller than or equal to all others, and where $\|\delta_\ell - \bar{\delta}\|^2 = c$.

Consider the cone of all vectors x in $\mathbb{R}^n$ for which $x_i \leq x_1$ for all $i \neq 1$ and $e'x = 0$. The extreme rays of the cone will be the $n - 1$ vectors of the form $\alpha e_i + \beta(e - e_i)$ with $i \neq 1$. Thus element i is equal to α and all other elements (including x_1) are equal to β . Thus we must have $\alpha > \beta$. The sum of the elements is $\alpha + (n - 1)\beta$, which must be zero. Thus $\alpha = -(n - 1)\beta$, which implies $\beta < 0$ and $\alpha > 0$. The extreme rays are positive multiples of the vectors $e_i - n^{-1}e$ with $i \neq 1$ and the cone is the set of all weighted sums with non-negative weights of these $n - 1$ vectors. Here is a small example.

```
a <- -cbind(1, -diag(4))
b <- (diag(5)-(1 / 5))[, -1]
print(a)
```

	[,1]	[,2]	[,3]	[,4]	[,5]
[1,]	-1	1	0	0	0
[2,]	-1	0	1	0	0
[3,]	-1	0	0	1	0
[4,]	-1	0	0	0	1

```
print(b)
```

	[,1]	[,2]	[,3]	[,4]
[1,]	-0.2	-0.2	-0.2	-0.2
[2,]	0.8	-0.2	-0.2	-0.2
[3,]	-0.2	0.8	-0.2	-0.2
[4,]	-0.2	-0.2	0.8	-0.2
[5,]	-0.2	-0.2	-0.2	0.8

```
print(a %*% b)
```

	[,1]	[,2]	[,3]	[,4]
[1,]	1	0	0	0
[2,]	0	1	0	0
[3,]	0	0	1	0
[4,]	0	0	0	1

For later reference the sum of squares of the vectors $e_i - n^{-1}e$ is $(n-1)/n$, so the vectors $\sqrt{n/(n-1)}(e_i - n^{-1}e)$ are the vectors of length 1 on the extreme rays.

Suppose y is a vector with n elements. The problem we study is to minimize the sum of squares $\sigma(x) := \|x - y\|^2$ over $x \in K$ and $x'Jx = 1$. Here

- K is the cone of vectors with $x_1 \leq x_2 \leq \dots \leq x_n$,
- J is the centering matrix $J := I - n^{-1}ee'$, where
- e is the vector with all n elements equal to one.

Change variables to $\tilde{x} = Jx$. Then $x'Jx = \tilde{x}'\tilde{x}$ and $x \in K$ if and only if $\tilde{x} \in K$. Also $\tilde{x} = Jx$ if and only if $x = \tilde{x} + \beta e$. So the transformed problem is to minimize $\|\tilde{x} + \beta e - y\|^2$ over β and $\tilde{x} \in K$, with $e'\tilde{x} = 0$ and $\tilde{x}'\tilde{x} = 1$. The minimizer for β is $\bar{y} := n^{-1}e'y$, and thus the problem becomes minimizing $\|\tilde{x} - Jy\|^2$ over $\tilde{x} \in K$, $\tilde{x}'\tilde{x} = 1$, and $e'\tilde{x} = 0$.

Forget about the constraint $e'\tilde{x} = 0$. If M is the monotone regression operator it follows that the solution for $\tilde{x}$ is $M(Jy)/\|M(Jy)\|$, which automatically satisfies $e'\tilde{x} = 0$. Thus the solution of our original problem is

$$\hat{x} = \frac{M(y - \bar{y})}{\|M(y - \bar{y})\|} + \bar{y}.$$

$M(y - \bar{y}) = M(y) - \bar{y}e$ and thus $\|M(y - \bar{y})\|^2 = \|M(y)\|^2 + n\bar{y}^2 - 2\bar{y}e'M(y)$ nor $e'M(y) = e'y = n\bar{y}$. Thus $\|M(y - \bar{y})\|^2 = \|M(y)\|^2 - n\bar{M}(y)^2$

$$\hat{x} = \frac{M(y) - \bar{y}e}{\|M(y - \bar{y})\|} + \bar{y}e = \frac{1}{\|M(y - \bar{y})\|} M(y) + \left\{ 1 - \frac{1}{\|M(y - \bar{y})\|} \right\} \bar{y}e$$

2.2 Majorization

$$A_{ij} := \begin{bmatrix} e_i e'_i & -e_i e'_j \\ -e_j e'_i & e_j e'_j \end{bmatrix}$$

$$Z := \begin{bmatrix} X \\ Y \end{bmatrix}$$

$$d_{ij}(X, Y) = \text{tr } Z' A_{ij} Z$$

$$V := \sum_{i=1}^n \sum_{j=1}^m A_{ij} = \begin{bmatrix} mI & -ee' \\ -ee' & nI \end{bmatrix}$$

If we define

$$V_1 := (n + m)^{-1} \begin{bmatrix} \frac{m}{n}E & -E \\ -E & \frac{n}{m}E \end{bmatrix}, \quad (9)$$

$$V_2 := \begin{bmatrix} J_n & 0 \\ 0 & 0 \end{bmatrix}, \quad (10)$$

$$V_3 := \begin{bmatrix} 0 & 0 \\ 0 & J_m \end{bmatrix}, \quad (11)$$

then

$$V = (n + m)V_1 + mV_2 + nV_3$$

Matrices V_1 , V_2 , and V_3 are symmetric projectors on orthogonal subspaces. It follows that V^+ , the Moore-Penrose inverse³ of V , is

$$V^+ = \frac{1}{n + m}V_1 + \frac{1}{m}V_2 + \frac{1}{n}V_3$$

$$B(X, Y) := \begin{bmatrix} B_{11}(X, Y) & B_{12}(X, Y) \\ B_{21}(X, Y) & B_{22}(X, Y) \end{bmatrix}$$

Software

3 Examples

References

- Aurenhammer, Franz, Rolf Klein, and Der-Tsai Lee. 2013. *Voronoi Diagrams and Delaunay Triangulations*. World Scientific Publishing Co.
- Bekker, Paul, and Jan De Leeuw. 1988. "Relation Between Variants of Nonlinear Principal Component Analysis." Chap. 1 in *Component and Correspondence Analysis*, edited by Jan L. A. Van Rijkevorsel and Jan De Leeuw. Wiley Series in Probability and Mathematical Statistics. Wiley.
- Borg, Ingwer, and Patrick J. F. Groenen. 2005. *Modern Multidimensional Scaling*. Second Edition. Springer.
- Busing, Frank M. T. A. 2010. "Advances in Multidimensional Unfolding." PhD thesis, Leiden University. <https://scholarlypublications.universiteitleiden.nl/access/item%3A2837800/view>.
- De Leeuw, Jan. 1977a. "Applications of Convex Analysis to Multidimensional Scaling." In *Recent Developments in Statistics*, edited by J. R. Barra, F. Brodeau, G. Romier, and B. Van Cutsem. North Holland Publishing Company.
- De Leeuw, Jan. 1977b. "Correctness of Kruskal's Algorithms for Monotone Regression with Ties." *Psychometrika* 42: 141–44.
- De Leeuw, Jan. 1983. *On Degenerate Multidimensional Unfolding Solutions*. Research Report. Department of Data Theory FSW/RUL. <https://jansweb.netlify.app/publication/deleeuw-r-06-a/deleeuw-r-06-a.pdf>.
- De Leeuw, Jan. 2004. *Homogeneity Analysis of Pavings*. Preprint Series No. 389. UCLA Department of Statistics. <https://jansweb.netlify.app/publication/deleeuw-r-04-e/deleeuw-r-04-e.pdf>.
- De Leeuw, Jan. 2023. "Deconstructing Multiple Correspondence Analysis." Chap. 22 in *Analysis of Categorical Data from Historical Perspectives. Essays in Honour of Shizuhiko Nishisato*, edited by Eric J. Beh, Rosaria Lombardo, and Jose G. Clavel. Springer.
- Gifi, Albert. 1990. *Nonlinear Multivariate Analysis*. Wiley.
- Greenacre, Michael J., and Jorg Blasius. 2006. *Multiple Correspondence Analysis and Related Methods*. Chapman; Hall.

- Guttman, Louis. 1941. "The Quantification of a Class of Attributes: A Theory and Method of Scale Construction." In *The Prediction of Personal Adjustment*, edited by Paul Horst. Social Science Research Council.
- Heiser, Willem J. 1981. "Unfolding Analysis of Proximity Data." PhD thesis, Leiden University.
- Husson, François, Julie Josse, and Gilbert Sapiro. 2016. "Jan de Leeuw and the French School of Data Analysis." *Journal of Statistical Software* 73 (6): 1–14. <https://www.jstatsoft.org/article/view/v073i06>.
- Kruskal, Joseph B. 1964a. "Multidimensional Scaling by Optimizing Goodness of Fit to a Nonmetric Hypothesis." *Psychometrika* 29: 1–27.
- Kruskal, Joseph B. 1964b. "Nonmetric Multidimensional Scaling: a Numerical Method." *Psychometrika* 29: 115–29.
- Le Roux, Brigitte, and Henry Rouanet. 2010. *Multiple Correspondence Analysis*. Sage.
- Michailidis, George, and Jan De Leeuw. 2001. "Data Visualization through Graph Drawing." *Computational Statistics* 16: 435–50.
- Okabe, Atsuyuki, Barry Boots, Kokichi Sugihara, and Chiu Sung-Nok. 2000. *Spatial Tessellations: Concepts and Applications of Voronoi Diagrams*. Second Edition. Wiley.
- Tenenhaus, Michel, and Forest W. Young. 1985. "An Analysis and Synthesis of Multiple Correspondence Analysis, Optimal Scaling, Dual Scaling, Homogeneity Analysis and Other Methods for Quantifying Categorical Multivariate Data." *Psychometrika* 50: 91–119.