

Voronoi Homogeneity Analysis of Categorical Data

Jan de Leeuw

August 29, 2026

TBD

Table of contents

1	Introduction	3
1.1	Multiple Correspondence Analysis	3
1.2	Voronoi Homogeneity Analysis	6
1.3	Vertical Homogeneity Analysis	8
2	Algorithm	9
2.1	Initial Configuration	9
2.2	Monotone Regression	9
2.3	Majorization	10
3	Constraints	14
3.1	External Unfolding	14
3.2	Column-Centering	14
3.3	Orthogonality	14
3.4	Rank Constraints	14
3.5	Centroid	15
4	Software	16
4.1	Main	16
4.2	Subroutines	17
4.3	Plots	17

5	Examples	18
5.1	Small Example	18
5.2	GALO	19
5.3	Cetacea	25
6	Appendix: Cetacea	34
	References	35

Note: This is a working manuscript which will be expanded/updated frequently. All suggestions for improvement are welcome. All Rmd, tex, html, pdf, R, and C files are in the public domain. Attribution will be appreciated, but is not required. The files can be found at <https://github.com/deleeuw/voronoi>

1 Introduction

1.1 Multiple Correspondence Analysis

Multiple Correspondence Analysis (MCA) can be introduced and implemented in many different ways. We refer to Guttman (1941), Tenenhaus and Young (1985), Bekker and De Leeuw (1988), Gifi (1990), Greenacre and Blasius (2006), Le Roux and Rouanet (2010), Husson et al. (2016), and De Leeuw (2023) for a bouquet of discussions and interpretations of the equations and algorithms defining the MCA technique.

First some terminology. A *variable* is a function that maps on a (finite or infinite) domain of *objects* into a (finite or infinite) codomain of *values*. In MCA the data are the values of m variables on the same *carrier*, a finite subset the domain consisting of n objects. Each variable j has a finite number k_j of *categories*. The categories constitute the *range* of the variable for the n objects in the carrier. Note that all variables have the same carrier. Each object maps into a unique category, and each category contains a subset of objects from the carrier. We also say that a category has a number of objects that “belong to” the category.

Here is a small example with $n = 10$ and $m = 3$, taken from Gifi (1990), page 65. In the github voronoi repository these data are in data/small.R.

	V1	V2	V3
1	a	p	u
2	b	q	v
3	a	r	v
4	a	p	u
5	b	p	v
6	c	p	v
7	a	p	u
8	a	p	v
9	c	p	v
10	a	p	v

The key mathematical object in MCA is the *indicator matrix*, which is a classical way to code categorical data. Coding variables using indicator matrices is also known as *dummy coding*. For a variable j with k_j categories it is an $n \times k_j$ binary matrix G_j in which each row i contains a single element equal to one, indicating into which category the object i maps for variable j . The cross product $G_j'G_j$ is diagonal, with the *marginal frequencies* of the categories on the diagonal. For two different variables the crossproduct $G_j'G_\ell$ is the $k_j \times k_\ell$ *cross table* or *contingency table* of the two variables. Here are the three indicator matrices of our small example, next to each other.

		a	b	c		p	q	r		u	v	
[1,]		1	0	0		1	0	0		1	0	
[2,]		0	1	0		0	1	0		0	1	
[3,]		1	0	0		0	0	1		0	1	
[4,]		1	0	0		1	0	0		1	0	
[5,]		0	1	0		1	0	0		0	1	
[6,]		0	0	1		1	0	0		0	1	
[7,]		1	0	0		1	0	0		1	0	
[8,]		1	0	0		1	0	0		0	1	
[9,]		0	0	1		1	0	0		0	1	
[10,]		1	0	0		1	0	0		0	1	

We now deviate from the usual Gifi and French approaches (Husson et al. (2016)) and define MCA as a form of non-metric multidimensional scaling, using the *stress* loss function (Kruskal (1964a), Kruskal (1964b)). Define¹

$$\sigma(X, Y_1, \dots, Y_m, \Delta_1, \dots, \Delta_m) := \sum_{j=1}^m \sum_{i=1}^n w_{ij} \sum_{\ell=1}^{k_j} (\delta_{i\ell}^j - d(x_i, y_\ell^j))^2, \quad (1)$$

with w_{ij} given non-negative weights, $d(x_i, y_\ell^j)$ the *Euclidean distance* between row i of X and row ℓ of Y_j , and $\delta_{i\ell}^j$ the corresponding *disparity*. In classical MCA there are no weights, i.e. all w_{ij} in (1) are equal to one. We have added weights to the definition of MCA so that, for example, we can easily incorporate missing data. Disparities are constrained to satisfy the measurement restrictions implied by the data. In MCA this means that we require that $\delta_{i\ell}^j = 0$ whenever $g_{i\ell}^j = 1$, or, using the elementwise product $\otimes$ of matrices, $\Delta_j \otimes G_j = 0$.

Define W_j as the diagonal matrix of weights for variable j , and $W_\star$ as the sum of the W_j . MCA minimizes (1) over the $n \times p$ configurations X , over the Y_j , and over the Δ_j that satisfy the measurement restrictions. There may be additional constraints on X and/or the Y_j , which define different variations of MCA.

Define

$$\sigma_\star(X, Y_1, \dots, Y_m) := \min_{\Delta_1, \dots, \Delta_m} \sigma(X, Y_1, \dots, Y_m, \Delta_1, \dots, \Delta_m). \quad (2)$$

The minimizing disparity $\delta_{i\ell}^j$ is equal to zero if $g_{i\ell}^j = 1$ and equal to $d(x_i, y_\ell^j)$ otherwise. Consequently

$$\sigma_\star(X, Y_1, \dots, Y_m) = \sum_{j=1}^m \sum_{i=1}^n w_{ij} \sum_{\ell=1}^{k_j} \{d^2(x_i, y_\ell^j) \mid g_{i\ell}^j = 1\}. \quad (3)$$

¹The symbol $:=$ is used for definitions

We can make a graphical representation of (3). For variable j we plot the n points x_i and the k_j points y_ℓ^j in a joint plot (or *biplot*). We then connect, with straight lines, each category point with the object points that belong to it. For each category this defines a figure in the plot which we call a *star*. The *size* of a star is the weighted sum of squares of the length of its lines. The loss function (3) for variable j is the weighted sum of squares of the length of all n lines in this plot, or equivalently the sum of the sizes of the k_j stars. The total loss is the sum of star-sizes over all m plots. Thus MCA aims to place the object-points and category-points in space in such a way that the stars are as small as possible, i.e. objects are as close as possible to the categories they belong to.

In matrix notation

$$\sigma_\star(X, Y_1, \dots, Y_m) = \sum_{j=1}^m \text{tr} \{X' W_j X - 2 X' W_j G_j Y_j + Y_j' D_j Y_j\} \quad (4)$$

Here D_j is the diagonal matrix with column-sums of $W_j G_j$ on the diagonal. In classical MCA $D_j = G_j' G_j$, but if there are weights this is no longer the case.

Next define

$$\sigma_{\star\star}(X) := \min_{Y_1, \dots, Y_m} \sigma_\star(X; Y_1, \dots, Y_m). \quad (5)$$


The minimum of (4) over the Y_j is attained at $\hat{Y}_j := D_j^{-1} G_j' W_j X$. Category-points are the weighted centroids of the object-points belonging to the category. This makes “stars” actually look like stars. If we substitute these optimal $\hat{Y}_j$ in (3) we see that $\sigma_{\star\star}(X)$ is indeed equal to the weighted sum of squares of all lines defining the stars. Minimizing $\sigma_{\star\star}(X)$ over $X' W_\star X = I$, and thus minimizing (1) over (X, Y_j, Δ_j) , defines MCA.

By writing

$$\sigma_{\star\star}(X) := \text{tr} X' \left\{ \sum_{j=1}^m (W_j - W_j G_j D_j^{-1} G_j' W_j) \right\} X \quad (6)$$

we see that minimizing (6) over $X' W_\star X = I$ is a straightforward eigenvalue-eigenvector problem.

A MCA in two dimensions of our small example produces the following three star plots. Objects are numbered and in BLUE, categories have bigger labels and are in RED. Note that there are ten objects, but the sets of objects (1, 4, 7), (8, 10), and (6, 9) have the same *profile* and consequently get mapped into the same object-point. Thus there are at most six different object-points. The analysis minimizes the sum of squares of the length of all black lines in the three plots, under the condition that the matrix of object scores is column-centered and orthonormal. We can think of the results as a form of *graph drawing*, where the graph is the union of the bipartite graphs of all m “belonging to” relations (De Leeuw and Michailidis (2000), Michailidis and De Leeuw (2001)).



1.2 Voronoi Homogeneity Analysis

Although using loss function (1) makes MCA a form of non-metric multidimensional scaling there are some important differences with the more familiar MDS versions minimizing stress. There is no local minima problem in MCA, because it turns out to be a symmetric eigenvalue problem. In MCA, unlike in MDS, we impose the constraint $X'X = I$. Also, and perhaps most importantly, in MCA we do not really expect the minimum loss function value to be zero, or even close to zero. The loss function measures deviations from a very unrealistic “model”. Zero loss would imply all object-points coincide with all category-points they belong to. This can only be realized, except in trivial cases, if all object and category-points map into a single point, which is prohibited by the orthonormality constraint on the object scores. Thus we get a perfectly determinate and stable solution, at low computational cost, but by imposing strong constraints on the disparities and a rather arbitrary normalization. We can get around this seemingly unfortunate situation by admitting we are not fitting or testing a “model”, but we are creating “best” drawings of the bipartite graphs that constitute the data.

The lack of a realistic model makes it interesting to continue use the more general loss function

$$\sigma(X, Y_1, \dots, Y_m, \Delta_1, \dots, \Delta_m) := \sum_{j=1}^m \sum_{i=1}^n \sum_{\ell=1}^{k_j} w_{i\ell}^j (\delta_{i\ell}^j - d(x_i, y_\ell^j))^2, \quad (7)$$

with the constraints that

$$\text{if } g_{i\ell}^j = 1 \text{ then } \delta_{i\ell}^j \leq \delta_{i\nu}^j \text{ for all } \nu = 1, \dots, k_j. \quad (8)$$

This is significantly weaker than the MCA constraints. If object i belongs to category ℓ of variable j we do no longer require the object-point to coincide with the category-point, but we merely require that the object-point is at least as close to category-point ℓ as to any of the other $k_j - 1$ category-points of that variable.

Constraints (8) are called the *horizontal constraints*. The *vertical constraints* are

$$\text{if } g_{i\ell}^j = 1 \text{ and } g_{k\ell}^j = 0 \text{ then } \delta_{i\ell}^j \leq \delta_{k\ell}^j. \quad (9)$$

Voronoi Homogeneity Analysis (VHA) of categorical data minimizes loss function (1) over X , over the Y_j and over the Δ_j satisfying the relaxed ordinal constraints. It is consequently another form of non-metric (ordinal) multidimensional scaling. To be more precise, we have m linked non-metric multidimensional unfolding problems for the binary matrices G_j , which are “linked” because they share the same X .

Because of the different ordinal constraints on the disparities, the plots corresponding with this analysis are different. As in MCA there is a plot for each variable j with both n object-points and k_j category-points. The k_j category-points are the *generators* of k_j *Voronoi regions* that partition the space. Voronoi region $\mathcal{V}_\ell^j$ is the region of space where the points are at least as close to generator y_ℓ^j as to any of the other generators of variable j . And loss (4) is zero (for a variable) if all object-points are in the “correct” Voronoi region, i.e. in the Voronoi region of the generator corresponding with the category they belong to.

One (inefficient, but conceptually easy) way of constructing Voronoi regions in the plane is to draw the perpendicular bisectors of the (hypothetical) lines connecting all pairs of category-points of a variable. In higher dimensions $p > 2$ the bisectors are planes of dimension $p - 1$. The convex polyhedral regions thus formed, some bounded and some unbounded, define the Voronoi regions². The literature on properties and computation of Voronoi regions is large, and we refer to the textbooks of Okabe et al. (2000) and Aurenhammer et al. (2013).

Since the category-point itself is obviously in its own Voronoi region and since Voronoi regions are convex it follows that if we make stars by connecting category-points with object points as in MCA then we actually require each star to be in its Voronoi region. It is no longer true that a category-point is the centroid of points of the objects belonging to the category. The category-point can even be outside the convex hull of the object-points belonging to the category.

The constraint $X'X = I$ from MCA is no longer imposed directly, but we somehow have to deal with the fact that in VHA the minimum of loss function (1) is always zero. The trivial solution, in which all points and regions collapse into a single point and all Δ_j are zero, is always available and not excluded by the constraints. But, perhaps more urgently, we have to deal with the types of degenerate solutions familiar from multidimensional unfolding (Heiser (1981), De Leeuw (1983), Busing (2010)). For example, we can collapse all category-points of a variable into a single point and put all object-points on a sphere with the category-point at its center. Then all distances between category and object-points are the same and we can choose all disparities equal to that distance value and obtain loss zero.

We will use some constraints that tend to counteract degeneracy. Also, we always start our iterations with the MCA analysis, which in most cases already has small loss. Then we perform iterations until either the distances do not change from one iteration to the next (up to some

²In the plane a Voronoi region is unbounded if and only if its generator is on the boundary of the convex hull of all generators.

epsilon) or the loss becomes zero (up to some epsilon) or we have reached the maximum number of iterations allowed.

1.3 Vertical Homogeneity Analysis

Constraints (8) are called the *horizontal constraints*. The *vertical constraints* are

$$\text{if } g_{i\ell}^j = 1 \text{ and } g_{k\ell}^j = 0 \text{ then } \delta_{i\ell}^j \leq \delta_{k\ell}^j. \quad (10)$$

Thus object-points belonging to a category must be inside a ball centered at the category point (or on its boundary), and objects not belonging to that category must be outside the ball (or on its boundary). Clearly an object cannot be in the interiors of two different balls of two categories of the same variable – if it was it would be in two categories of that variable. Thus convex hulls of the point clouds in balls corresponding with different categories of the same variable are disjoint, which implies that object points are in the correct Voronoi regions.

2 Algorithm

We use an *Alternating Least Squares* or *ALS* algorithm, similar to the approach used in many other non-metric scaling methods, to minimize loss (5). For context, see Borg and Groenen (2005). We alternate minimization over $\Delta_1, \dots, \Delta_m$ for given X and $Y_1 \dots Y_m$ with minimization over X and $Y_1 \dots Y_m$ for given $\Delta_1, \dots, \Delta_m$. Using superscript (μ) for the iteration index we can write for an ALS iteration

$$(\Delta_1^{(\mu)}, \dots, \Delta_m^{(\mu)}) = \underset{\Delta_1, \dots, \Delta_m}{\operatorname{argmin}} \sigma(X^{(\mu)}, Y_1^{(\mu)}, \dots, Y_m^{(\mu)}, \Delta_1, \dots, \Delta_m), \quad (11a)$$

$$(X^{(\mu+1)}, Y_1^{(\mu+1)}, \dots, Y_m^{(\mu+1)}) = \underset{X, Y_1, \dots, Y_m}{\operatorname{argmin}} \sigma(X, Y_1, \dots, Y_m, \Delta_1^{(\mu)}, \dots, \Delta_m^{(\mu)}). \quad (11b)$$

The first substep (11a) of iteration (μ) solves a monotone regression problem for a simple partial order, in which one of k_j elements must be smaller than the other $k_j - 1$ elements. The second substep (11b) solves a metric multidimensional scaling problem. This second subproblem cannot be solved completely, because it would need an infinite iterative process by itself. Thus the second substep (11b) consists of a finite number of *inner iterations* within the second substep of a single ALS *outer iteration*.

2.1 Initial Configuration

In all our analysis we use MCA as the initial configuration for both object-scores and category-scores. The algorithm for MCA is the ALS method used in Gifi (1990) to compute the truncated SVD. The repository includes mca.R

2.2 Monotone Regression

The nm monotone regressions are applied for each object i and each variable j . Forgetting about all these indices for the moment we need to minimize the sum of squares $\|\delta - d\|^2$ over δ , where one of the elements, say δ_ν , must be smaller than or equal to all others.

This can be solved by using monotone regression with the primary approach to ties (Kruskal (1964b), De Leeuw (1977)), treating all δ_ℓ with $\ell \neq \nu$ as ties.

Suppose y and $w \geq 0$ are given vectors with n elements. Suppose N_1 and N_0 are a partition of $N := \{1, 2, \dots, n\}$. The Binary Monotone Regression Problem is to find x with $x_i \leq x_k$ for all $i \in N_0$ and $k \in N_1$ such that

$$\sigma(x) = \frac{1}{2} \sum_{i \in N} w_i (x_i - y_i)^2$$

is minimized. Equivalently we want to find α such that $x_i \leq \alpha$ for $i \in N_0$ and $x_i \geq \alpha$ for $i \in N_1$. Thus loss can be rewritten as a function of the single variable α .

$$\sigma(\alpha) = \frac{1}{2} \sum_{i \in N_0} w_i \min(\alpha - y_i, 0)^2 + \frac{1}{2} \sum_{i \in N_1} w_i \max(\alpha - y_i, 0)^2.$$

This function is piecewise quadratic, convex, and differentiable (although not twice differentiable). Its derivative, which is increasing, is

$$\mathcal{D}\sigma(\alpha) = \sum_{i \in N_0} w_i (\min(\alpha - y_i, 0) + \sum_{i \in N_1} w_i (\max(\alpha - y_i, 0))$$

The function `binaryMonotoneRegression()`, with the alias `bmr()`, minimizes σ over α . We sort y , set α equal to the smallest y_i , and compute $\mathcal{D}\sigma(\alpha)$. If the derivative is zero, we are done. If it is negative we set α to the next smallest y_i . We continue this until we arrive at an α for which the derivative is either zero or positive. If it is zero we stop. If it is positive then the minimum is between this α and the previous α , for which the derivative was still negative. Since the derivative is linear on the interval between the two values of α we can use linear interpolation to find the value for which it is zero.

2.3 Majorization

We study the off-diagonal smacof algorithm for MDS, without taking into account that the loss function (1) comes from n observations on m categorical variables. This improves and generalizes the smacof treatment of the metric unfolding problem in Heiser and De Leeuw (1979).

We rewrite stress as

$$\sigma(X, Y) := \sum_{i=1}^n \sum_{k=1}^K w_{ik} (\delta_{ik} - d(x_i, y_k))^2. \quad (12)$$

The distance is

$$d(x_i, y_k) = \sqrt{(X'e_i - Y'e_k)'(X'e_i - Y'e_k)}. \quad (13)$$

We will construct a majorization of (12) at $\tilde{X}$ and $\tilde{Y}$. Think of $\tilde{X}$ and $\tilde{Y}$ as the current best estimates in a sequence of iterations. By Cauchy-Schwartz,

$$d(x_i, y_k) \geq \frac{1}{d(\tilde{x}_i, \tilde{y}_k)} (X'e_i - Y'e_k)'(\tilde{X}'e_i - \tilde{Y}'e_k). \quad (14)$$

Define the $n \times K$ matrix-valued function $B(X, Y)$ by

$$b_{ik}(X, Y) := \begin{cases} w_{ik} \frac{\delta_{ik}}{d(x_i, y_k)} & \text{if } d(x_i, y_k) > 0, \\ 0 & \text{if } d(x_i, y_k) = 0. \end{cases} \quad (15)$$

Also define the diagonal matrices $B_\star(X, Y)$ with the row-sums of $B(X, Y)$ and $B^\star(X, Y)$ with the column-sums.

If we define, as we usually do in smacof,

$$\rho(X, Y) := \sum_{i=1}^n \sum_{k=1}^K w_{ij} \delta_{ij} d_{ij}(x_i, y_k) \quad (16a)$$

then $\rho(X, Y) \geq \rho(X, Y; \tilde{X}, \tilde{Y})$, where

$$\rho(X, Y; \tilde{X}, \tilde{Y}) := \text{tr } X' B_\star(\tilde{X}, \tilde{Y}) \tilde{X} - \text{tr } X' B(\tilde{X}, \tilde{Y}) \tilde{Y} - \text{tr } Y' B'(\tilde{X}, \tilde{Y}) \tilde{X} + \text{tr } Y' B^\star(\tilde{X}, \tilde{Y}) \tilde{Y}. \quad (16b)$$

Also define

$$\eta^2(X, Y) := \sum_{i=1}^n \sum_{k=1}^K w_{ik} \delta_{ij} d_{ij}^2(x_i, y_k) = \text{tr } X' W_\star X - 2 \text{tr } X' W Y + \text{tr } Y' W^\star Y, \quad (16c)$$

with $W_\star$ and $W^\star$ the diagonal matrices with row-sums and column-sums of W , as well as the constant

$$C := \sum_{i=1}^n \sum_{k=1}^K w_{ik} \delta_{ik}^2 \quad (16d)$$

With these definitions

$$\sigma(X, Y) = C - 2 \rho(X, Y) + \eta^2(X, Y), \quad (17)$$

and $\sigma(X, Y) \leq \sigma(X, Y; \tilde{X}, \tilde{Y})$, where

$$\sigma(X, Y; \tilde{X}, \tilde{Y}) := C - 2 \rho(X, Y; \tilde{X}, \tilde{Y}) + \eta^2(X, Y), \quad (18)$$

with equality if $X = \tilde{X}$ and $Y = \tilde{Y}$. In a majorization (or MM) iteration we minimize $\sigma(X, Y; \tilde{X}, \tilde{Y})$ over X and Y . If the minimum is attained at $\tilde{X}$ and $\tilde{Y}$ we stop the algorithm, because we have found a stationary point. If not we have found updates X^+ and Y^+ such that

$$\sigma(X^+, Y^+) \leq \sigma(X^+, Y^+; \tilde{X}, \tilde{Y}) < \sigma(\tilde{X}, \tilde{Y}; \tilde{X}, \tilde{Y}) = \sigma(\tilde{X}, \tilde{Y}). \quad (19)$$

The stationary equations for minimizing $\sigma(X, Y; \tilde{X}, \tilde{Y})$ are

$$X = W_{\star}^{-1}(WY + B_{\star}(\tilde{X}, \tilde{Y})\tilde{X} - B(\tilde{X}, \tilde{Y})\tilde{Y}), \quad (20a)$$

$$Y = \{W^{\star}\}^{-1}(W'X + B^{\star}(\tilde{X}, \tilde{Y})\tilde{Y} - B'(\tilde{X}, \tilde{Y})\tilde{X}). \quad (20b)$$

Alternatively

$$\begin{bmatrix} W_{\star} & -W \\ -W' & W^{\star} \end{bmatrix} \begin{bmatrix} X \\ Y \end{bmatrix} = \begin{bmatrix} B_{\star}(\tilde{X}, \tilde{Y}) & -B(\tilde{X}, \tilde{Y}) \\ -B'(\tilde{X}, \tilde{Y}) & B^{\star}(\tilde{X}, \tilde{Y}) \end{bmatrix} \begin{bmatrix} \tilde{X} \\ \tilde{Y} \end{bmatrix} \quad (21)$$

The next step in each iteration could be to directly solve the system (21) for X and Y . This approach has the disadvantage that the order is $n + m$, which can be quite large in applications. We also have to take into account that the coefficient matrix of (21) is singular. Alternatively we can substitute the expression for X from (20a) in (20b) and then solve the resulting equation for Y . Or we can solve the system in (20a) and (20b) iteratively by ALS. Choose a Y , then compute X from (20a), then use that X in (20b) to find a new Y , and so on.

In this paper we use an approach similar to ALS, which could be called MALS for Majorization by Alternating Least Squares. In the first step of a MALS cycle we majorize $\sigma(X, \tilde{Y})$ and minimize $\sigma(X, \tilde{Y}; \tilde{X}, \tilde{Y})$. Thus we treat the loss as a function of X only, and by the earlier results in this section we have

$$X^+ = W_{\star}^{-1}(W\tilde{Y} + B_{\star}(\tilde{X}, \tilde{Y})\tilde{X} - B(\tilde{X}, \tilde{Y})\tilde{Y}). \quad (22a)$$

The next step, in the same MALS iteration, is to use this new X^+ and minimize $\sigma(X^+, Y; X^+, \tilde{Y})$ over Y , giving

$$Y^+ = \{W^{\star}\}^{-1}(W'X^+ + B^{\star}(X^+, \tilde{Y})\tilde{Y} - B'(X^+, \tilde{Y})X^+) \quad (22b)$$

Then proceed to the next two-step MALS iteration, using the new X^+ and Y^+ .

One advantage of MALS, shared with ALS, is that the only matrix inversion needed is of the diagonal weight marginals. Also there is no need to work with square symmetric matrices of order $n + m$. But the main advantage of MALS is that it can easily incorporate various constraints on X and Y . Remember that

$$\sigma(X, \tilde{Y}; \tilde{X}, \tilde{Y}) = \text{tr } X'W_{\star}X - 2 \text{tr } X'\{W\tilde{Y} - \text{tr } X'B_{\star}(\tilde{X}, \tilde{Y})\tilde{X} + \text{tr } X'B(\tilde{X}, \tilde{Y})\tilde{Y}\} + C, \quad (23)$$

where C is now used generically for terms not involving X . Define

$$\bar{X} := W\tilde{Y} - \text{tr } X'B_{\star}(\tilde{X}, \tilde{Y})\tilde{X} + \text{tr } X'B(\tilde{X}, \tilde{Y})\tilde{Y} \quad (24)$$

Then, after completing the square,

$$\sigma(X, \tilde{Y}; \tilde{X}, \tilde{Y}) = \text{tr} (X - \bar{X})' W_{\star} (X - \bar{X}) + C. \quad (25a)$$

Thus minimizing $\sigma(X, \tilde{Y}; \tilde{X}, \tilde{Y})$ over X that must satisfy some constraints means projecting $\bar{X}$ in the diagonal $W_{\star}$ metric on the set defined by the constraints. Of course in the same way

$$\sigma(\tilde{X}, Y; \tilde{X}, \tilde{Y}) = \text{tr} (Y - \bar{Y})' W^{\star} (Y - \bar{Y}) + C. \quad (25b)$$

3 Constraints

3.1 External Unfolding

$X = ZA$ with Z a known matrix.

3.2 Column-Centering

If we require $X'W_r e = 0$ then the stationary equations are $W_r(X - \bar{X}) = W_r e \lambda'$ and thus $X = \bar{X} + e \lambda'$ with $\lambda' = -e' W_r \bar{X} / e' W_r e$ or

$$X = (I - \frac{ee' W_r}{e' W_r e}) \bar{X}$$

This is different from updating X as $J_n \tilde{X}$ and Y as $\tilde{Y} - n^{-1} ee' \tilde{X}$. With this alternative update, in which we subtract the column-means of $\tilde{X}$ from both $\tilde{X}$ and $\tilde{Y}$, the distances are the same before and after centering. Thus the distances in all iterations, and consequently the loss function values, are the same with or without this type of centering.

3.3 Orthogonality

If we require $X'W_\star X = I$ we have the stationary equations $\bar{X} = XM$, with M a symmetric matrix of Lagrange multipliers. Thus

$$X = \text{procrustus}(\bar{X}) := \bar{X}(\bar{X}' W_\star \bar{X})^{-\frac{1}{2}}.$$

If we require both $X'W_\star X = I$ and $X'W_\star e = 0$ we have

$$X = \text{procrustus}((I - \frac{ee' W_\star}{e' W_\star e}) \bar{X})$$

3.4 Rank Constraints

One of the main themes in Gifi (1990) is that the $k_j \times p$ matrices of category-scores can be constrained to be of rank $r < p$. We will restrict ourselves in this paper to $r = 1$, which defines what Gifi calls *single quantifications*. If all variables are restricted to have single quantifications then a MCA becomes equivalent to a PCA of the transformed variables.

In VHA the Voronoi regions of single variables become parallel strips. The category-points of a variable are on a line through the origin, and the Voronoi lines or planes are parallel and perpendicular to the line with the category-points.

$$\text{tr} (Y - za')' W_c (Y - za') = -2z' W_c Y a + z' W_c z . a' a$$

So rank one approximation of $\tilde{Y} + n^{-1}ee'(X - \tilde{X})$. If X is column-centered then of course $\tilde{Y} - ee'\tilde{X}$.

3.5 Centroid

A more complicated restriction is $Y = D^{-1}G'X$, possibly together with $X'X = I$ and $e'X = 0$. Minimization over X and the Y_j is no longer separated. In fact, the *centroid constraints* effectively eliminate the Y_j from the optimization problem. We have to adapt the majorization method to take these new constraints into account.

If $Y_j = D_j^{-1}G'_jX$ then $d_{i\ll}^j(X, Y_j)$ only depends on X and we have

$$\{d_{i\ell}^j(X)\}^2 = \text{tr} X' \{e_i e'_i - 2 \{d_\ell^j\}^{-1} g_\ell^j e'_i + \{d_\ell^j\}^{-2} g_\ell^j \{g_\ell^j\}'\} X. \quad (26a)$$

Moreover for each pair X and $\tilde{X}$ we have, by Cauchy=Schwartz,

$$d_{i\ell}^j(X) \geq \frac{1}{d_{i\ell}^j(\tilde{X})} \text{tr} X' \{e_i e'_i - 2 \{d_\ell^j\}^{-1} g_\ell^j e'_i + \{d_\ell^j\}^{-2} g_\ell^j \{g_\ell^j\}'\} \tilde{X} \quad (26b)$$

Suppose R_W^j and C_W^j are diagonal matrices with row-sums and column=column-sums of W_j . Then

$$\sum_{i=1}^n \sum_{\ell=1}^{k_j} w_{i\ell}^j \{d_{i\ell}^j(X)\}^2 = \text{tr} X' \{R_W^j - G_j D_j^{-1} W_j' - W_j D_j^{-1} G_j' + G_j (C_W^j D_j^{-2}) G_j'\} X, \quad (27a)$$

and

$$\sum_{i=1}^n \sum_{\ell=1}^{k_j} w_{i\ell}^j d_{i\ell}^j(X) \geq \text{tr} X' \{R_{\tilde{B}}^j - G_j D_j^{-1} \tilde{B}_j' - \tilde{B}_j D_j^{-1} G_j' + G_j (C_{\tilde{B}}^j D_j^{-2}) G_j'\} \tilde{X}. \quad (27b)$$

In (27b) the matrix $\tilde{B}_j$ is defined as

$$\tilde{b}_{i\ell}^j := \begin{cases} w_{i\ell}^j \frac{\delta_{i\ell}^j}{d_{i\ell}^j(\tilde{X})} & \text{if } d_{i\ell}^j(\tilde{X}) > 0, \\ 0 & \text{if } d_{i\ell}^j(\tilde{X}) = 0, \end{cases} \quad (28)$$

and $R_{\tilde{B}}^j$ and $C_{\tilde{B}}^j$ are the diagonal matrices of its row-sums and column-sums.

4 Software

4.1 Main

The main program `voronoiHomogeneityAnalysis()`, aliased as `vha()`, has the following arguments (with default values).

```
voronoiHomogeneityAnalysis <- function(theData,
                                       ndim = 2,
                                       wght = NULL,
                                       xextr = NULL,
                                       xcent = FALSE,
                                       xnorm = FALSE,
                                       yrank = ndim,
                                       itmax = 1000,
                                       eps = 1e-6,
                                       aps = 1e-6,
                                       verbose = TRUE)
```

- *theData* is an $n \times m$ matrix with n observations on m categorical variables.
- *ndim* is the number of dimensions of the solution.
- *wght* is either NULL or a list of $n \times k_j$ matrices. If *wght* is NULL all weights are one.
- *dnorm* if TRUE the monotone regression is normalized.
- *xcent* if TRUE object scores are W-centered.
- *xnorm* if TRUE object scores are W-orthonormal.
- *yrank* rank restrictions on category-scores. If the length of *yrank* is one it is expanded to a vector of length m .
- *itmax* is the maximum number of iterations.
- *eps* is the cutoff for the change in distances from one iteration to another.
- *aps* is the cutoff for the loss function value. If loss is smaller than *aps* the program stops.
- *verbose* if TRUE iteration information is printed to stdout.

The value returned by the program is a list.

```
return(
  list(
    xmat = xnew,
    ymat = ynew,
    dmat = dnew,
    dhat = dhat,
```

```

    xini = xini,
    yini = yini,
    itel = itel,
    loss = snew,
    indi = indi,
    ncat = ncat,
    wght = wght
  )
)

```

- *xmat* object scores, an $n \times p$ matrix, with p equal to $ndim$.
- *ymat* category-scores, a list of m matrices of dimension $k_j \times p$.
- *dmat* distances, a list of m matrices of dimension $n \times k_j$.
- *dhat* disparities, a list of m matrices of dimension $n \times k_j$.
- *xini* the MCA solution for the object scores,
- *yini* the MCA solution for the category-scores,
- *itel* number of iterations.
- *loss* final value of stress.
- *indi* indicators, a list of m matrices of dimension $n \times k_j$.
- *ncat* a vector with the number of categories of the m variables.
- *wght* weights, a list of m matrices of dimension $n \times k_j$.

4.2 Subroutines

4.3 Plots


The `plotVHA()` routine makes joint plots of object scores and category scores for a single variable. It can label the points in different ways and it can draw some additional geometric objects in the plots. The arguments are

- *xmat* $n \times p$, object scores,
- *ymat* $k_j \times p$, category scores variable j ,
- *indi* $n \times k_j$, indicator matrix variable j ,
- *dmat* $n \times k_j$, distance matrix variable j ,
- *type* “voronoi” (default), “circle”, or “hull”,
- *main* NULL (default), or plot label,
- *ylab* = 1,
- *xlab* = 1,
- *pdf* plot device, default FALSE)


5 Examples

5.1 Small Example


We start with the small example we used before to illustrate MCA. We first do an analysis without additional configuration constraints, using the horizontal (row-wise) ordinal constraints on the dissimilarities.



Loss is 0.0000000000 after 443 iterations. If we compare the plots from the VHA with the star plots from the initial MCA for this example we see that the algorithm did not have to change much to obtain perfect fit. One thing that does change is that the iterations moves the category-points “p” and “r” for variable 2 very close to each other. At the point where we stop the iterations they are still distinguishable, and thus their perpendicular bisector is still a line in the Voronoi plot. If we would continue iterating “p” and “r” would coincide and there would only be two Voronoi regions: one containing category-point “q” and object-point 2, and one containing all other points.



Next we repeat the analysis, but this time with vertical (column-wise) constraints. We show the circle-plots for this solution.



Note that for variable 2 the p-circle is within the r-circle, and there is a p-object on the boundary of the r-circle and an r-object on the boundary of the p-circle. The intersection of the interiors of these two circle do not contain any objects. For the third variable the three u-objects (which all have the same “apu” profile and thus have a circle with zero radius) are on the boundary of the v-circle. Loss is 0.0000000000 after 173 iterations.

5.2 GALO

Our next example use the GALO data from Peschar (1975). In the github voronoi repository the data are in data/galo.R. Subjects are 1290 children who in 1959/1960 were in sixth grade of primary school in the city of Groningen. Variables are SEX, IQ (in nine ordered categories), ADV (sixth grade teachers advice on the form of secondary education appropriate for the student), and SES. The categories of ADV are as follows (alphabetically).

- *Agr* Agricultural.
- *Ext* Extended primary (like middle school).
- *Gen* General (like high school).
- *Grls* Extended primary for girls (like middle school).
- *Man* Vocational, including housekeeping.
- *None* No secondary education.

The categories of SES are as follows (again alphabetically).

- *LowWC* Lower white collar.
- *MidWC* Middle white collar.
- *Prof* Professional.
- *Shop* Shopkeepers.
- *Skil* Skilled manual.

- *Unsk Unskilled Manual.*

We start with an MCA of the galo data. The initial value of the loss function is 0.3158339145. The loss function value is computed by using

$$\sigma_{\star}(X, Y_1, \dots, Y_m) := \min_{\Delta_1, \dots, \Delta_m} \sigma(X, Y_1, \dots, Y_m, \Delta_1, \dots, \Delta_m), \quad (29)$$

where X and the Y_j are the MCA solution, and where the minimum over the Δ_j is computed by requiring that $\delta_{iv}^j \leq \delta_{il}^j$ if $g_{iv}^j = 1$. This is *not* the MCA loss function (7), in which the minimum over the Δ_j is computed by requiring that $\delta_{il}^j = 0$ whenever $g_{il}^j = 1$. Thus we evaluate the MCA solution using the VGA loss function.

For any MCA or VGA solution we can compute a Shepard diagram by plotting the distances and disparities, pooled over all objects (individuals) and variables. The weak regression requirements imply that many disparities will be equal to the corresponding distances. This is illustrated by the diagonal of Figure 1.

The MCA solution can also be used to produce Voronoi plots, which are given in Figure 2 for IQ. We see the familiar horseshoe shape of the object score cloud (the Guttman effect, see De Leeuw (2023)). The category-scores are largely in order along the horseshoe, and as a consequence the Voronoi regions are unbounded and emanate from a point inside the horseshoe. There are some relatively minor deviations from that pattern. Specifically category 1 does not fit the pattern, but that is not too serious because there are only five individuals in that category. There are quite a few misclassifications, i.e. objects in the “wrong” Voronoi region.

The VHA solution has loss equal to 0.0000818331 after 1000 iterations. The voronoi function stopped because the maximum number of iterations was reached. The $d - \hat{d}$ plot in Figure 3 shows a near-perfect fit.

Figure 4 shows some characteristic properties of Voronoi plots. If the objects in two categories are not well separated the object-points will be on the perpendicular bisector of the category pair. This is partial degeneracy due to requiring weak monotonicity. The algorithm cannot impose strict monotonicity by placing an object in the interior of its Voronoi region, and compromises by putting it on the boundary. Thus the VHA plot will show object-points on lines. Some category scores are very close, which makes the Voronoi diagram unstable. The solution for the category-scores for IQ is

	dim1	dim2	freq
1	-0.03551679	0.0038947482	5
2	-0.02703758	-0.0491982477	17
3	-0.02706029	-0.0427253760	121
4	-0.02290451	-0.0006391076	234
5	-0.02289474	-0.0006162247	390




Figure 1: Galo D-Dhat Plot MCA




Figure 2: Galo IQ MCA




Figure 3: Galo D-Dhat Plot VHA




Figure 4: Galo IQ VHA

6	0.02215996	0.0203126796	244
7	0.02216069	0.0203143237	197
8	0.05467150	-0.0480675505	63
9	0.06483906	-0.0642872376	19

We see that categories 2 and 3 map into basically the same point. The same is true for categories 4 and 5, and for categories 6 and 7. So instead of nine points there are really only six. We can clean up the Voronoi diagram by using only those six categories. Note that we do not recode the data and redo the analysis, we just use the same plot with the same object-scores and just change the diagram. The result is in Figure 5.

5.3 Cetacea

The data in Vescia (1985) describe 36 whales, porpoises and dolphins using 15 categorical variables. In the github voronoi repository the data are in data/cetacea.R. Appendix C gives the common names of the 36 species and the names of the variables.

The variables are mostly physical characteristics, but they also include habitat and feeding. The average number of categories per variable is about four, and almost all variables are “nominal” (i.e. not “ordinal” and not “numerical”). In the same book Van der Burg (1985) presents a secondary analysis of the data with MCA and Guttman (1985) applies his MSA-I technique (which uses MCA as an initial configuration for its iterations).

We start with an MCA of the cetacea data. The initial value of the loss function ... is 0.9945577132.




Figure 5: Galo IQ Corrected VHA




Figure 6: Cetacea D-Dhat Plot MCA




Figure 7: Cetacea Object Scores MCA




Figure 8: Cetacea Flippers MCA

Loss is 0.0000964845 after 1000 iterations.




Figure 9: Cetacea D-Dhat Plot VHA

	dim1	dim2	freq
small	-0.03818544	-0.01751674	12
large and short	-0.03733595	-0.03078537	9
medium size	0.18051322	0.15150857	9
long and narrow	-0.08505453	0.10776408	6




Figure 10: Cetacea Flippers

Loss is 0.0003024926 after 1000 iterations.




Figure 11: Cetacea D-Dhat Plot VHA XX




Figure 12: Cetacea Flippers VHA XX

6 Appendix: Cetacea

[1]	Bowhead whales	Rorquals
[3]	Blue whale	Giant bottle-nosed whales
[5]	Commerson's Dolphins	White whales
[7]	Common dolphins	Grey whales
[9]	Right whales	Pilot whales
[11]	Risso's dolphins	Bottle-nosed whales
[13]	Amazon dolphins	Pygmy sperm whales
[15]	White-sided dolphins	Chinese river dolphins
[17]	Right whale dolphins	Humpback whales
[19]	Sowerby's whales	Narwhals
[21]	Pygmy right whales	Finless black porpoises
[23]	Irawady dolphins	Killer whales
[25]	Common porpoises	Sperm whales
[27]	Gangetic dolphins	False killer whales
[29]	Guyanian river dolphins	Cameroun's dolphins
[31]	Spotted dolphins	Rough toothed dolphins
[33]	La Plata's dolphins	Shepherd's beaked whales
[35]	Bottle-nosed dolphins	Goosebeak whales

[1]	NECK	FORM OF THE HEAD
[3]	SIZE OF THE HEAD	BEAK
[5]	DORSAL FIN	FLIPPERS
[7]	SET OF TEETH	FEEDING
[9]	BLOW HOLE	COLOR
[11]	CERVICAL VERTEBRAE	LACRYMAL AND JUGAL BONES
[13]	HABITAT	LONGITUDINAL FURROWS ON THE THROAT
[15]	HEAD BONES	

References

- Aurenhammer, Franz, Rolf Klein, and Der-Tsai Lee. 2013. *Voronoi Diagrams and Delaunay Triangulations*. World Scientific Publishing Co.
- Bekker, Paul, and Jan De Leeuw. 1988. “Relation Between Variants of Nonlinear Principal Component Analysis.” Chap. 1 in *Component and Correspondence Analysis*, edited by Jan L. A. Van Rijkevorsel and Jan De Leeuw. Wiley Series in Probability and Mathematical Statistics. Wiley.
- Borg, Ingwer, and Patrick J. F. Groenen. 2005. *Modern Multidimensional Scaling*. Second Edition. Springer.
- Busing, Frank M. T. A. 2010. “Advances in Multidimensional Unfolding.” PhD thesis, Leiden University. <https://scholarlypublications.universiteitleiden.nl/access/item%3A2837800/view>.
- De Leeuw, Jan. 1977. “Correctness of Kruskal’s Algorithms for Monotone Regression with Ties.” *Psychometrika* 42: 141–44.
- De Leeuw, Jan. 1983. *On Degenerate Multidimensional Unfolding Solutions*. Research Report. Department of Data Theory FSW/RUL. <https://jansweb.netlify.app/publication/deleeuw-r-06-a/deleeuw-r-06-a.pdf>.
- De Leeuw, Jan. 2023. “Deconstructing Multiple Correspondence Analysis.” Chap. 22 in *Analysis of Categorical Data from Historical Perspectives. Essays in Honour of Shizuhiko Nishisato*, edited by Eric J. Beh, Rosaria Lombardo, and Jose G. Clavel. Springer.
- De Leeuw, Jan, and George Michailidis. 2000. “Graph Layout Techniques and Multidimensional Data Analysis.” In *Game Theory. Optimal Stopping, Probability and Statistics*, edited by F. T. Bruss and L. LeCam. Lecture Notes – Monographs 35. Institute of Mathematical Statistics.
- Gifi, Albert. 1990. *Nonlinear Multivariate Analysis*. Wiley.
- Greenacre, Michael J., and Jorg Blasius. 2006. *Multiple Correspondence Analysis and Related Methods*. Chapman; Hall.
- Guttman, Louis. 1941. “The Quantification of a Class of Attributes: A Theory and Method of Scale Construction.” In *The Prediction of Personal Adjustment*, edited by Paul Horst.

Social Science Research Council.

- Guttman, Louis. 1985. "Multidimensional Structuple Analysis (MSA-I) for the Classification of Cetacea: Whales, Porpoises and Dolphins." In *Data Analysis in Real Life Environment: Ins and Outs of Solving Problems*, edited by J.-F. Marcotorchino, J. M. Proth, and J. Janssen. North-Holland.
- Heiser, Willem J. 1981. "Unfolding Analysis of Proximity Data." PhD thesis, Leiden University.
- Heiser, Willem J., and Jan De Leeuw. 1979. "Metric Multidimensional Unfolding." *Methoden en Data Nieuwsbrief SWS/VVS* 4: 26–50. <https://jansweb.netlify.app/publication/heiser-deleeuw-a-79/heiser-deleeuw-a-79.pdf>.
- Husson, François, Julie Josse, and Gilbert Saporta. 2016. "Jan de Leeuw and the French School of Data Analysis." *Journal of Statistical Software* 73 (6): 1–14. <https://www.jstatsoft.org/article/view/v073i06>.
- Kruskal, Joseph B. 1964a. "Multidimensional Scaling by Optimizing Goodness of Fit to a Nonmetric Hypothesis." *Psychometrika* 29: 1–27.
- Kruskal, Joseph B. 1964b. "Nonmetric Multidimensional Scaling: a Numerical Method." *Psychometrika* 29: 115–29.
- Le Roux, Brigitte, and Henry Rouanet. 2010. *Multiple Correspondence Analysis*. Sage.
- Michailidis, George, and Jan De Leeuw. 2001. "Data Visualization through Graph Drawing." *Computational Statistics* 16: 435–50.
- Okabe, Atsuyuki, Barry Boots, Kokichi Sugihara, and Chiu Sung-Nok. 2000. *Spatial Tessellations: Concepts and Applications of Voronoi Diagrams*. Second Edition. Wiley.
- Peschar, Julien L. 1975. *School, Milieu, Beroep*. Tjeek Willink. <https://pure.rug.nl/ws/portalfiles/portal/14528340/peschar.PDF>.
- Tenenhaus, Michel, and Forest W. Young. 1985. "An Analysis and Synthesis of Multiple Correspondence Analysis, Optimal Scaling, Dual Scaling, Homogeneity Analysis and Other Methods for Quantifying Categorical Multivariate Data." *Psychometrika* 50: 91–119.
- Van der Burg, E. 1985. "HOMALS Classification of Whales, Porpoises and Dolphins." In *Data Analysis in Real Life Environment: Ins and Outs of Solving Problems*, edited by J.-F.

Marcotorchino, J. M. Proth, and J. Janssen. North-Holland.

Vescia, G. 1985. "Descriptive Classification of Cetacea: Whales, Porpoises and Dolphins." In *Data Analysis in Real Life Environment: Ins and Outs of Solving Problems*, edited by J.-F. Marcotorchino, J. M. Proth, and J. Janssen. North-Holland.