

Ćirić type fixed point theorems

Adrian Petruşel

Babeş-Bolyai University Cluj-Napoca
Faculty of Mathematics and Computer Science

January, 2014

- 1 Introduction
 - Abstract
- 2 Some known results
 - The single-valued case
 - The multi-valued case
- 3 Fixed point theorems in metric spaces endowed with a partial order
 - Basic concepts
 - Some abstract results
 - Fixed point theorems for φ -contractions
- 4 Fixed point theorems in metric spaces endowed with a graph
 - Basic concepts
 - The single-valued case
 - The multi-valued case
- 5 Main works

- 1 Introduction
 - Abstract
- 2 Some known results
 - The single-valued case
 - The multi-valued case
- 3 Fixed point theorems in metric spaces endowed with a partial order
 - Basic concepts
 - Some abstract results
 - Fixed point theorems for φ -contractions
- 4 Fixed point theorems in metric spaces endowed with a graph
 - Basic concepts
 - The single-valued case
 - The multi-valued case
- 5 Main works

The purpose of this talk is to present some fixed point and strict results for (single-valued and multi-valued) generalized contractions of Ćirić type.

The purpose of this talk is to present some fixed point and strict results for (single-valued and multi-valued) generalized contractions of Ćirić type.

Connections with other results given in:

L.B. Ćirić: *Generalized contraction and fixed point theorems*, Publ. Inst. Math., 12 (1971), 19-26.

L.B. Ćirić: *Generalized contraction and fixed point theorems*, Publ. Inst. Math., 12 (1971), 19-26.

L.B. Ćirić: *Fixed points for generalized multi-valued contractions*, Math. Vesnik, 9(24) (1972), 265-272.

L.B. Ćirić: *Generalized contraction and fixed point theorems*, Publ. Inst. Math., 12 (1971), 19-26.

L.B. Ćirić: *Fixed points for generalized multi-valued contractions*, Math. Vesnik, 9(24) (1972), 265-272.

L.B. Ćirić: *A generalization of Banach's contraction principle*, Proc. Amer. Math. Soc., 45 (1974), 267-273.

L.B. Ćirić: *Generalized contraction and fixed point theorems*, Publ. Inst. Math., 12 (1971), 19-26.

L.B. Ćirić: *Fixed points for generalized multi-valued contractions*, Math. Vesnik, 9(24) (1972), 265-272.

L.B. Ćirić: *A generalization of Banach's contraction principle*, Proc. Amer. Math. Soc., 45 (1974), 267-273.

L.B. Ćirić: *Contractive type non-self mappings on metric spaces of hyperbolic type*, J. Math. Anal. Appl., 317 (2006), 28-42.

L.B. Ćirić: *Generalized contraction and fixed point theorems*, Publ. Inst. Math., 12 (1971), 19-26.

L.B. Ćirić: *Fixed points for generalized multi-valued contractions*, Math. Vesnik, 9(24) (1972), 265-272.

L.B. Ćirić: *A generalization of Banach's contraction principle*, Proc. Amer. Math. Soc., 45 (1974), 267-273.

L.B. Ćirić: *Contractive type non-self mappings on metric spaces of hyperbolic type*, J. Math. Anal. Appl., 317 (2006), 28-42.

G.E. Hardy and T.D. Rogers: *A generalization of a fixed point theorem of Reich*, Canad. Math. Bull., 16(1973), 201-206.

S. Reich: *Fixed point of contractive functions*, Boll. Un. Mat. Ital., 5 (1972), 26-42.

L.B. Ćirić: *Generalized contraction and fixed point theorems*, Publ. Inst. Math., 12 (1971), 19-26.

L.B. Ćirić: *Fixed points for generalized multi-valued contractions*, Math. Vesnik, 9(24) (1972), 265-272.

L.B. Ćirić: *A generalization of Banach's contraction principle*, Proc. Amer. Math. Soc., 45 (1974), 267-273.

L.B. Ćirić: *Contractive type non-self mappings on metric spaces of hyperbolic type*, J. Math. Anal. Appl., 317 (2006), 28-42.

G.E. Hardy and T.D. Rogers: *A generalization of a fixed point theorem of Reich*, Canad. Math. Bull., 16(1973), 201-206.

S. Reich: *Fixed point of contractive functions*, Boll. Un. Mat. Ital., 5 (1972), 26-42.

I.A. Rus: *Generalized Contractions and Applications*, Transilvania Press, 2001.

I.A. Rus: *Generalized Contractions and Applications*, Transilvania Press, 2001.

I. Beg, A.R. Butt and S. Radojevic: *The contraction principle for set-valued mappings on a metric space with a graph*, Computers Math. Appl., 60(2010), 1214-1219.

I.A. Rus: *Generalized Contractions and Applications*, Transilvania Press, 2001.

I. Beg, A.R. Butt and S. Radojevic: *The contraction principle for set-valued mappings on a metric space with a graph*, Computers Math. Appl., 60(2010), 1214-1219.

A. Petrușel, G. Petrușel: Multivalued Picard operator, J. Nonlinear Convex Anal., 2012

I.A. Rus: *Generalized Contractions and Applications*, Transilvania Press, 2001.

I. Beg, A.R. Butt and S. Radojevic: *The contraction principle for set-valued mappings on a metric space with a graph*, Computers Math. Appl., 60(2010), 1214-1219.

A. Petruşel, G. Petruşel: Multivalued Picard operator, J. Nonlinear Convex Anal., 2012

T. Dinevari and M. Frigon: *Fixed point results for multivalued contractions on a metric space with a graph*, J. Math. Anal. Appl., 405(2013), 2, 507-517.

I.A. Rus: *Generalized Contractions and Applications*, Transilvania Press, 2001.

I. Beg, A.R. Butt and S. Radojevic: *The contraction principle for set-valued mappings on a metric space with a graph*, Computers Math. Appl., 60(2010), 1214-1219.

A. Petruşel, G. Petruşel: Multivalued Picard operator, J. Nonlinear Convex Anal., 2012

T. Dinevari and M. Frigon: *Fixed point results for multivalued contractions on a metric space with a graph*, J. Math. Anal. Appl., 405(2013), 2, 507-517.

C. Chifu, G. Petruşel and M. Bota, *Fixed points and strict fixed points for multivalued contractions of Reich type on metric spaces endowed with a graph*, Fixed Point Theory Appl. 2013, 2013:203, doi:10.1186/1687-1812-2013-203.

I.A. Rus: *Generalized Contractions and Applications*, Transilvania Press, 2001.

I. Beg, A.R. Butt and S. Radojevic: *The contraction principle for set-valued mappings on a metric space with a graph*, Computers Math. Appl., 60(2010), 1214-1219.

A. Petruşel, G. Petruşel: Multivalued Picard operator, J. Nonlinear Convex Anal., 2012

T. Dinevari and M. Frigon: *Fixed point results for multivalued contractions on a metric space with a graph*, J. Math. Anal. Appl., 405(2013), 2, 507-517.

C. Chifu, G. Petruşel and M. Bota, *Fixed points and strict fixed points for multivalued contractions of Reich type on metric spaces endowed with a graph*, Fixed Point Theory Appl. 2013, 2013:203, doi:10.1186/1687-1812-2013-203.

- 1 Introduction
 - Abstract
- 2 Some known results
 - The single-valued case
 - The multi-valued case
- 3 Fixed point theorems in metric spaces endowed with a partial order
 - Basic concepts
 - Some abstract results
 - Fixed point theorems for φ -contractions
- 4 Fixed point theorems in metric spaces endowed with a graph
 - Basic concepts
 - The single-valued case
 - The multi-valued case
- 5 Main works

- 1 Introduction
 - Abstract
- 2 Some known results
 - The single-valued case
 - The multi-valued case
- 3 Fixed point theorems in metric spaces endowed with a partial order
 - Basic concepts
 - Some abstract results
 - Fixed point theorems for φ -contractions
- 4 Fixed point theorems in metric spaces endowed with a graph
 - Basic concepts
 - The single-valued case
 - The multi-valued case
- 5 Main works

- 1 Introduction
 - Abstract
- 2 Some known results
 - The single-valued case
 - The multi-valued case
- 3 Fixed point theorems in metric spaces endowed with a partial order
 - Basic concepts
 - Some abstract results
 - Fixed point theorems for φ -contractions
- 4 Fixed point theorems in metric spaces endowed with a graph
 - Basic concepts
 - The single-valued case
 - The multi-valued case
- 5 Main works

- 1 Introduction
 - Abstract
- 2 Some known results
 - The single-valued case
 - The multi-valued case
- 3 Fixed point theorems in metric spaces endowed with a partial order
 - Basic concepts
 - Some abstract results
 - Fixed point theorems for φ -contractions
- 4 Fixed point theorems in metric spaces endowed with a graph
 - Basic concepts
 - The single-valued case
 - The multi-valued case
- 5 Main works

- 1 Introduction
 - Abstract
- 2 Some known results
 - The single-valued case
 - The multi-valued case
- 3 Fixed point theorems in metric spaces endowed with a partial order
 - Basic concepts
 - Some abstract results
 - Fixed point theorems for φ -contractions
- 4 Fixed point theorems in metric spaces endowed with a graph
 - Basic concepts
 - The single-valued case
 - The multi-valued case
- 5 Main works

- **Definition.** Let X be a nonempty set. Then, by definition (X, \rightarrow, \leq) is an ordered L-space if and only if:
- (i) (X, \rightarrow) is an L-space;
 - (ii) (X, \leq) is a partially ordered set;
 - (iii) $(x_n)_{n \in \mathbb{N}} \rightarrow x$, $(y_n)_{n \in \mathbb{N}} \rightarrow y$ and $x_n \leq y_n$, for each $n \in \mathbb{N} \Rightarrow x \leq y$.
- If (X, d) is a metric space, then the triple (X, d, \leq) will be called an ordered metric space.

- ▶ Let (X, \leq) be a partially ordered set.

Denote

$$X_{\leq} := \{(x, y) \in X \times X \mid x \leq y \text{ or } y \leq x\}.$$

- ▶ Let (X, \leq) be a partially ordered set.

Denote

$$X_{\leq} := \{(x, y) \in X \times X \mid x \leq y \text{ or } y \leq x\}.$$

- ▶ In the same setting, consider $f : X \rightarrow X$. Then:
 $(LF)_f := \{x \in X \mid x \leq f(x)\}$ is the lower fixed point set of f ,
 $(UF)_f := \{x \in X \mid x \geq f(x)\}$ is the upper fixed point set of f .

- ▶ Let (X, \leq) be a partially ordered set.

Denote

$$X_{\leq} := \{(x, y) \in X \times X \mid x \leq y \text{ or } y \leq x\}.$$

- ▶ In the same setting, consider $f : X \rightarrow X$. Then:
 $(LF)_f := \{x \in X \mid x \leq f(x)\}$ is the lower fixed point set of f ,
 $(UF)_f := \{x \in X \mid x \geq f(x)\}$ is the upper fixed point set of f .
- ▶ If $f : X \rightarrow X$ and $g : Y \rightarrow Y$, then the cartesian product of f and g is denoted by $f \times g$ and it is defined in the following way:

$$f \times g : X \times Y \rightarrow X \times Y, (f \times g)(x, y) := (f(x), g(y)).$$

Definition. (I.A. Rus) Let (X, \rightarrow) be an L-space.

An operator $f : X \rightarrow X$ is, by definition, a Picard operator if:

- (i) $F_f = \{x^*\}$;
- (ii) $(f^n(x))_{n \in \mathbb{N}} \rightarrow x^*$ as $n \rightarrow \infty$, for all $x \in X$.

Theorem. (Ran and Reurings-2004) *Let X be a partially ordered set such that every pair $x, y \in X$ has a lower and an upper bound. Let d be a metric on X such that the metric space (X, d) is complete. Let $f : X \rightarrow X$ be a continuous and monotone (i. e., either decreasing or increasing) operator. Suppose that the following two assertions hold:*

- 1) there exists $a \in]0, 1[$ such that $d(f(x), f(y)) \leq a \cdot d(x, y)$, for each $x, y \in X$ with $x \leq y$*
- 2) $(LF)_f \cup (UF)_f \neq \emptyset$.*

Then f is a Picard operator.

Theorem. (Nieto and Rodríguez-López, 2005) *Let X be a partially ordered set such that every pair $x, y \in X$ has a lower or an upper bound. Let d be a metric on X such that the metric space (X, d) is complete. Let $f : X \rightarrow X$ be an increasing operator. Suppose that the following two assertions hold:*

- 1) there exists $a \in]0, 1[$ such that $d(f(x), f(y)) \leq a \cdot d(x, y)$, for each $x, y \in X$ with $x \leq y$;*
- 2) there exists $x_0 \in X$ such that $x_0 \leq f(x_0)$;*
- 3) if an increasing sequence (x_n) converges to x in X , then $x_n \leq x$ for all $n \in \mathbb{N}$.*

Then f is a Picard operator.

- 1 Introduction
 - Abstract
- 2 Some known results
 - The single-valued case
 - The multi-valued case
- 3 Fixed point theorems in metric spaces endowed with a partial order
 - Basic concepts
 - **Some abstract results**
 - Fixed point theorems for φ -contractions
- 4 Fixed point theorems in metric spaces endowed with a graph
 - Basic concepts
 - The single-valued case
 - The multi-valued case
- 5 Main works

Lemma.

(A. Petruşel and Rus, 2006)

Let (X, \rightarrow) be an L -space and U a symmetric subset of $X \times X$ such that $\Delta(X) \subset U$. Let $f : X \rightarrow X$ be an operator. Suppose that:

- (i) for each $x, y \in X$ with $(x, y) \notin U$ there exists $z \in X$ such that $(x, z) \in U$ and $(y, z) \in U$;
- (ii) there exist $x_0, x^* \in X$ such that $x_0 \in A_f(x^*)$;
- (iii) $(x, y) \in U$ and $x \in A_f(x^*)$ implies $y \in A_f(x^*)$.

Then $A_f(x^*) = X$.

Moreover, if f is orbitally continuous, then f is a Picard operator.

where

$$A_f(x^*) := \{x \in X : (f^n(x))_{n \in \mathbb{N}} \rightarrow x^* \text{ as } n \rightarrow \infty\}.$$

A natural consequence of the above result follows by choosing $U := X_{\leq}$.

Lemma.

(A. Petruşel and Rus, 2006)

Let (X, \rightarrow, \leq) be an ordered L -space and $f : X \rightarrow X$ be an operator.

Suppose that:

- (i) for each $x, y \in X$ with $(x, y) \notin X_{\leq}$ there exists $z \in X$ such that $(x, z) \in X_{\leq}$ and $(y, z) \in X_{\leq}$;
- (ii) there exist $x_0, x^* \in X$ such that $x_0 \in A_f(x^*)$;
- (iii) $(x, y) \in X_{\leq}$ and $x \in A_f(x^*)$ implies $y \in A_f(x^*)$;
- (iv) f is orbitally continuous

Then f is a Picard operator.

- 1 Introduction
 - Abstract
- 2 Some known results
 - The single-valued case
 - The multi-valued case
- 3 Fixed point theorems in metric spaces endowed with a partial order
 - Basic concepts
 - Some abstract results
 - Fixed point theorems for φ -contractions
- 4 Fixed point theorems in metric spaces endowed with a graph
 - Basic concepts
 - The single-valued case
 - The multi-valued case
- 5 Main works

Theorem.

(O'Regan and A. Petruşel, 2008)

Let (X, d, \leq) be an ordered metric space and $f : X \rightarrow X$ be an operator.

We suppose that:

(i) For each $x, y \in X$ with $(x, y) \notin X_{\leq}$ there exists $c(x, y) \in X$ such that $(x, c(x, y)) \in X_{\leq}$ and $(y, c(x, y)) \in X_{\leq}$;

(ii) $f : (X, \leq) \rightarrow (X, \leq)$ is increasing;

(iii) there exists $x_0 \in X$ such that $x_0 \leq f(x_0)$;

(iv)_a f is orbitally continuous

or

(iv)_b if an increasing sequence (x_n) converges to x in X , then $x_n \leq x$ for all $n \in \mathbb{N}$;

(v) there exists a comparison function $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that $d(f(x), f(y)) \leq \varphi(d(x, y))$, for each $(x, y) \in X_{\leq}$;

(vi) the metric d is complete.

Then f is a Picard operator.

1 Introduction

- Abstract

2 Some known results

- The single-valued case
- The multi-valued case

3 Fixed point theorems in metric spaces endowed with a partial order

- Basic concepts
- Some abstract results
- Fixed point theorems for φ -contractions

4 Fixed point theorems in metric spaces endowed with a graph

- Basic concepts
- The single-valued case
- The multi-valued case

5 Main works

- 1 Introduction
 - Abstract
- 2 Some known results
 - The single-valued case
 - The multi-valued case
- 3 Fixed point theorems in metric spaces endowed with a partial order
 - Basic concepts
 - Some abstract results
 - Fixed point theorems for φ -contractions
- 4 Fixed point theorems in metric spaces endowed with a graph
 - Basic concepts
 - The single-valued case
 - The multi-valued case
- 5 Main works

Let (X, d) be a metric space and Δ the diagonal of $X \times X$.

Let (X, d) be a metric space and Δ the diagonal of $X \times X$.

Let G be a directed graph such that one can identify G with the pair $(V(G), E(G))$, where the set $V(G)$ of its vertices coincides with X and the set $E(G)$ of the edges of the graph contains Δ .

Let (X, d) be a metric space and Δ the diagonal of $X \times X$.

Let G be a directed graph such that one can identify G with the pair $(V(G), E(G))$, where the set $V(G)$ of its vertices coincides with X and the set $E(G)$ of the edges of the graph contains Δ .

If x and y are vertices of G , then a path in G from x to y of length $k \in \mathbb{N}$ is a finite sequence $(x_n)_{n \in \{0, 1, 2, \dots, k\}}$ of vertices such that

$$x_0 = x, x_k = y \text{ and } (x_{i-1}, x_i) \in E(G), \text{ for } i \in \{1, 2, \dots, k\}.$$

Let (X, d) be a metric space and Δ the diagonal of $X \times X$.

Let G be a directed graph such that one can identify G with the pair $(V(G), E(G))$, where the set $V(G)$ of its vertices coincides with X and the set $E(G)$ of the edges of the graph contains Δ .

If x and y are vertices of G , then a path in G from x to y of length $k \in \mathbb{N}$ is a finite sequence $(x_n)_{n \in \{0, 1, 2, \dots, k\}}$ of vertices such that

$$x_0 = x, x_k = y \text{ and } (x_{i-1}, x_i) \in E(G), \text{ for } i \in \{1, 2, \dots, k\}.$$

A graph G is connected if there is a path between any two vertices and it is weakly connected if \tilde{G} is connected, where \tilde{G} denotes the undirected graph obtained from G by ignoring the direction of edges.

Definition. (J. Jachymski-2008)

An operator $f : X \rightarrow X$ is called a Banach G -contraction if and only if:

Definition. (J. Jachymski-2008)

An operator $f : X \rightarrow X$ is called a Banach G -contraction if and only if:

- (a) f is edge preserving, i.e., for each $x, y \in X$ with $(x, y) \in E(G)$ we have $(f(x), f(y)) \in E(G)$;

Definition. (J. Jachymski-2008)

An operator $f : X \rightarrow X$ is called a Banach G -contraction if and only if:

- (a) f is edge preserving, i.e., for each $x, y \in X$ with $(x, y) \in E(G)$ we have $(f(x), f(y)) \in E(G)$;
- (b) there exists $\alpha \in]0, 1[$ such that for each $x, y \in X$ the following implication holds:

$$(x, y) \in E(G) \text{ implies } d(f(x), f(y)) \leq \alpha d(x, y).$$

Definition. (J. Jachymski-2008)

An operator $f : X \rightarrow X$ is called a Banach G -contraction if and only if:

- (a) f is edge preserving, i.e., for each $x, y \in X$ with $(x, y) \in E(G)$ we have $(f(x), f(y)) \in E(G)$;
- (b) there exists $\alpha \in]0, 1[$ such that for each $x, y \in X$ the following implication holds:

$$(x, y) \in E(G) \text{ implies } d(f(x), f(y)) \leq \alpha d(x, y).$$

Examples:

- 1) Any Banach contraction is a G_0 -contraction, where the graph G_0 is defined by $E(G_0) := X \times X$.

Definition. (J. Jachymski-2008)

An operator $f : X \rightarrow X$ is called a Banach G -contraction if and only if:

- (a) f is edge preserving, i.e., for each $x, y \in X$ with $(x, y) \in E(G)$ we have $(f(x), f(y)) \in E(G)$;
- (b) there exists $\alpha \in]0, 1[$ such that for each $x, y \in X$ the following implication holds:

$$(x, y) \in E(G) \text{ implies } d(f(x), f(y)) \leq \alpha d(x, y).$$

Examples:

- 1) Any Banach contraction is a G_0 -contraction, where the graph G_0 is defined by $E(G_0) := X \times X$.
- 2) Let \leq be a partial order in X . Define the graph G_1 by

$$E(G_1) := \{(x, y) \in X \times X : x \leq y\}.$$

Definition. (J. Jachymski-2008)

An operator $f : X \rightarrow X$ is called a Banach G -contraction if and only if:

- (a) f is edge preserving, i.e., for each $x, y \in X$ with $(x, y) \in E(G)$ we have $(f(x), f(y)) \in E(G)$;
- (b) there exists $\alpha \in]0, 1[$ such that for each $x, y \in X$ the following implication holds:

$$(x, y) \in E(G) \text{ implies } d(f(x), f(y)) \leq \alpha d(x, y).$$

Examples:

1) Any Banach contraction is a G_0 -contraction, where the graph G_0 is defined by $E(G_0) := X \times X$.

2) Let \leq be a partial order in X . Define the graph G_1 by

$$E(G_1) := \{(x, y) \in X \times X : x \leq y\}.$$

3) Let \leq be a partial order in X . Define the graph G_2 by

$$E(G_2) := \{(x, y) \in X \times X : x \leq y \text{ or } y \leq x\}.$$

- 1 Introduction
 - Abstract
- 2 Some known results
 - The single-valued case
 - The multi-valued case
- 3 Fixed point theorems in metric spaces endowed with a partial order
 - Basic concepts
 - Some abstract results
 - Fixed point theorems for φ -contractions
- 4 Fixed point theorems in metric spaces endowed with a graph
 - Basic concepts
 - The single-valued case
 - The multi-valued case
- 5 Main works

Theorem. (J. Jachymski-2008)

Let (X, d) be a complete metric space and let G be a directed graph G such that the triple (X, d, G) has property (P):

for any sequence $(x_n)_{n \in \mathbb{N}} \subset X$, if $x_n \rightarrow x$ as $n \rightarrow +\infty$ and
 (P) $(x_n, x_{n+1}) \in E(G)$, for each $n \in \mathbb{N}$, then there exists a subsequence
 $(x_{k_n})_{n \in \mathbb{N}}$ of $(x_n)_{n \in \mathbb{N}}$ such that $(x_{k_n}, x) \in E(G)$, for each $n \in \mathbb{N}$.

Let $f : X \rightarrow X$ be a G -contraction. Then the following statements hold:

1) $F_f \neq \emptyset$ if and only if $X_f \neq \emptyset$,

where $X_f := \{x \in X : (x, f(x)) \in E(G)\}$;

2) if $X_f \neq \emptyset$ and G is weakly connected, then f is a Picard operator.

- 1 Introduction
 - Abstract
- 2 Some known results
 - The single-valued case
 - The multi-valued case
- 3 Fixed point theorems in metric spaces endowed with a partial order
 - Basic concepts
 - Some abstract results
 - Fixed point theorems for φ -contractions
- 4 Fixed point theorems in metric spaces endowed with a graph
 - Basic concepts
 - The single-valued case
 - The multi-valued case
- 5 Main works

Theorem.

(Nicolae-O'Regan-A. Petruşel, 2011)

Let (X, d) be a complete metric space and G be a directed graph such that the triple (X, d, G) satisfy property (P) . Let $T : X \rightarrow P_{cl}(X)$ be a multi-valued operator. Suppose the following assertions hold:

(i) there exists $\alpha \in (0, 1)$ such that

$$H(T(x), T(y)) \leq \alpha d(x, y) \text{ for all } (x, y) \in E(G).$$

(ii) for each $(x, y) \in E(G)$, each $u \in T(x)$ and $v \in T(y)$ satisfying the condition $d(u, v) \leq \alpha d(x, y)$, for some $\alpha \in (0, 1)$, we have $(u, v) \in E(G)$; Then $F_T \neq \emptyset$ if and only if $X_T \neq \emptyset$, where

$$X_T := \{x \in X; \text{ there exists } y \in T(x) \text{ such that } (x, y) \in E(G)\},$$

- 1 Introduction
 - Abstract
- 2 Some known results
 - The single-valued case
 - The multi-valued case
- 3 Fixed point theorems in metric spaces endowed with a partial order
 - Basic concepts
 - Some abstract results
 - Fixed point theorems for φ -contractions
- 4 Fixed point theorems in metric spaces endowed with a graph
 - Basic concepts
 - The single-valued case
 - The multi-valued case

5 Main works

This part is based on:

This part is based on:

A.C.M. Ran, M.C. Reurings:

A fixed point theorem in partially ordered sets and some applications to matrix equations, *Proc. Amer. Math. Soc.* 132(2004) 1435-1443.

This part is based on:

A.C.M. Ran, M.C. Reurings:

A fixed point theorem in partially ordered sets and some applications to matrix equations, *Proc. Amer. Math. Soc.* 132(2004) 1435-1443.

J.J. Nieto, R. Rodríguez-López:

Contractive mapping theorems in partially ordered sets and applications to ordinary differential equations, *Order* 22(2005) 223-239.

This part is based on:

A.C.M. Ran, M.C. Reurings:

A fixed point theorem in partially ordered sets and some applications to matrix equations, *Proc. Amer. Math. Soc.* 132(2004) 1435-1443.

J.J. Nieto, R. Rodríguez-López:

Contractive mapping theorems in partially ordered sets and applications to ordinary differential equations, *Order* 22(2005) 223-239.

A. Petrușel, I.A. Rus:

Fixed point theorems in ordered L -spaces, *Proc. Amer. Math. Soc.* 134(2006) 411-418.

This part is based on:

A.C.M. Ran, M.C. Reurings:

A fixed point theorem in partially ordered sets and some applications to matrix equations, *Proc. Amer. Math. Soc.* 132(2004) 1435-1443.

J.J. Nieto, R. Rodríguez-López:

Contractive mapping theorems in partially ordered sets and applications to ordinary differential equations, *Order* 22(2005) 223-239.

A. Petruşel, I.A. Rus:

Fixed point theorems in ordered L -spaces, *Proc. Amer. Math. Soc.* 134(2006) 411-418.

D. O'Regan, A. Petruşel:

Fixed point theorems in ordered metric spaces, *J. Math. Anal. Appl.* 341(2008) 1241-1252.

J. Jachymski:

The contraction principle for mappings on a metric space with a graph,
Proc. Amer. Math. Soc. 136(2008) 1359-1373.

J. Jachymski:

The contraction principle for mappings on a metric space with a graph, *Proc. Amer. Math. Soc.* 136(2008) 1359-1373.

A. Nicolae, D. O'Regan, A. Petruşel:

Fixed point theorems for singlevalued and multivalued generalized contractions in metric spaces endowed with a graph, *J. Georgian Math. Soc.*, 18(2011), 307327.