

Rscience

Rscience

2026-01-06

Table of contents

1	Anova Output	2
1.1	Anova Table	2
1.2	Tukey Table and Plot	2
1.3	Requeriments - Residuals Normality	4
1.4	Requeriments - Residuals Homogeneity	4
2	Reporting Level I - Summary Anova	5
2.1	Summary Table	5
2.2	Phrases	5
3	Reporting Level II - Decision-Making Process in ANOVA	6
3.1	Step 01: Normality Verification	6
3.2	Step 02: Homogeneity Verification	6
3.3	Step 03: Global Requirements Check	7
3.4	Step 04: ANOVA Validation	7
3.5	Step 05: ANOVA Interpretation	7
3.6	Step 06: Tukey Test Validation	7
3.7	Step 07: Tukey Test Interpretation	8
3.8	Step 08: ANOVA-Tukey Consistency Analysis	9
4	Reporting Level III - Automatic Statistic Asesor (ASA)	10
4.1	General	10
4.2	Requeriments	10
4.3	Anova	10
4.4	Tukey	11

1 Anova Outpus


1.1 Anova Table

	Sources of Variation	Df	Sum Sq	Mean Sq	F value	Pr(>F)
cyl	Factor	2	824.7846	412.39229	39.69752	0
Residuals	Error	29	301.2626	10.38837	NA	NA
Total	Total	31	1126.0472	36.32410	NA	NA


1.2 Tukey Table and Plot

order	level	mean	group
1	4	26.66364	a
2	6	19.74286	b
3	8	15.10000	c

Plot 007 - Mean y model standard error



Plot 007 - Mean y model standard error



1.3 Requeriments - Residuals Normality

variable	test	statistic(W)	p_value
residuals	Shapiro-Wilk normality test	0.9706547	0.517665

1.4 Requeriments - Residuals Homogeneity

variable	test	statistic(Bartlett's K-squared)	p_value
residuals	Bartlett test of homogeneity of variances	8.393396	0.0150452

2 Reporting Level I - Summary Anova

2.1 Summary Table

name	test	variable	p_value	alpha_val	Decision	Valid
Shapiro-Wilk test	Normality	residuals	0.5176650	0.05	Ho no rejected	Yes, it is.
Bartlett test	Homogeneity	residuals	0.0150452	0.05	Ho Rejected	Yes, it is.
Anova 1 way	Analysis of Variance	mpg	0.0000000	0.05	Ho Rejected	No, is not!!!
Tukey	Mean groups	mpg	NA	0.05	At least two groups of means.	No, is not!!!

2.2 Phrases

The null hypothesis of normal distribution of residuals is not rejected.

The hypothesis of homogeneity of variances (homoscedasticity) is rejected.

Not all requirements for the model are met.

It is NOT valid to draw conclusions from the ANOVA test.

As the model requirements are not met, the ANOVA and Tukey analyses must be discarded, irrespective of the statistical values obtained. The literal and decontextualized interpretation of the ANOVA test and the Tukey test is detailed below for demonstrative purposes only, but holds no validity for drawing conclusions from them.

The null hypothesis of the ANOVA test is rejected. There are statistically significant differences in at least one level of the factor.

Since the ANOVA null hypothesis is rejected, the use of a multiple comparison test to accompany the ANOVA test is valid.

The Tukey test is selected as the multiple comparison test for this script.

The Tukey test indicates that there are at least 2 groups among the factor levels. The number of groups and their structure must be analyzed in detail using the Tukey table.

ANOVA indicates that all factor levels are equal, while Tukey has found at least 2 groups. Tukey is subject to the rejection of the ANOVA hypothesis. Since the ANOVA null hypothesis is not rejected, the Tukey test must not be taken into account for decision-making. All factor levels are equal.

3 Reporting Level II - Decision-Making Process in ANOVA

The ANOVA test is specifically performed to make decisions.

The decision-making process has a set of sequential steps, in a particular order.

Step 01: Normality Verification

Step 02: Homogeneity Verification

Step 03: Global Requirements Check

Step 04: ANOVA Validation

Step 05: ANOVA Interpretation

Step 06: Tukey Test Validation

Step 07: Tukey Test Interpretation

Step 08: ANOVA-Tukey Consistency Analysis

Step 09: Final Conclusion

3.1 Step 01: Normality Verification

variable	test	statistic(W)	p_value
residuals	Shapiro-Wilk normality test	0.9706547	0.517665

The p value about normality distribution is 0.517664951040294.

Alpha value is 0.05.

The p value is equal or mayor than alpha value.

The null hypothesis of normal distribution of residuals is not rejected.

Los residuos poseen distribución normal.

3.2 Step 02: Homogeneity Verification

variable	test	statistic(Bartlett's K-squared)	p_value
residuals	Bartlett test of homogeneity of variances	8.393396	0.0150452

The p value about homogeneity variances from residuals is 0.0150451770493758.

Alpha value is 0.05.

The p value is less than alpha value.

The null hypothesis of homogeneity of variances from residuals is rejected.

Los residuos son homogéneos.

3.3 Step 03: Global Requirements Check

Not all requirements for the model are met.

3.4 Step 04: ANOVA Validation

It is NOT valid to draw conclusions from the ANOVA test. This dataset must be analyzed with another statistical tool such as the Kruskal-Wallis test. If a more sophisticated tool is desired, the possibility of utilizing generalized linear mixed models, generalized linear models, generalized linear models, exact distribution linear models, etc., could be evaluated.

It is NOT valid to draw conclusions from the ANOVA test. As the model requirements are not met, the ANOVA and Tukey analyses must be discarded, irrespective of the statistical values obtained. The literal and decontextualized interpretation of the ANOVA test and the Tukey test is detailed below for demonstrative purposes only, but holds no validity for drawing conclusions from them.

3.5 Step 05: ANOVA Interpretation

	Sources of Variation	Df	Sum Sq	Mean Sq	F value	Pr(>F)
cyl	Factor	2	824.7846	412.39229	39.69752	0
Residuals	Error	29	301.2626	10.38837	NA	NA
Total	Total	31	1126.0472	36.32410	NA	NA

The p-value for the ANOVA test is 4.97891917440017e-09. The alpha value is 0.05. The p-value is less than the alpha value. The null hypothesis of equal means for the ANOVA test is rejected. There are statistically significant differences in at least one level of the factor. By rejecting the null hypothesis, the ANOVA test guarantees statistically significant differences in at least one level of the factor with the lowest mean and the level of the factor with the highest mean.

3.6 Step 06: Tukey Test Validation

Since the ANOVA null hypothesis is rejected, the use of a multiple comparison test to accompany the ANOVA test is valid.

The Tukey test is selected as the multiple comparison test for this script.

3.7 Step 07: Tukey Test Interpretation

The Tukey test details at least 2 statistically different groups. The group structure detailed by the Tukey test must now be considered in order to provide a recommendation in your area of work.

order	level	mean	group
1	4	26.66364	a
2	6	19.74286	b
3	8	15.10000	c

```
Error in `initialize()`:
! `width` must be a whole number, not the string "100%".
```


3.8 Step 08: ANOVA-Tukey Consistency Analysis

ANOVA indicates that all factor levels are equal, while Tukey has found at least 2 groups. Tukey is subject to the rejection of the ANOVA hypothesis. Since the ANOVA null hypothesis is not rejected, the Tukey test must not be taken into account for decision-making. All factor levels are equal.

The means of each factor level are detailed with the standard error derived from the model. The statistical groups from the Tukey test are detailed for each factor level.

4 Reporting Level III - Automatic Statistic Asesor (ASA)

4.1 General

When not all model requirements (Normality and/or Homogeneity) are met, the ANOVA analysis is immediately discarded. This invalidation occurs irrespective of whether the failure is due to non-normality of the residuals, non-homogeneity, or both requirements. Indistinctly of the statistical values obtained in the ANOVA test and the Tukey test, it is NOT valid to draw conclusions of any type; the entire analysis is completely discarded. An alternative statistical tool or path must be sought to analyze the data, with the quickest recommended tool being the Kruskal-Wallis test, although a better analysis may allow for the use of more robust tools such as Generalized Linear Mixed Models or Exact Statistics; the alternative statistical path is to perform a transformation on the response variable and attempt the 1-Factor ANOVA test again. The statistical options suggested must be verified and justified for their feasibility of use. Continuing with statistical interpretations without fulfilling the model's requirements is a serious error common among those new to statistics, and a common practice among those lacking the sufficient statistical knowledge to correctly analyze the reality they intend to study. If, despite all these warnings, you continue with the analysis, you expose your work to severe criticism from colleagues, evaluation committees, and publication reviewers.

4.2 Requeriments

The ANOVA statistical test is based on 3 points for decision-making: shape, dispersion, and position. The shape is the normal distribution of the residuals. The dispersion is the homogeneous variances in the residuals, and the position is the mean of the response variable for one of the factor levels. By guaranteeing that the shape and dispersion of the residuals have a certain particular structure, ANOVA manages to test the position in the response variable. This is why the fulfillment of normality and homogeneity of variances of the residuals is a requirement, and only with the simultaneous fulfillment of both requirements is it valid to make an interpretation of the means. If all requirements are not met simultaneously, the ANOVA analysis must be discarded, irrespective of the results it may have yielded. There is no escape, and you must seek another statistical tool to interpret these results.

4.3 Anova

The model requirements are not met. The ANOVA statistical test has the indispensable requirement of residual normality and homogeneity of variances. Both requirements must be met jointly for it to be valid to draw conclusions about the factor levels. Continuing with the statistical interpretation without fulfilling all the requirements is a catastrophic error. The ANOVA model requirements are neither optional nor can they be assumed. The model requirements are the fundamental basis upon which the ANOVA test relies to make decisions

about the means of the factor levels. In this case, you must seek another statistical tool to perform the analysis of your data.

4.4 Tukey

The use of the Tukey test is subject to the validity of the ANOVA model and the rejection of the ANOVA hypothesis. As the model requirements are not met, the ANOVA test is not valid, and therefore the Tukey test is not valid either. The information related to the Tukey test must be discarded. As detailed before, in this context, continuing with the statistical interpretation is a serious statistical error that demonstrates a total lack of statistical criteria for decision-making.