

# Introduction of SAV method

Liang Zeng

February 21, 2019

# Contents

Analysis of numerical methods

Introduction of Gradient Flow

Different method to solve Gradient Flow

SAV Versions of Several Discrete Formats

Advantage and Disadvantage of SAV

Numerical result

Improvement of SAV

Reference

# Convergence ,Error, Existence and Uniqueness of solution

- ▶ Linear equation:  
Lax Theorem: Constancy = Convergence  
when Compatibility
- ▶ Nonlinear equation: ?

# Physical property

Dissipation: Energy analysis method  
Other conservative properties

# Computational overhead

linear?

Constant/variable coefficients?

Parallelism?

# Introduction of Gradient flows

A gradient flow is determined not only by the driving free energy, but also the dissipation mechanism. Given a free energy functional  $E[\phi(x)]$  bounded from below. Denote its variational derivative as  $\mu = \delta E / \delta \phi$ . The general form of the gradient flow can be written as

$$\frac{\partial \phi}{\partial t} = G\mu, \quad (1)$$

supplemented with suitable boundary conditions. To simplify the presentation, we assume throughout the paper that the boundary conditions are chosen such that all boundary terms will vanish when integrating by parts are performed. This is true with periodic boundary conditions or homogeneous Neumann boundary conditions.

# Convex splitting approach

A very popular approach for gradient flow is the so called convex splitting method which appears to be introduced by [1] and popularized by [2]. Assuming the free energy density  $F(\phi)$  can be split as the difference of two convex functions, namely,  $F(\phi) = F_c(\phi) - F_e(\phi)$  with  $F_c''(\phi), F_e''(\phi) \geq 0$ .

Then, the first-order convex splitting scheme reads:

$$\begin{aligned} \frac{\phi^{n+1} - \phi^n}{\delta t} &= \delta \mu^{n+1}, \\ \mu^{n+1} &= -\Delta \phi^{n+1} + (F'_c(\phi^{n+1}) - F'_e(\phi^n)). \end{aligned} \tag{2}$$

# Convex splitting approach

One can easily show that the above scheme is unconditionally energy stable in the sense that

$$E(\phi^{n+1}) - E(\phi^n) \leq -\delta t \|\nabla \mu^{n+1}\|^2.$$



## Stabilized approach

Another widely used approach is the stabilized scheme, introduced in [3] (see also [4]). The main idea is to introduce an artificial stabilization term to balance the explicit treatment of the nonlinear term. A first-order stabilized scheme for 1 reads:

$$\begin{aligned}\frac{1}{\delta t}(\phi^{n+1} - \phi^n) &= \Delta \mu^{n+1}, \\ \mu^{n+1} &= -\Delta \phi^{n+1} + S(\phi^{n+1} - \phi^n) + F\end{aligned}\tag{3}$$

where  $S$  is a suitable stabilization parameter. It is shown in [18] that, under the assumption  $\|F''(\phi)\|_\infty \leq L$ , the above scheme is unconditionally stable for all  $S \geq L_2$ .

# Invariant energy quadratization (IEQ) approaches

The approach is proposed in [5] for dealing with AllenCahn and CahnHilliard equations with double well free energy. It is based on a Lagrange multiplier approach introduced in [6].

Very recently, X. Yang and his collaborators [7], [8], [9], [10], [11], [12], [13] made a big leap in generalizing the Lagrange multiplier approach to the so called invariant energy quadratization (IEQ) approach which is applicable to a large class of free energies.

Assuming that there exists  $C_0 \geq 0$  such that  $F(\phi) \geq -C_0$ , one then introduces a Lagrange multiplier (auxiliary variable)  $q(t, x; \phi) = \sqrt{F(\phi) + C_0}$ , and rewrite the equation as

# Invariant energy quadratization (IEQ) approaches

$$\begin{aligned}\phi_t &= \Delta\mu, \\ \mu &= -\Delta\phi + \frac{q}{\sqrt{F(\phi) + C_0}} F'(\phi), \\ q_t &= \frac{F'(\phi)}{2\sqrt{F(\phi) + C_0}} \phi_t.\end{aligned}\tag{4}$$

Taking the inner products of the above with  $\mu$ ,  $\phi_t$  and  $2q$ , respectively, we see that the above system satisfies a modified energy dissipation law:

$$\frac{d}{dt} \left( \frac{1}{2} \|\nabla\phi\|^2 + \int_{\Omega} q^2 dx \right) = -\|\nabla\mu\|^2.$$

# Invariant energy quadratization (IEQ) approaches

The above formulation is amenable to simple and efficient numerical schemes. Consider for instance,

$$\begin{aligned}\frac{\phi^{n+1} - \phi^n}{\delta t} &= \Delta \mu^{n+1}, \\ \mu^{n+1} &= -\Delta \phi^{n+1} + \frac{q^{n+1}}{\sqrt{F(\phi^n) + C_0}} F'(\phi^n), \\ \frac{q^{n+1} - q^n}{\delta t} &= \frac{F'(\phi^n)}{2\sqrt{F(\phi^n) + C_0}} \frac{\phi^{n+1} - \phi^n}{\delta t}.\end{aligned}\tag{5}$$

## Invariant energy quadratization (IEQ) approaches

Taking the inner products of the above with  $\mu^{n+1}$ ,  $\frac{\phi^{n+1}-\phi^n}{\delta t}$  and  $2q^{n+1}$ , respectively, one obtains immediately:

$$\begin{aligned} \frac{1}{\delta t} \left[ \frac{1}{2} \|\nabla \phi^{n+1}\|^2 + \int_{\Omega} (q^{n+1})^2 dx - \frac{1}{2} \|\nabla \phi^n\|^2 - \int_{\Omega} (q^n)^2 dx \right. \\ \left. + \frac{1}{2} \|\nabla(\phi^{n+1} - \phi^n)\|^2 + \int_{\Omega} (q^{n+1} - q^n)^2 dx \right] = -\|\nabla \mu^{n+1}\|^2, \end{aligned} \quad (6)$$

which indicates that the above scheme is unconditionally stable with respect to the modified energy.

# The scalar auxiliary variable (SAV) approach

we now only assume  $E_1(\phi) := \int_{\Omega} F(\phi) dx$  is bounded from below, i.e.,  $E_1(\phi) \geq C_0$ , which is necessary for the free energy to be physically sound, and introduce a scalar auxiliary variable (SAV):

$$r(t) = \sqrt{E_1(\phi) + C_0}.$$

Then, (1) can be rewritten as:

$$\begin{aligned}\phi_t &= \Delta \mu, \\ \mu &= -\Delta \phi + \frac{r}{\sqrt{E_1[\phi] + C_0}} F'(\phi), \\ r_t &= \frac{1}{2\sqrt{E_1[\phi] + C_0}} \int_{\Omega} F'(\phi) \phi_t dx.\end{aligned}\tag{7}$$

# The scalar auxiliary variable (SAV) approach

Taking the inner products of the above with  $\mu$ ,  $\frac{\partial \phi}{\partial t}$  and  $2r$ , respectively, we obtain the modified energy dissipation law:

$$\frac{d}{dt} \left( \frac{1}{2} \|\nabla \phi\|^2 + r^2(t) \right) = -\|\nabla \mu\|^2.$$

We now construct a semi-implicit second-order BDF scheme for the above system.

# The scalar auxiliary variable (SAV) approach

$$\begin{aligned}\frac{3\phi^{n+1} - 4\phi^n + \phi^{n-1}}{2\delta t} &= \Delta\mu^{n+1}, \\ \mu^{n+1} &= -\Delta\phi^{n+1} + \frac{r^{n+1}}{\sqrt{E_1[\bar{\phi}^{n+1}] + C_0}} F'(\bar{\phi}^{n+1}), \\ \frac{3r^{n+1} - 4r^n + r^{n-1}}{2\delta t} &= \int_{\Omega} \frac{F'(\bar{\phi}^{n+1})}{2\sqrt{E_1[\bar{\phi}^{n+1}] + C_0}} \frac{3\phi^{n+1} - 4\phi^n + \phi^{n-1}}{2\delta t} dx.\end{aligned}\tag{8}$$

where  $\bar{\phi}^{n+1}$  is any explicit  $O(\delta t^2)$  approximation for  $\phi(t^{n+1})$ , which can be flexible according to the problem, and which we will specify in our numerical results.



We can eliminate  $\mu^{n+1}$  and  $r^{n+1}$  from (8) to obtain

$$\begin{aligned} \frac{3\phi^{n+1} - 4\phi^n + \phi^{n-1}}{2\delta t} = & -\Delta^2 \phi^{n+1} + \frac{\Delta F'(\bar{\phi}^{n+1})}{3\sqrt{E_1[\bar{\phi}^{n+1}] + C_0}}(4r^n \\ & - r^{n-1} + \int_{\Omega} \frac{F'(\bar{\phi}^{n+1})}{2\sqrt{E_1[\bar{\phi}^{n+1}] + C_0}}(3\phi^{n+1} - 4\phi^n + \phi^{n-1})dx). \end{aligned} \quad (9)$$

Denote

$$\begin{aligned} b^n &= \frac{F'(\bar{\phi}^{n+1})}{\sqrt{E_1[\bar{\phi}^{n+1}] + C_0}}, \\ A &= I + (2\delta t/3)\Delta^2 \\ g^n &= \frac{1}{3}(4\phi^n - 3\phi^{n-1}) \\ &+ \frac{2\delta t}{9}[4r^n - r^{n-1} - \frac{1}{2}(b^n, 4\phi^n - \phi^{n-1})]\Delta b^n \end{aligned} \quad (10)$$

We can get

$$A\phi^{n+1} - \frac{\delta t}{3}(b^n, \phi^{n+1})\Delta b^n = g^n, \quad (11)$$

Then

$$(b^n, \phi^{n+1}) + \frac{\delta t}{3}\gamma^n(b^n, \phi^{n+1}) = (b^n, A^{-1}g^n),, \quad (12)$$

where  $\gamma = -(b^n, A^{-1}\Delta b^n) \geq 0$ , so

$$(b^n, \phi^{n+1}) = \frac{(b^n, A^{-1}g^n)}{1 + \delta t\gamma^n/3}. \quad (13)$$

Finally, we can solve  $\phi^{n+1}$  from (11).

## Unconditional energy stability of SAV/BDF2

The scheme (8) is second-order accurate, unconditionally energy stable in the sense that

$$\begin{aligned} & \frac{1}{\Delta t} \{ \tilde{E}[(\phi^{n+1}, r^{n+1}), (\phi^n, r^n)] - \tilde{E}[(\phi^n, r^n), (\phi^{n-1}, r^{n-1})] \} \\ & + \frac{1}{\Delta t} \left\{ \frac{1}{4} (\phi^{n+1} - 2\phi^n + \phi^{n-1}, -\Delta(\phi^{n+1} - 2\phi^n + \phi^{n-1})) \right. \\ & \left. + \frac{1}{2} (r^{n+1} - 2r^n + r^{n-1})^2 \right\} = (\mu, \Delta\mu), \end{aligned} \quad (14)$$

where the modified discrete energy is defined as

$$\begin{aligned} \tilde{E}[(\phi^{n+1}, r^{n+1}), (\phi^n, r^n)] &= \frac{1}{4} ((\phi^{n+1}, -\Delta\phi^{n+1}) \\ &+ (2\phi^{n+1} - \phi^n, -\Delta(2\phi^{n+1} - \phi^n))) + \frac{1}{2} ((r^{n+1})^2 + (2r^{n+1} - r^n)^2), \end{aligned} \quad (15)$$

## SAV/Crank-Nicolson

A semi-implicit second-order SAV scheme based on CrankNicolson is as follows:

$$\begin{aligned}\frac{\phi^{n+1} - \phi^n}{\delta t} &= \Delta \mu^{n+1/2}, \\ \mu^{n+1/2} &= -\Delta \frac{1}{2}(\phi^{n+1} + \phi^n) + \frac{r^{n+1} + r^n}{2\sqrt{E_1[\bar{\phi}^{n+1/2}] + C_0}} F'(\bar{\phi}^{n+1/2}), \\ \frac{r^{n+1} - r^n}{\delta t} &= \int_{\Omega} \frac{F'(\bar{\phi}^{n+1/2})}{2\sqrt{E_1[\bar{\phi}^{n+1/2}] + C_0}} \frac{\phi^{n+1} - \phi^n}{\delta t} dx.\end{aligned}\tag{16}$$

where  $\bar{\phi}^{n+1/2}$  is any explicit  $O(\delta t^2)$  approximation for  $\Phi^{n+1/2}$ .

## SAV/BDF3

$$\begin{aligned}
 11\phi^{n+1} - 18\phi^n + 9\phi^{n-1} - 2\phi^{n-2} &= 6\delta t \Delta \mu^{n+1}, \\
 \mu^{n+1} &= -\Delta \phi^{n+1} + \frac{r^{n+1}}{\sqrt{E_1[\bar{\phi}^{n+1}] + C_0}} F'(\bar{\phi}^{n+1}), \\
 11r^{n+1} - 18r^n + 9r^{n-1} - 2r^{n-2} &= \\
 \int_{\Omega} \frac{F'(\bar{\phi}^{n+1})}{2\sqrt{E_1[\bar{\phi}^{n+1}] + C_0}} (11\phi^{n+1} - 18\phi^n + 9\phi^{n-1} - 2\phi^{n-2}) dx.
 \end{aligned} \tag{17}$$

where  $\bar{\phi}^{n+1}$  is any explicit  $O(\delta t^2)$  approximation for  $\Phi^{n+1}$ .

## SAV/BDF3

To obtain  $\bar{\phi}^{n+1}$  in BDF3, we can use the extrapolation (BDF3A):

$$\bar{\phi}^{n+1} = 3\phi^n - 3\phi^{n-1} + \phi^{n-2},$$

or prediction by one BDF2 step (BDF3B):

$$\bar{\phi}^{n+1} = \text{BDF2} \{ \phi^n, \phi^{n-1}, \Delta t \}.$$

It is noticed that using the prediction with a lower order BDF step will double the total computation cost.

# Advantage of SAV

We presented the SAV approach for gradient flows, which is inspired by the Lagrange multiplier/IEQ methods. It preserves many of their advantages, plus:

- ▶ It leads to linear, decoupled equations with **CONSTANT** coefficients. So fast direct solvers are often available!
- ▶ It only requires the nonlinear energy functional, instead of nonlinear energy density, be bounded from below, so it applies to a larger class of gradient flows.
- ▶ For gradient flows with multiple components, the scheme will lead to decoupled equations with constant coefficients to solve at each time step.

# Advantage of SAV

- ▶ A particular advantage of unconditionally energy stable scheme is that it can be coupled with an adaptive time stepping strategy.
- ▶ The proofs are based on variational formulation with simple test functions, so that they can be extended to full discrete discretization with Galerkin approximation in space.
- ▶ We have performed rigorous error analysis to show that, under mild conditions, the solution of proposed schemes converge to the solution of the original problem.[14]



# Disadvantage of SAV

- ▶ Restrictions on the free energy
- ▶ Extrapolation scheme
- ▶ Convergence and error analysis

# Numerical result I

Example: Evolutions of coarsening process and of energy

In this example, we simulate the Cahn-Hilliard equation on  $[0, 2\pi)^2$ , starting from

$$\phi(x, y, 0) = 0.25 + 0.4\mathbf{Rand}(x, y) \quad (18)$$

We choose  $\beta = 4$ ,  $C_0 = 0$  and discretize the space by the Fourier spectral method with  $256 \times 256$  modes.

We investigate the coarsening process with  $\epsilon^2 = 0.01$  and  $\delta t = 2 \times 10^{-5}$ . The reference solution and the results of SAV/BDF2 are shown in Fig.1, no visible difference is observed.

## Numerical result II




Figure: Coarsening process with SAV/BDF2.

# Improvement of SAV

- ▶ BDF4C, BDF4D?
- ▶ ?
- ▶ ?

# Reference I

- [1] C. Elliott and A. Stuart. “The Global Dynamics of Discrete Semilinear Parabolic Equations”. In: *SIAM Journal on Numerical Analysis* 30.6 (1993), pp. 1622–1663. DOI: 10.1137/0730084. eprint: <https://doi.org/10.1137/0730084>. URL: <https://doi.org/10.1137/0730084>.
- [2] David J. Eyre. “Unconditionally gradient stable time marching the Cahn-Hilliard equation”. In: vol. 529. cited By 157. 1998, pp. 39–46. URL: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-0032321134&partnerID=40&md5=c4e46ec1e406a0dac18bf0b43f479fe9>.

## Reference II

- [3] J. Zhu et al. “Coarsening kinetics from a variable-mobility cahn-hilliard equation: Application of a semi-implicit fourier spectral method”. In: *Physical Review E - Statistical Physics, Plasmas, Fluids, and Related Interdisciplinary Topics* 60.4 (1999). cited By 272, pp. 3564–3572. DOI: 10.1103/PhysRevE.60.3564. URL: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-0000707473&doi=10.1103%2fPhysRevE.60.3564&partnerID=40&md5=15be1e4eaf692db7d2124aa4edb22062>.

## Reference III

- [4] J. Shen and X. Yang. “Numerical approximations of allen-cahn and cahn-hilliard equations”. In: *Discrete and Continuous Dynamical Systems* 28.4 (2010). cited By 177, pp. 1669–1691. DOI: 10.3934/dcds.2010.28.1669. URL: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-77957918727&doi=10.3934%2fdcds.2010.28.1669&partnerID=40&md5=e65a742b562bc14cc0a1336341237a8f>.
- [5] F. Guilln-gonzlez and G. Tierra. “On linear schemes for a cahn-hilliard diffuse interface model”. In: *Journal of Computational Physics* 234.1 (2013). cited By 43, pp. 140–171. DOI: 10.1016/j.jcp.2012.09.020. URL: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-84870538130&doi=10.1016%2fj.jcp.2012.09.020&partnerID=40&md5=62937e68204bc7cdfc0dd52da544a484>.

## Reference IV

- [6] S. Badia, F. Guilln-Gonzlez, and J.V. Gutierrez-Santacreu. “Finite element approximation of nematic liquid crystal flows using a saddle-point structure”. In: *Journal of Computational Physics* 230.4 (2011). cited By 19, pp. 1686–1706. DOI: 10.1016/j.jcp.2010.11.033. URL: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-78650568203&doi=10.1016%2fj.jcp.2010.11.033&partnerID=40&md5=f563e70af2e5e364ffcd04ac0ed328f>.



## Reference V

- [7] X. Yang. “Linear, first and second-order, unconditionally energy stable numerical schemes for the phase field model of homopolymer blends”. In: *Journal of Computational Physics* 327 (2016). cited By 35, pp. 294–316. DOI: 10.1016/j.jcp.2016.09.029. URL: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-84988916643&doi=10.1016%2fj.jcp.2016.09.029&partnerID=40&md5=e285d30dc1c5d4246baaf1db0c315fc8>.

## Reference VI

- [8] X. Yang and L. Ju. “Efficient linear schemes with unconditional energy stability for the phase field elastic bending energy model”. In: *Computer Methods in Applied Mechanics and Engineering* 315 (2017). cited By 30, pp. 691–712. DOI: 10.1016/j.cma.2016.10.041. URL: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-85003422191&doi=10.1016%2fj.cma.2016.10.041&partnerID=40&md5=4e00dffa7aefc938a621604de16cce21>.

## Reference VII

- [9] X. Yang, J. Zhao, and Q. Wang. “Numerical approximations for the molecular beam epitaxial growth model based on the invariant energy quadratization method”. In: *Journal of Computational Physics* 333 (2017). cited By 28, pp. 104–127. DOI: 10.1016/j.jcp.2016.12.025. URL: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-85008879638&doi=10.1016%2fj.jcp.2016.12.025&partnerID=40&md5=412203253ec6fad16da1ff0da83872d8>.

## Reference VIII

- [10] H. Yu and X. Yang. “Numerical approximations for a phase-field moving contact line model with variable densities and viscosities”. In: *Journal of Computational Physics* 334 (2017). cited By 17, pp. 665–686. DOI: 10.1016/j.jcp.2017.01.026. URL: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-85010289968&doi=10.1016%2fj.jcp.2017.01.026&partnerID=40&md5=96c02deb5b12f218f0082115aeb92cb7>.

## Reference IX

- [11] J. Zhao et al. “A novel linear second order unconditionally energy stable scheme for a hydrodynamic Q-tensor model of liquid crystals”. In: *Computer Methods in Applied Mechanics and Engineering* 318 (2017). cited By 23, pp. 803–825. DOI: 10.1016/j.cma.2017.01.031. URL: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-85014165227&doi=10.1016%2fj.cma.2017.01.031&partnerID=40&md5=a17679a798ef871261b4d482beb50e90>.

## Reference X

- [12] X. Yang and L. Ju. “Linear and unconditionally energy stable schemes for the binary fluids surfactant phase field model”. In: *Computer Methods in Applied Mechanics and Engineering* 318 (2017). cited By 22, pp. 1005–1029. DOI: 10.1016/j.cma.2017.02.011. URL: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-85014457724&doi=10.1016%2fj.cma.2017.02.011&partnerID=40&md5=0cf2d76e7653ac544ae9235574e468ff>.
- [13] Xiaofeng Yang et al. “Numerical approximations for a three-component Cahn-Hilliard phase-field model based on the invariant energy quadratization method”. In: *Mathematical Models and Methods in Applied Sciences* 27.11 (2017), pp. 1993–2030. DOI: 10.1142/S0218202517500373. eprint: <https://doi.org/10.1142/S0218202517500373>. URL: <https://doi.org/10.1142/S0218202517500373>.

## Reference XI

- [14] J. Shen and J. Xu. “Convergence and Error Analysis for the Scalar Auxiliary Variable (SAV) Schemes to Gradient Flows”. In: *SIAM Journal on Numerical Analysis* 56.5 (2018), pp. 2895–2912. DOI: 10.1137/17M1159968. eprint: <https://doi.org/10.1137/17M1159968>. URL: <https://doi.org/10.1137/17M1159968>.