


Chapter 3.5

Markov State Process

Performance Evaluation of the Internet of Things (IoT)

Module Course: Performance Evaluation of Distributed Systems

Prof. Tobias Hoßfeld, Summer Semester 2022



Disclaimer and Copyright Notice

Lecture slides, figures, and scripts are based on the open access text book "Performance Modeling and Analysis of Communication Networks". The book and scripts are licensed under the Creative Commons License Attribution-ShareAlike 4.0 International ([CC BY-SA 4.0](https://creativecommons.org/licenses/by-sa/4.0/)). If you remix, transform, or build upon the material, you must distribute your contributions under the same license as the original.

The book must be cited and the disclaimer attached when using lectures slides or scripts.

*Tran-Gia, P. & Hossfeld, T. (2021).
Performance Modeling and Analysis of Communication
Networks - A Lecture Note. Würzburg University Press.
<https://doi.org/10.25972/WUP-978-3-95826-153-2>*

Website to download book, exercises, slides and scripts:
<https://modeling.systems/>

Chapter 3

3 Elementary Random Processes

3.1 Stochastic Processes

3.1.1 Definition

3.1.2 Markov Processes

3.1.3 Elementary Processes in Performance Models

3.2 Renewal Processes

3.2.1 Definition

3.2.2 Analysis of Recurrence Time

3.3 Poisson Process

3.3.1 Definition of a Poisson Process

3.3.2 Properties of the Poisson Process

3.3.3 Poisson Arrivals during Arbitrarily Distributed Interval

3.4 Superposition of Independent Renewal Processes

3.4.1 Superposition of Poisson Processes

3.4.2 Palm-Khintchine Theorem

3.5 Markov State Process

3.5.1 Definition of Continuous-Time Markov Chain

3.5.2 Transition Behavior of Markovian State Processes

3.5.3 State Equations and State Probabilities

3.5.4 Examples of Transition Probability Densities

3.5.5 Birth-and-Death Processes

DEFINITION OF CONTINUOUS-TIME MARKOV CHAIN

Continuous-Time Markov Chain (CTMC)

▶ Continuous-time Markov process

- Stochastic process $\{X(t), t \geq 0\}$ with Markov property

▶ Continuous-time Markov chain (CMTC) is defined by

- discrete state space S is finite or countable; e.g. number of customers in system
- transition rates $q_{ij} \geq 0$ for $i \neq j$ and $i, j \in S$
- initial state $X(0)$, i.e. probability distribution of initial state

▶ Probability $x(i, t) = P(X(t) = i)$ that the system is in state $[X = i]$ at time t


▶ State vector $\mathbf{X}(t) = (x(0, t), x(1, t), \dots)$

▶ Definition of **rate matrix** \mathbf{Q} with $q_{ii} = -\sum_{i \neq j} q_{ij}$

- allows compact notation (Kolmogorov equations.)
- row-wise sums of \mathbf{Q} are 0

Illustration of CTMC: Example

- ▶ State space $S = \{0, 1, 2, \dots, n\}$



- ▶ Transition rates


- $q_{i,i+1} = \lambda_i$ for $i = 0, 1, \dots, n-1$
- $q_{i,i-1} = \mu_i$ for $i = 1, \dots, n$
- otherwise: $q_{i,j} = 0$ for $i \neq j$
- $q_{ii} = -\sum_{i \neq j} q_{i,j}$

- ▶ Transition matrix

$$Q = \begin{pmatrix} q_{00} & q_{01} & \dots & q_{0j} & \dots \\ q_{10} & q_{11} & \dots & q_{1j} & \dots \\ \vdots & \vdots & \ddots & \vdots & \\ q_{j0} & q_{j1} & \dots & q_{jj} & \dots \\ \vdots & \vdots & & \vdots & \ddots \end{pmatrix}$$


Transition Behavior of CTMC

- ▶ System remains in state i for time T_i
 - exponentially distributed with rate $q_i = -q_{ii} > 0$
 - time to change from i to j : $T_{ij} \sim \text{EXP}(q_{ij})$
 - $T_i = \min_{i \neq j} \{T_{ij}\} \sim \text{EXP}(q_i)$




- ▶ When process leaves state i , the state j is reached with probability

- $p_{ij} = \frac{q_{ij}}{q_i}$




Analysis of Continuous-time Markov Chain (CMTC)

independent outside observer at time t_n sees system state probability x_n



state $[X(t_n) = x_n]$



future development of process only depends on state x_n


Overview on the Analysis

- ▶ Markov state process and memoryless property
- ▶ **Homogeneous systems**: transition probabilities independent of time instant
- ▶ **Chapman-Kolmogorov equation** for state transition probabilities: $\mathcal{P}(t+\Delta t) = \mathcal{P}(t) \cdot \mathcal{P}(\Delta t)$
 - Kolmogorov forward equation for **probability densities** $\lim_{\Delta t \rightarrow 0} \mathcal{P}(t+\Delta t) \longrightarrow \frac{d\mathcal{P}(t)}{dx} = \mathcal{P}(t) \cdot Q$
 - Kolmogorov forward equation for **state probabilities** $\mathcal{X}(t) = \mathcal{X}(0) \cdot \mathcal{P}(t) \longrightarrow \frac{d\mathcal{X}(t)}{dx} = \mathcal{X}(t) \cdot Q$
- ▶ **Stationary system** $\lim_{t \rightarrow \infty} \longrightarrow \frac{\partial}{\partial t} x(j, t) = 0 \quad \mathcal{X} \cdot Q = 0$
- ▶ Example: **birth-and-death process**

TRANSITION BEHAVIOR OF MARKOVIAN STATE PROCESSES

Chapman-Kolmogorov equation

Transition Behavior of Markovian State Processes



Transition Probability

- ▶ State transition $i \rightarrow j$ during interval $\Delta t = t_{n+1} - t_n$ occurs with probability

$$p_{ij}(t_n, t_{n+1}) = P(X(t_{n+1})=j \mid X(t_n)=i)$$


- ▶ **Time-homogeneous state process**

- transition behavior is identical for each process point in time
- transition probability is independent of the observation instant

$$p_{ij}(t_n, t_{n+1}) = p_{ij}(t_{n+1} - t_n) = p_{ij}(\Delta t)$$

- standardization condition for all i


$$\sum_j p_{ij}(\Delta t) = 1, \quad \Delta t \geq 0$$



Transition Matrix

- ▶ Transition probabilities $\{p_{ij}(\Delta t), i, j = 0, 1, \dots\}$ form the transition matrix

$$\mathcal{P}(\Delta t) = \begin{pmatrix} p_{00}(\Delta t) & p_{01}(\Delta t) & \dots & p_{0j}(\Delta t) & \dots \\ p_{10}(\Delta t) & p_{11}(\Delta t) & \dots & p_{1j}(\Delta t) & \dots \\ \vdots & \vdots & & \vdots & \\ p_{i0}(\Delta t) & p_{i1}(\Delta t) & \dots & p_{ij}(\Delta t) & \dots \\ \vdots & \vdots & & \vdots & \end{pmatrix}$$




Chapman-Kolmogorov Equation

$$\mathcal{P}(t + \Delta t) = \mathcal{P}(t) \cdot \mathcal{P}(\Delta t)$$

or

$$p_{ij}(t + \Delta t) = \sum_k p_{ik}(t) p_{kj}(\Delta t)$$



Chapman-Kolmogorov Equation (f.)

Lecture

KOLMOGOROV FORWARD EQUATION FOR TRANSITION PROBABILITIES

Kolmogorov Forward Equation for Transition Probabilities

- ▶ Chapman-Kolmogorov equation

$$p_{ij}(t + \Delta t) = \sum_k p_{ik}(t) p_{kj}(\Delta t)$$

- ▶ can be formulated as

$$p_{ij}(t + \Delta t) = \sum_{k \neq j} p_{ik}(t) p_{kj}(\Delta t) + p_{ij}(t) p_{jj}(\Delta t)$$

$$\frac{p_{ij}(t + \Delta t) - p_{ij}(t)}{\Delta t} = \sum_{k \neq j} p_{ik}(t) \cdot \frac{p_{kj}(\Delta t)}{\Delta t} - p_{ij}(t) \cdot \frac{1 - p_{jj}(\Delta t)}{\Delta t}$$

- ▶ Next: limiting process $\Delta t \rightarrow 0$

Kolmogorov Forward Equation: Limiting Process

$$\frac{p_{ij}(t + \Delta t) - p_{ij}(t)}{\Delta t} = \sum_{k \neq j} p_{ik}(t) \cdot \frac{p_{kj}(\Delta t)}{\Delta t} - p_{ij}(t) \cdot \frac{1 - p_{jj}(\Delta t)}{\Delta t}$$

► Limiting process $\Delta t \rightarrow 0$

$$\lim_{\Delta t \rightarrow 0} \frac{p_{ij}(t + \Delta t) - p_{ij}(t)}{\Delta t} = \frac{d}{dt} p_{ij}(t)$$

first derivative of transition probabilities $p_{ij}(t)$ at time t

$$\lim_{\Delta t \rightarrow 0} \frac{p_{kj}(\Delta t)}{\Delta t} = q_{kj}, \quad k \neq j$$

transition probability density for the transition $k \rightarrow j$

$$\lim_{\Delta t \rightarrow 0} \frac{1 - p_{jj}(\Delta t)}{\Delta t} = q_j = \sum_{k \neq j} q_{jk}$$

transition probability density for leaving the state j

► **Kolmogorov forward equation for transition probabilities**

$$\frac{d}{dt} p_{ij}(t) = \sum_{k \neq j} q_{kj} p_{ik}(t) - q_j p_{ij}(t)$$

Kolmogorov Forward Equation: Matrix Notation

- ▶ Kolmogorov forward equation for transition probabilities

$$\frac{d}{dt} p_{ij}(t) = \sum_{k \neq j} q_{kj} p_{ik}(t) - q_j p_{ij}(t)$$

- ▶ Matrix for transition probability densities is defined

$$Q = \begin{pmatrix} q_{00} & q_{01} & \cdots & q_{0j} & \cdots \\ q_{10} & q_{11} & \cdots & q_{1j} & \cdots \\ \vdots & \vdots & \ddots & \vdots & \\ q_{j0} & q_{j1} & \cdots & q_{jj} & \cdots \\ \vdots & \vdots & & \vdots & \end{pmatrix} \quad \begin{array}{l} \text{with } \sum_k q_{jk} = 0 \quad \text{row-wise sum is 0} \\ \text{and } q_{jj} = -\sum_{k \neq j} q_{jk} = -q_j \quad \text{probability density to remain in state } j \end{array}$$



$$\frac{d\mathcal{P}(t)}{dt} = \mathcal{P}(t) \cdot Q$$

- ▶ Note:

- „ $-q_j$ “ does not imply that there is a negative rate.
- solely notation for the rate to stay in state

Solution of Kolmogorov Forward Equation

- ▶ Kolmogorov forward equation for transition probabilities in matrix notation

$$\frac{d\mathcal{P}(t)}{dt} = \mathcal{P}(t) \cdot Q$$

- ▶ Solution requires the computation of the matrix exponential of the matrix $t \cdot Q$

$$\mathcal{P}(t) = e^{tQ} = \sum_{k=0}^{\infty} \frac{(tQ)^k}{k!}$$

- ▶ See notebook script "3.5 [Markov processes](https://modeling.systems/): nonstationary and stationary analysis [\[ipynb\]](#)"
<https://modeling.systems/>

STATE EQUATIONS AND STATE PROBABILITIES


Kolmogorov Forward Equation for State Probabilities

Kolmogorov Forward Equation for State Probabilities

- Probability $x(j, t)$ for the system to be in state j at time t .

$x(j, t) = P(X(t) = j)$ process is in state j at time t


$x(i, 0)$ initial state at time $t = 0$



$$x(j, t) = \sum_i P(X(t) = j | X(0) = i) \cdot P(X(0) = i) = \sum_i x(i, 0) \cdot p_{ij}(t)$$

- Kolmogorov forward equation **for transition probabilities**

$$\frac{d}{dt} p_{ij}(t) = \sum_{k \neq j} q_{kj} p_{ik}(t) - q_j p_{ij}(t)$$



$$\sum_i \frac{d}{dt} p_{ij}(t) x(i, 0) = \sum_{k \neq j} q_{kj} \sum_i (p_{ik}(t) x(i, 0)) - \sum_i (q_j p_{ij}(t) x(i, 0))$$

- Kolmogorov forward equation **for state probabilities**

$$\frac{\partial}{\partial t} x(j, t) = \sum_{k \neq j} q_{kj} x(k, t) - q_j x(j, t), \quad j = 0, 1, \dots, \quad \left(\sum_j x(j, t) = 1 \right)$$

Kolmogorov Forward Eq. for State Probabilities: Matrix Notation

- ▶ Kolmogorov forward equation **for state probabilities** is system of differential equations

$$\frac{\partial}{\partial t} x(j, t) = \sum_{k \neq j} q_{kj} x(k, t) - q_j x(j, t), \quad j = 0, 1, \dots,$$

- ▶ State probabilities as vector $\mathbf{X}(t) = (x(0, t), x(1, t), \dots, x(j, t), \dots)$

- ▶ Initial state and transition probability matrix $\mathbf{X}(t) = \mathbf{X}(0) \cdot \mathcal{P}(t)$

- ▶ **Matrix notation** of Kolmogorov forward equation for state probabilities

$$\frac{d}{dt} \mathbf{X}(t) = \mathbf{X}(t) \cdot \mathcal{Q}$$

Stationary State Probabilities

- ▶ State process is **stationary** if
 - system state stops changing (statistical equilibrium)
 - state probability no longer depends on time t

$$\frac{d}{dt} P(X(t) = j) = \frac{\partial}{\partial t} x(j, t) = 0$$

- ▶ Stationary state probability

$$x(j) = \lim_{t \rightarrow \infty} P(X(t) = j)$$


- ▶ Results in stationary equation system

$$\frac{\partial}{\partial t} x(j, t) = \sum_{k \neq j} q_{kj} x(k, t) - q_j x(j, t), \quad j = 0, 1, \dots,$$



$$q_j x(j) = \sum_{k \neq j} q_{kj} \cdot x(k), \quad j = 0, 1, \dots,$$

$$\sum_j x(j) = 1.$$



Principle of Maintaining Statistical Equilibrium

$$\sum_j x(j) = 1$$

$$q_j x(j) = \sum_{k \neq j} q_{kj} x(k)$$




rate for
leaving
state j



rate for
reaching
state j

► Principle of maintaining statistical equilibrium

- Stationary system: flows of weighted probability densities for reaching and leaving a state must be in equilibrium, i.e., they are the same
- state probability no longer changes in time




Stationary Equation System: Matrix Notation

$$\sum_j x(j) = 1$$

$$q_j x(j) = \sum_{k \neq j} q_{kj} x(k)$$

↑
↑

rate for leaving state j = rate for reaching state j



► Stationary state probability vector

$$\mathcal{X} = \{x(0), x(1), \dots, x(j), \dots\}$$

► **State equation system**

$$\mathcal{X} \cdot Q = 0$$

$$\left(\sum_j x(j) = 1 \quad \text{or} \quad \mathcal{X} e = 1 \right)$$

Linear Dependency of State Equations

- ▶ Consider a finite state space $\{0,1,2, \dots, N\}$
- ▶ Stationary system of equations

$$\sum_{j=1}^N x(j) = 1$$
$$q_j x(j) = \sum_{k \neq j} q_{kj} x(k), \quad j=0,1,\dots,N$$

- $(N + 2)$ equations for $(N + 1)$ unknowns (state probabilities)
- Any arbitrarily chosen equation can be omitted due to linear dependency to solve the equation system

Macro States


- ▶ Single state that cannot be further decomposed is also called a **micro state**
- ▶ Combination of any set of micro states leads to a **macro state**

- ▶ Example

$$+ \begin{cases} (q_{i1} + q_{ij} + q_{ik})x(i) = q_{1i}x(1) + q_{ji}x(j) + q_{ki}x(k) \\ (q_{j2} + q_{jk} + q_{ji})x(j) = q_{2j}x(2) + q_{kj}x(k) + q_{ij}x(i) \\ (q_{k3} + q_{ki} + q_{kj})x(k) = q_{3k}x(3) + q_{ik}x(i) + q_{jk}x(j) \end{cases}$$

$$q_{i1}x(i) + q_{j2}x(j) + q_{k3}x(k) = q_{1i}x(1) + q_{2j}x(2) + q_{3k}x(3)$$

- ▶ Appropriate choice of macro states often provides simpler system of equations for computing the (micro) state probabilities



Global Equilibrium Equation

- State equation for an arbitrary macro state

$$\underbrace{\sum_{j \in S, u \notin S} q_{ju} x(j)}_{\text{weighted rates for leaving the macro state } S} = \underbrace{\sum_{u \notin S, j \in S} q_{uj} x(u)}_{\text{weighted rates for reaching the macro state } S}$$

- Relates transition probability densities between a macro state and rest in the state space
- Global equilibrium equations are also referred to as full or **global balance equations**

Example: Stationary State Equations

Example: Stationary Macro State Equations

Lecture


Summary: Stationary Equation System

Lecture

EXAMPLES OF TRANSITION PROBABILITY DENSITIES


Transition Probability Densities

- What is the transition probability density of a Markov state process?




- Example: M/M/n-0

- transition rates of a Poisson process? $q_{i,i+1} = \lambda$.



Transition Probability Density of Poisson Arrival Process



- Transition $i \rightarrow i + 1$: arriving customer is accepted

$$\begin{aligned}
 q_{i,i+1} &= \lim_{dt \rightarrow 0} \frac{p_{i,i+1}(dt)}{dt} \\
 &= \lim_{dt \rightarrow 0} \frac{P(R \leq dt)}{dt} = \lim_{dt \rightarrow 0} \frac{1 - e^{-\lambda dt}}{dt} \stackrel{\text{L'Hospital}}{=} \lambda
 \end{aligned}$$

$$p_{i,i+1}(dt) = P(X(t + dt) = i + 1 | X(t) = i)$$

Transition Probability Density for Exponential Service Time

- ▶ k servers with exponential service times

$$R(t) = B(t) = 1 - e^{-\mu t}$$


- ▶ Interval R^* until next service termination

$$R^* = \min\{\underbrace{R, \dots, R}_{k\text{-times}}\}$$

$$R^*(t) = 1 - \prod_{i=1}^k (1 - R(t)) = 1 - e^{-k\mu t}$$

- ▶ Transition $k \rightarrow k - 1$


$$\begin{aligned} q_{k,k-1} &= \lim_{dt \rightarrow 0} \frac{p_{k,k-1}(dt)}{dt} \\ &= \lim_{dt \rightarrow 0} \frac{P(R^* \leq dt)}{dt} = \lim_{dt \rightarrow 0} \frac{1 - e^{-k\mu dt}}{dt} \stackrel{\text{L'Hospital}}{=} k\mu \end{aligned}$$



BIRTH-AND-DEATH PROCESSES

Birth-and-Death Processes

- ▶ Markov processes in which only transitions between neighboring states occur
- ▶ State transition diagram of a finite-state, one-dimensional BD process



$$q_{ij} = \begin{cases} \lambda_i & i = 0, 1, \dots, n-1, \quad j = i+1 & \text{birth rate} \\ \mu_i & i = 1, 2, \dots, n, \quad j = i-1 & \text{death rate} \\ 0 & \text{otherwise} \end{cases}$$

- ▶ Pure birth process: all $\mu_i = 0$
 - equilibrium: $x(n) = 1, x(i) = 0$ otherwise
- ▶ Pure death process: all $\lambda_i = 0$
 - equilibrium: $x(0) = 1, x(i) = 0$ otherwise


Example: No BD Process

Non-stationary Birth-and-Death Processes

- ▶ Transient (time-dependent) state probabilities of the BD process

$$x(i, t) = P(X(t) = i)$$

$$\Rightarrow \left\{ \begin{array}{l} \frac{\partial}{\partial t} x(0, t) = -\lambda_0 x(0, t) + \mu_1 x(1, t) \\ \frac{\partial}{\partial t} x(i, t) = -(\lambda_i + \mu_i) x(i, t) + \lambda_{i-1} x(i-1, t) + \mu_{i+1} x(i+1, t), \quad i = 1, \dots, n-1 \\ \frac{\partial}{\partial t} x(n, t) = -\mu_n x(n, t) + \lambda_{n-1} x(n-1, t) \end{array} \right.$$



- ▶ Solution of this differential equation system with the initial conditions $\{x(i, 0), i = 0, \dots, n\}$


$$\mathbf{X}(t) = (x(0, t), x(1, t), \dots, x(n, t))$$

- ▶ Example: Poisson Process as pure birth process

Example: Poisson Process as Pure Birth Process

Stationary Birth-and-Death Processes

- Equation system for the micro states of the state space




$$\left. \begin{array}{ll} 0 : & \lambda_0 x(0) = \mu_1 x(1) \\ i \in (1, n-1) : & (\lambda_i + \mu_i) x(i) = \lambda_{i-1} x(i-1) + \mu_{i+1} x(i+1), \\ n : & \lambda_{n-1} x(n-1) = \mu_n x(n) \end{array} \right\}$$

because of linear dependency, one of these equations can be omitted

$$\sum_{i=0}^n x(i) = 1$$

Macro State Equation System



- State equation for macro state S

$$i \in (1, n) \quad \lambda_{i-1} x(i-1) = \mu_i x(i),$$


$$\left(\sum_{i=0}^n x(i) = 1 \right)$$



$$x(i) = x(0) \cdot \frac{\prod_{k=0}^{i-1} \lambda_k}{\prod_{k=1}^i \mu_k}, \quad i = 1, 2, \dots, n$$

- Normalization yields $x(0)$

$$1 = \sum_{i=0}^n x(i) = x(0) + x(0) \sum_{i=1}^n \frac{\prod_{k=0}^{i-1} \lambda_k}{\prod_{k=1}^i \mu_k}$$




$$x(0) = \left(1 + \sum_{i=1}^n \frac{\prod_{k=0}^{i-1} \lambda_k}{\prod_{k=1}^i \mu_k} \right)^{-1}$$

Solving the Macro State Equation System

Lecture

Example: Delay-Loss System with State-dependent Rates

- ▶ Delay-loss system $M(x)/M/2-1$ as an example for birth-and-death process



- ▶ Customers arrive with state-dependent arrival rates
 - not a Poisson process
 - PASTA property is not valid: $x_A(i) \neq x^*(i)$
- ▶ State equation system yields arbitrary-time state probabilities $x^*(i)$
 - e.g. blocking probability requires $x_A(i)$ for arriving customers
 - strong law of large numbers for Markov chains can be applied

Strong Law of Large Numbers for Markov Chains

- ▶ Expected number of arrivals when the system is in state $[X = i]$ in interval of length T is $n_A(i, T)$
- ▶ Expected total number $n_A(T)$ of arrivals

- ▶ **Strong law of large numbers for Markov chains**

$$x_A(i) = \lim_{T \rightarrow \infty} \frac{n_A(i, T)}{n_A(T)} = \frac{\lambda_i \cdot x(i)}{\sum_k \lambda_k \cdot x(k)}$$

- ▶ For birth-and-death process, as in the example, $x_A(i)$ and characteristics like blocking probability or waiting probability can be derived accordingly

