

# FREE RESOLUTIONS OVER A COMPLETE HYPERSURFACE (AND FRIENDS)

BASED ON “HOMOLOGICAL ALGEBRA ON A COMPLETE  
INTERSECTION, WITH AN APPLICATION TO GROUP  
REPRESENTATIONS.” BY DAVID EISENBUD

David DeMark

MATH 8212 University of Minnesota

2 May 2018

**Throughout this talk, we let...**

**Throughout this talk, we let...**

- $(A, \mathfrak{m})$  be a regular local ring

**Throughout this talk, we let...**

- $(A, \mathfrak{m})$  be a regular local ring
- $x_1, \dots, x_n$  a regular  $A$ -sequence of finite length

**Throughout this talk, we let...**

- $(A, \mathfrak{m})$  be a regular local ring
- $x_1, \dots, x_n$  a regular  $A$ -sequence of finite length
- $B := A/\langle x_1, \dots, x_n \rangle$

**Throughout this talk, we let...**

- $(A, \mathfrak{m})$  be a regular local ring
- $x_1, \dots, x_n$  a regular  $A$ -sequence of finite length
- $B := A/\langle x_1, \dots, x_n \rangle$

Such a  $B$  is a *complete intersection of codimension  $n$* .

**Throughout this talk, we let...**

- $(A, \mathfrak{m})$  be a regular local ring
- $x_1, \dots, x_n$  a regular  $A$ -sequence of finite length
- $B := A/\langle x_1, \dots, x_n \rangle$

Such a  $B$  is a *complete intersection of codimension  $n$* .

We shall study the structure of  $B$ -free resolutions of  $B$ -modules, relating these to their liftings to  $A$ .

# CLARIFICATION: $B$ -FREE

We do **NOT** mean. . .



# CLARIFICATION: $B$ -FREE

We do **NOT** mean...




FIGURE: Another “ $B$ -Free” Object.

# WHY DO WE CARE?

# WHY DO WE CARE?

- Classifying maximal CM modules over complete hypersurfaces! (case  $n = 1$ )

# WHY DO WE CARE?

- Classifying maximal CM modules over complete hypersurfaces! (case  $n = 1$ )
- Expanding upon a familiar characterization!

# WHY DO WE CARE?

- Classifying maximal CM modules over complete hypersurfaces! (case  $n = 1$ )
- Expanding upon a familiar characterization!

## THEOREM (AUSLANDER-BUCHSBAUM-SERRE)

*For  $(R, \mathfrak{m})$  local,  $R$  is regular  $\iff \operatorname{gl dim}(R) < \infty$ .*

# WHY DO WE CARE?

- Classifying maximal CM modules over complete hypersurfaces! (case  $n = 1$ )
- Expanding upon a familiar characterization!

## THEOREM (AUSLANDER-BUCHSBAUM-SERRE)

*For  $(R, \mathfrak{m})$  local,  $R$  is regular  $\iff \text{gl dim}(R) < \infty$ .*

We will give a similar (albeit more complicated) characterization of codim-1 quotients of regular local rings.

## CODIM $n$ : SET-UP

Recall that  $B := A/\langle x_1, \dots, x_n \rangle$ . Let  $M$  be a  $B$ -module, and  $\mathbf{F}$  a free resolution of  $B$ .

$$\mathbf{F} : \dots \xrightarrow{\partial_3} F_2 \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} F_0 \xrightarrow{\partial_0} 0$$

# CODIM $n$ : SET-UP

Recall that  $B := A/\langle x_1, \dots, x_n \rangle$ . Let  $M$  be a  $B$ -module, and  $\mathbf{F}$  a free resolution of  $B$ .

$$\mathbf{F} : \dots \xrightarrow{\partial_3} F_2 \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} F_0 \xrightarrow{\partial_0} 0$$

**Now let...**



## CODIM $n$ : SET-UP

Recall that  $B := A/\langle x_1, \dots, x_n \rangle$ . Let  $M$  be a  $B$ -module, and  $\mathbf{F}$  a free resolution of  $B$ .

$$\mathbf{F} : \dots \xrightarrow{\partial_3} F_2 \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} F_0 \xrightarrow{\partial_0} 0$$

Now let...

- $\tilde{F}_i$  be a free  $A$ -module with  $\text{rank}_A \tilde{F}_i = \text{rank}_B F_i$

## CODIM $n$ : SET-UP

Recall that  $B := A/\langle x_1, \dots, x_n \rangle$ . Let  $M$  be a  $B$ -module, and  $\mathbf{F}$  a free resolution of  $B$ .

$$\mathbf{F} : \dots \xrightarrow{\partial_3} F_2 \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} F_0 \xrightarrow{\partial_0} 0$$

Now let...

- $\tilde{F}_i$  be a free  $A$ -module with  $\text{rank}_A \tilde{F}_i = \text{rank}_B F_i$
- $\tilde{\partial}_i$  denote an arbitrary lifting of (the entries of)  $\partial_i$  to  $A$

**Consider**

$$\tilde{\mathbf{F}} : \dots \xrightarrow{\tilde{\partial}_3} \tilde{F}_2 \xrightarrow{\tilde{\partial}_2} \tilde{F}_1 \xrightarrow{\tilde{\partial}_1} \tilde{F}_0 \xrightarrow{\tilde{\partial}_0} 0$$

Consider

$$\tilde{\mathbf{F}} : \dots \xrightarrow{\tilde{\partial}_3} \tilde{F}_2 \xrightarrow{\tilde{\partial}_2} \tilde{F}_1 \xrightarrow{\tilde{\partial}_1} \tilde{F}_0 \xrightarrow{\tilde{\partial}_0} 0$$

- $\tilde{\mathbf{F}}$  is *not* necessarily a complex, but  
 $\text{Im}(\tilde{\partial}_i \circ \tilde{\partial}_{i+1}) \subset \langle x_1, \dots, x_n \rangle F_{i-1}$

Consider

$$\tilde{\mathbf{F}} : \dots \xrightarrow{\tilde{\partial}_3} \tilde{F}_2 \xrightarrow{\tilde{\partial}_2} \tilde{F}_1 \xrightarrow{\tilde{\partial}_1} \tilde{F}_0 \xrightarrow{\tilde{\partial}_0} 0$$

- $\tilde{\mathbf{F}}$  is *not* necessarily a complex, but  
 $\text{Im}(\tilde{\partial}_i \circ \tilde{\partial}_{i+1}) \subset \langle x_1, \dots, x_n \rangle F_{i-1}$
- Write  $\tilde{\partial}_i \otimes \tilde{\partial}_{i+1} = \sum_{j=1}^n x_j \tilde{t}_j$  for some transformation  
 $\tilde{t}_j := \tilde{t}_j(A, \{x_i\}, \mathbf{F}) : \tilde{\mathbf{F}} \rightarrow \tilde{\mathbf{F}}.$

## Consider

$$\tilde{\mathbf{F}} : \dots \xrightarrow{\tilde{\partial}_3} \tilde{F}_2 \xrightarrow{\tilde{\partial}_2} \tilde{F}_1 \xrightarrow{\tilde{\partial}_1} \tilde{F}_0 \xrightarrow{\tilde{\partial}_0} 0$$

- $\tilde{\mathbf{F}}$  is *not* necessarily a complex, but  $\text{Im}(\tilde{\partial}_i \circ \tilde{\partial}_{i+1}) \subset \langle x_1, \dots, x_n \rangle F_{i-1}$
- Write  $\tilde{\partial}_i \otimes \tilde{\partial}_{i+1} = \sum_{j=1}^n x_j \tilde{t}_j$  for some transformation  $\tilde{t}_j := \tilde{t}_j(A, \{x_i\}, \mathbf{F}) : \tilde{\mathbf{F}} \rightarrow \tilde{\mathbf{F}}$ .
- Finally, let  $t_j = t_j(A, \{x_i\}, \mathbf{F}) : \mathbf{F} \rightarrow \mathbf{F}$  be defined by  $t_j = B \otimes \tilde{t}_j$ .

# THE $t_j$ HAVE MANY NICE PROPERTIES!

## THEOREM (SUMMARY OF §1)

*The  $t_j$  have many nice properties! These include:*

# THE $t_j$ HAVE MANY NICE PROPERTIES!

## THEOREM (SUMMARY OF §1)

*The  $t_j$  have many nice properties! These include:*

- *The  $t_j$  are unique (w/r/t fixed  $A, \mathbf{F}, \{x_i\}$ ) up to homotopy.*



# THE $t_j$ HAVE MANY NICE PROPERTIES!

## THEOREM (SUMMARY OF §1)

*The  $t_j$  have many nice properties! These include:*

- *The  $t_j$  are unique (w/r/t fixed  $A, \mathbf{F}, \{x_i\}$ ) up to homotopy.*
- *The  $t_j$  are natural in the sense that they commute with morphisms of complexes up to homotopy.*

# THE $t_j$ HAVE MANY NICE PROPERTIES!

## THEOREM (SUMMARY OF §1)

*The  $t_j$  have many nice properties! These include:*

- *The  $t_j$  are unique (w/r/t fixed  $A, \mathbf{F}, \{x_i\}$ ) up to homotopy.*
- *The  $t_j$  are natural in the sense that they commute with morphisms of complexes up to homotopy.*
- *The  $t_j$  commute with each other up to homotopy.*

# THE $t_j$ HAVE MANY NICE PROPERTIES!

## THEOREM (SUMMARY OF §1)

*The  $t_j$  have many nice properties! These include:*

- *The  $t_j$  are unique (w/r/t fixed  $A, \mathbf{F}, \{x_i\}$ ) up to homotopy.*
- *The  $t_j$  are natural in the sense that they commute with morphisms of complexes up to homotopy.*
- *The  $t_j$  commute with each other up to homotopy.*
- *If  $M, N$  are  $B$ -modules, with  $B$ -free resolutions  $\mathbf{F} \rightarrow M$ ,  $\mathbf{G} \rightarrow N$   $t_j(\mathbf{F})$  and  $t_j(\mathbf{G})$  induce the same map on  $\mathrm{Tor}_{\bullet}^B(M, N)$ .*

# THE $t_j$ HAVE MANY NICE PROPERTIES!

## THEOREM (SUMMARY OF §1)

*The  $t_j$  have many nice properties! These include:*

- *The  $t_j$  are unique (w/r/t fixed  $A, \mathbf{F}, \{x_i\}$ ) up to homotopy.*
- *The  $t_j$  are natural in the sense that they commute with morphisms of complexes up to homotopy.*
- *The  $t_j$  commute with each other up to homotopy.*
- *If  $M, N$  are  $B$ -modules, with  $B$ -free resolutions  $\mathbf{F} \rightarrow M$ ,  $\mathbf{G} \rightarrow N$   $t_j(\mathbf{F})$  and  $t_j(\mathbf{G})$  induce the same map on  $\mathrm{Tor}_\bullet^B(M, N)$ .*
- *The  $t_j$  play only a little less well with ring morphisms  $\alpha : A \rightarrow A'$ .*

# INTERLUDE: MATRIX FACTORIZATIONS

Let  $x \in A$  and  $\phi : F \rightarrow G$  a morphism of  $A$ -modules.

# INTERLUDE: MATRIX FACTORIZATIONS

Let  $x \in A$  and  $\phi : F \rightarrow G$  a morphism of  $A$ -modules.

## DEFINITION

The ordered pair  $(\psi, \phi)$  is a *matrix factorization of  $x$*  if  $\psi \circ \phi = 1_F$  and  $\phi \circ \psi = x \cdot 1_G$ .

# INTERLUDE: MATRIX FACTORIZATIONS

Let  $x \in A$  and  $\phi : F \rightarrow G$  a morphism of  $A$ -modules.

## DEFINITION

The ordered pair  $(\psi, \phi)$  is a *matrix factorization* of  $x$  if  $\psi \circ \phi = 1_F$  and  $\phi \circ \psi = x \cdot 1_G$ .

## THEOREM (5.5)

Given  $\phi$ , there exists a  $\psi$  such that  $(\psi, \phi)$  is a matrix factorization if and only if **each** of the following hold:

- $\text{rank } F = \text{rank } G$
- $\det \phi$  is a non-zero divisor
- $x \cdot \text{Fitt}_1(\phi) \subset \langle \det \phi \rangle$ , i.e.  $x$  annihilates  $\text{coker} \phi$  i.e.  $\phi$  is  $f$

# A QUICK COMPUTATION

- $A := k[W, Y, Z]$  with  $k$  a field of characteristic 0. Set  $x = W(Y^2 - Z^2)$ .



# A QUICK COMPUTATION

- $A := k[W, Y, Z]$  with  $k$  a field of characteristic 0. Set  $x = W(Y^2 - Z^2)$ .

$$\phi = \begin{bmatrix} Y - Z & 0 & Y - Z \\ Y - Z & 0 & 0 \\ 0 & W & 0 \end{bmatrix}$$

# A QUICK COMPUTATION

- $A := k[W, Y, Z]$  with  $k$  a field of characteristic 0. Set  $x = W(Y^2 - Z^2)$ .

$$\phi = \begin{bmatrix} Y - Z & 0 & Y - Z \\ Y - Z & 0 & 0 \\ 0 & W & 0 \end{bmatrix} \quad \det \phi = W(Y - Z)^2$$

# A QUICK COMPUTATION

- $A := k[W, Y, Z]$  with  $k$  a field of characteristic 0. Set  $x = W(Y^2 - Z^2)$ .

$$\phi = \begin{bmatrix} Y - Z & 0 & Y - Z \\ Y - Z & 0 & 0 \\ 0 & W & 0 \end{bmatrix} \quad \det \phi = W(Y - Z)^2$$

$$\phi^c = \begin{bmatrix} 0 & W(Y - Z) & 0 \\ 0 & 0 & (Y - Z)^2 \\ W(Y - Z) & W(Z - Y) & 0 \end{bmatrix}$$

# A QUICK COMPUTATION

- $A := k[W, Y, Z]$  with  $k$  a field of characteristic 0. Set  $x = W(Y^2 - Z^2)$ .

$$\phi = \begin{bmatrix} Y - Z & 0 & Y - Z \\ Y - Z & 0 & 0 \\ 0 & W & 0 \end{bmatrix} \quad \det \phi = W(Y - Z)^2$$

$$\phi^c = \begin{bmatrix} 0 & W(Y - Z) & 0 \\ 0 & 0 & (Y - Z)^2 \\ W(Y - Z) & W(Z - Y) & 0 \end{bmatrix}$$

- Thus,  $\text{Fitt}_1(\phi) = \langle W(Y - Z), (Y - Z)^2 \rangle$  and indeed  $x \cdot \text{Fitt}_1 \subset \langle \det \phi \rangle$

# A QUICK COMPUTATION

- $A := k[W, Y, Z]$  with  $k$  a field of characteristic 0. Set  $x = W(Y^2 - Z^2)$ .

$$\phi = \begin{bmatrix} Y - Z & 0 & Y - Z \\ Y - Z & 0 & 0 \\ 0 & W & 0 \end{bmatrix} \quad \det \phi = W(Y - Z)^2$$

$$\phi^c = \begin{bmatrix} 0 & W(Y - Z) & 0 \\ 0 & 0 & (Y - Z)^2 \\ W(Y - Z) & W(Z - Y) & 0 \end{bmatrix}$$

- Thus,  $\text{Fitt}_1(\phi) = \langle W(Y - Z), (Y - Z)^2 \rangle$  and indeed  $x \cdot \text{Fitt}_1 \subset \langle \det \phi \rangle$
- $\psi$  is uniquely determined by  $\phi$ :  $\psi = \frac{x}{\det \phi} \phi^c = \frac{(Y+Z)}{(Y-Z)} \phi^c$

# FROM MATRIX FACTORIZATIONS TO FREE RESOLUTIONS

Let  $x \in A$  be a nonzero divisor with  $\langle x \rangle / \langle x \rangle^2$  free over  $B = A / \langle x \rangle$  and  $(\psi, \phi)$  a matrix factorization of  $x$ .

# FROM MATRIX FACTORIZATIONS TO FREE RESOLUTIONS

Let  $x \in A$  be a nonzero divisor with  $\langle x \rangle / \langle x \rangle^2$  free over  $B = A / \langle x \rangle$  and  $(\psi, \phi)$  a matrix factorization of  $x$ .

- We can construct a minimal  $B$ -free resolution  $\mathbf{F}(\psi, \phi)$  of  $\text{coker} \phi$  with differentials  $\phi, \psi$ .

$$\mathbf{F}(\psi, \phi) : \dots \xrightarrow{\psi} F \xrightarrow{\phi} G \xrightarrow{\psi} F \xrightarrow{\phi} G \rightarrow 0$$

# FROM MATRIX FACTORIZATIONS TO FREE RESOLUTIONS

Let  $x \in A$  be a nonzero divisor with  $\langle x \rangle / \langle x \rangle^2$  free over  $B = A / \langle x \rangle$  and  $(\psi, \phi)$  a matrix factorization of  $x$ .

- We can construct a minimal  $B$ -free resolution  $\mathbf{F}(\psi, \phi)$  of  $\text{coker} \phi$  with differentials  $\phi, \psi$ .

$$\mathbf{F}(\psi, \phi) : \dots \xrightarrow{\psi} F \xrightarrow{\phi} G \xrightarrow{\psi} F \xrightarrow{\phi} G \rightarrow 0$$

- In such a resolution, necessarily,  $\text{rank}_B G = \text{rank}_B F$  (5.3-5.4)



# FROM MATRIX FACTORIZATIONS TO FREE RESOLUTIONS

Let  $x \in A$  be a nonzero divisor with  $\langle x \rangle / \langle x \rangle^2$  free over  $B = A / \langle x \rangle$  and  $(\psi, \phi)$  a matrix factorization of  $x$ .

- We can construct a minimal  $B$ -free resolution  $\mathbf{F}(\psi, \phi)$  of  $\text{coker} \phi$  with differentials  $\phi, \psi$ .

$$\mathbf{F}(\psi, \phi) : \dots \xrightarrow{\psi} F \xrightarrow{\phi} G \xrightarrow{\psi} F \xrightarrow{\phi} G \rightarrow 0$$

- In such a resolution, necessarily,  $\text{rank}_B G = \text{rank}_B F$  (5.3-5.4)
- If  $\det_A \phi = x^k u$  with  $u \notin \langle x \rangle$ , then  $\text{rank}_B \text{coker} \phi = k$ . (5.6)

# So What?

# SO WHAT?

- These classify maximal CM modules over complete hypersurfaces!

# SO WHAT?

- These classify maximal CM modules over complete hypersurfaces!

## THEOREM (6.1)

*Let  $x \in A$ ,  $d = \dim A$ , and  $B := A/\langle x \rangle$ . Then,*

# SO WHAT?

- These classify maximal CM modules over complete hypersurfaces!

## THEOREM (6.1)

Let  $x \in A$ ,  $d = \dim A$ , and  $B := A/\langle x \rangle$ . Then,

- 1 For any  $B$ -module  $M$  with minimal free resolution  $\mathbf{F}$ , the truncation at  $F_{d+1}$  is periodic with period 2.

# SO WHAT?

- These classify maximal CM modules over complete hypersurfaces!

## THEOREM (6.1)

Let  $x \in A$ ,  $d = \dim A$ , and  $B := A/\langle x \rangle$ . Then,

- 1 For any  $B$ -module  $M$  with minimal free resolution  $\mathbf{F}$ , the truncation at  $F_{d+1}$  is periodic with period 2.
- 2  $\mathbf{F}$  periodic  $\iff M$  is a maximal CM module w/o a free summand.

# SO WHAT?

- These classify maximal CM modules over complete hypersurfaces!

## THEOREM (6.1)

Let  $x \in A$ ,  $d = \dim A$ , and  $B := A/\langle x \rangle$ . Then,

- 1 For any  $B$ -module  $M$  with minimal free resolution  $\mathbf{F}$ , the truncation at  $F_{d+1}$  is periodic with period 2.
- 2  $\mathbf{F}$  periodic  $\iff M$  is a maximal CM module w/o a free summand.
- 3 If so,  $\mathbf{F}$  is induced by a matrix factorization.

# SKETCH OF PROOF OF THEOREM 6.1

- $(2) \implies (1)$



# SKETCH OF PROOF OF THEOREM 6.1

- (2)  $\implies$  (1)

## LEMMA (0.1–WELL-KNOWN)

*For  $B$  a local ring and  $B$ -free resolution*

$$\mathbf{F} : \dots \xrightarrow{\partial_3} F_2 \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} F_0 \xrightarrow{\partial_0} 0$$

- 1 If  $x_1, \dots, x_r$  is a  $B$ -sequence, then it is a regular sequence on  $\operatorname{Im} \partial_k$  for all  $k \geq r$ .
- 2 If  $\mathbf{F}$  is minimal,  $\operatorname{Im} \partial_k$  has no free summands for  $k \geq 1 + \operatorname{depth} B$

# SKETCH OF PROOF OF THEOREM 6.1

- $(2) \implies (1)$

## LEMMA (0.1–WELL-KNOWN)

*For  $B$  a local ring and  $B$ -free resolution*

$$\mathbf{F} : \dots \xrightarrow{\partial_3} F_2 \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} F_0 \xrightarrow{\partial_0} 0$$

- 1 If  $x_1, \dots, x_r$  is a  $B$ -sequence, then it is a regular sequence on  $\operatorname{Im} \partial_k$  for all  $k \geq r$ .
  - 2 If  $\mathbf{F}$  is minimal,  $\operatorname{Im} \partial_k$  has no free summands for  $k \geq 1 + \operatorname{depth} B$
- $(2 \implies )$  Note that  $M = \operatorname{coker}(G \rightarrow F)$ .

# SKETCH OF PROOF OF THEOREM 6.1

- (2  $\Leftarrow$  ) Consider  $M$  as an  $A$ -module. Use  $\mathbf{F}^\# : 0 \rightarrow F \rightarrow G \rightarrow M$  be its minimal  $A$ -free resolution.

# SKETCH OF PROOF OF THEOREM 6.1

- (2  $\Leftarrow$  ) Consider  $M$  as an  $A$ -module. Use  $\mathbf{F}^\# : 0 \rightarrow F \rightarrow G \rightarrow M$  be its minimal  $A$ -free resolution.
- Show  $F \rightarrow G$  satisfies conditions for matrix factorization with corresponding  $\psi$ .

# SKETCH OF PROOF OF THEOREM 6.1

- (2  $\Leftarrow$  ) Consider  $M$  as an  $A$ -module. Use  $\mathbf{F}^\# : 0 \rightarrow F \rightarrow G \rightarrow M$  be its minimal  $A$ -free resolution.
- Show  $F \rightarrow G$  satisfies conditions for matrix factorization with corresponding  $\psi$ .
- Show if  $\mathbf{F}(\phi, \psi)$  is not minimal,  $M$  contains a free summand.

# SKETCH OF PROOF OF THEOREM 6.1

- (3)

# SKETCH OF PROOF OF THEOREM 6.1

- (3)
- Show that for any periodic resolution  $\mathbf{F}$  over  $B$ ,  $t(A, x, \mathbf{F})$  can be chosen to be the identity!

# SO WHAT?

- Expanding upon a familiar characterization!

## THEOREM (AUSLANDER-BUCHSBAUM-SERRE)

*For  $(R, \mathfrak{m})$  local,  $R$  is regular  $\iff \text{gl dim}(R) < \infty$ .*



# SO WHAT?

- Expanding upon a familiar characterization!

## THEOREM (6.1)

*For  $(R, \mathfrak{m})$  local with  $\dim R = d$ , TFAE:*

- *For some  $x_1, \dots, x_{d+1} \in R$ ,  $\mathfrak{m} = \langle x_1, \dots, x_{d+1} \rangle$  and  $\hat{O}$  is unmixed in  $\hat{R}^{\mathfrak{m}}$  (i.e. all associated primes of  $\hat{O}$  are minimal).*
- *For any f.g.  $R$ -module  $M$  with minimal free resolution  $\mathbf{F}$ , the truncation of  $\mathbf{F}$  at degree  $d + 1$  is periodic of period 2.*
- *There exists a free resolution  $\mathbf{F} : \dots \rightarrow F_1 \rightarrow F_0 \rightarrow 0$  of  $R_{\mathfrak{m}}/\mathfrak{m}R_{\mathfrak{m}}$  where for some  $n$ ,  $\text{rank} F_n < n$ .*

*We call such an  $R$  an abstract hypersurface.*

# SO WHAT?

- Expanding upon a familiar characterization!

## THEOREM (6.1)

*For  $(R, \mathfrak{m})$  local with  $\dim R = d$ , TFAE:*

- *For some  $x_1, \dots, x_{d+1} \in R$ ,  $\mathfrak{m} = \langle x_1, \dots, x_{d+1} \rangle$  and  $\hat{O}$  is unmixed in  $\hat{R}^{\mathfrak{m}}$  (i.e. all associated primes of  $\hat{O}$  are minimal).*
- *For any f.g.  $R$ -module  $M$  with minimal free resolution  $\mathbf{F}$ , the truncation of  $\mathbf{F}$  at degree  $d + 1$  is periodic of period 2.*
- *There exists a free resolution  $\mathbf{F} : \dots \rightarrow F_1 \rightarrow F_0 \rightarrow 0$  of  $R_{\mathfrak{m}}/\mathfrak{m}R_{\mathfrak{m}}$  where for some  $n$ ,  $\text{rank} F_n < n$ .*

*We call such an  $R$  an abstract hypersurface.*

# Thank you for listening!